



ESTEEM ALL IN™

Superior control of hard-to-kill broadleaf weeds like thistles, dandelions and hemp-nettle – all in a convenient formulation with flexible use rates!



GROUP 4

Active Ingredients

Fluroxypyr 59.7 g/L +
Clopyralid 40 g/L +
MCPA 2 EH Ester 240.5 g/L
= EC

Packaging

Case: 2 × 10.93 L jugs

Application Rates and Acres Treated

Low rate: 730 ml/ac

High rate: 970 ml/ac

Acres Treated: 11–15 ac/jug

Water Volume

Ground: 20–40 L/ac
(5–10 US gal/ac)

Aerial: 12–20 L/ac
(3–5 US gal/ac)

Rainfastness

4 hours

REGISTERED CROPS

- Barley
- Oats
- Wheat (spring, winter)

KEY WEEDS CONTROLLED*

At the low rate of 730 ml/ac:

- Burdock
- Canada thistle (low infestations)
- Cleavers
- Cocklebur
- Field Horsetail (top growth)
- Flixweed
- Kochia
- Lamb's quarters
- Plantain (top growth)
- Prickly lettuce
- Ragweed
- Shepherd's purse
- Stinkweed
- Stork's bill
- Wild buckwheat
- Wild mustard
- Vetch

At the high rate of 970 ml/ac, the above weeds plus:

- Annual sow thistle
- Canada thistle (medium to high infestations, season-long control)
- Chickweed
- Common Groundsel
- Dandelion (spring rosettes only)
- Hemp-nettle
- Perennial sow thistle (season-long control)
- Redroot pigweed
- Russian Pigweed
- Scentless Chamomile
- Smartweed
- Tartary buckwheat
- Volunteer canola

* for complete list of weeds controlled, refer to the label.



ESTEEM ALL IN™

HOW IT WORKS

ESTEEM ALL IN™ moves within the plant to control exposed and underground plant tissues. It mimics naturally occurring plant hormones which control weeds by disrupting normal plant growth patterns. Symptoms of effect include epinasty (twisting of the stems) and swollen nodes.

CROP STAGING

Spring wheat, barley and oats: 3-leaf to just before the flag leaf

Winter wheat (in the spring): 3-tiller stage to just before the flag leaf

SUPPORTED TANK MIXES

- Liquid Achieve™ SC¹
- BISON® 400 L
- Puma® Advance
- Traxos®
- Simplicity™ GoDRI™²
- Varro®

¹Tank mixes with Liquid Achieve™ SC may cause temporary injury if applied before the 4-leaf stage, however, yield will not normally be affected.

²Use only the 730 ml/acre (low) rate when mixing with Simplicity™ GoDRI™.

This product may be tank-mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank-mix partner labels, the most restrictive label must be followed. Do not tank-mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

QUICK TIPS

Flexible rates are available — choose what works best for you:

- 970 ml/ac for heavy weed pressure OR hard-to-control weeds;
- 730 ml/ac for standard weed pressure.

MIXING INSTRUCTIONS

1. Fill sprayer tank ½ full of water.
2. Start sprayer tank agitation.
3. Add any dry formulation tank mix partners.
4. Add the required amount of grassy weed tank-mix partner.
5. Add the required amount of ESTEEM ALL IN™.
6. Add any required adjuvants of surfactants for tank-mix partners.
7. Complete filling tank with water.

PRE-HARVEST INTERVALS

- Forage: 7 days
- Wheat, barley, oats: 60 days

ADJUVANT RATE

Adjuvant only required, if tank-mixing with Liquid Achieve™ SC: Turbocharge® @ 0.5% v/v

CROP ROTATIONS

Fields previously treated with ESTEEM ALL IN™ can be:

- seeded to wheat, barley, oats and rye (not underseeded to forage legumes, clover or alfalfa), canola, field peas, flax, forage grasses, mustard, or;
- summer-fallowed.

Field peas: a rainfall of 140 mm (5.5") between herbicide application and August 31 (in the year of application), as well as an annual precipitation greater than 175 mm (6.9") is required.

GRAZING RESTRICTIONS

Do not cut or graze treated fields of wheat, barley or oats within 7 days after application.

Withdraw meat animals from treated fields at least 3 days before slaughter.

STORAGE

Do not freeze.

WANT TO KNOW MORE?

Click on the QR code to access all the product details including the label and SDS documents



ADAMA

Always read and follow label directions.
Toll-free: 1.855.264.6262
Website: ADAMA.COM/CANADA

®/™ BISON® is a registered trademark and ESTEEM ALL IN™ is a trademark of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies. © 2024 ADAMA Agricultural Solutions Canada Ltd.
04/24