




INSECTICIDE



ADAMA

CORMORAN®

Multiple modes of action for codling moth control in apples and Colorado potato beetle control in potatoes as well as a wide range of other insects in specialty crops.



GROUP 4 & 15

Active Ingredients:

Novaluron 100 g/L and Acetamiprid 80 g/L = EC

Application Rates and Acres Treated:

Rate: 180–840 ml/ac

Acres Treated: 12–56 ac/jug

Packaging:

Case: 2 x 10.08 L jugs

Water Volume:

Ground: 80–400 L/ac
(20–105 US gal/ac)

Aerial: Do not apply by air.

Rainfastness:

Avoid application when heavy rain is forecast.

REGISTERED CROPS:

- Alfalfa (grown for seed)
- Apples
- Brassica Leafy Greens (crop sub-group 4-13B)
- Bushberries (crop sub-group 13-07B)
- Head and Stem Brassica Vegetable Crops (crop group 5-13)
- Peppers (bell and non-bell)
- Potatoes
- Strawberries
- Stone Fruits (crop group 12-09)
- Sweet Corn

KEY INSECTS CONTROLLED:

- Alfalfa Looper
- Alfalfa Plant Bug
- Aphids
- Apple Maggot
- Armyworm
- Blueberry Flea Beetle
- Blueberry Gall Midge (cranberry tip worm)
- Blueberry Maggot Fly
- Blueberry Spanworm
- Cabbage Looper
- Cherry Fruit Fly (suppression)
- Leafhopper
- Lygus Bug
- Cherry Fruitworm
- Codling Moth
- Colorado Potato Beetle
- Cranberry Fruitworm
- Diamondback Moth
- Dogwood Borer
- European Apple Sawfly
- European Corn Borer
- Green Fruitworm
- Gypsy Moth
- Imported Cabbageworm
- Japanese Beetle
- Leafhopper
- Leafroller
- Lesser Appleworm
- Lygus Bug
- Mullein Bug
- Oriental Fruit Moth
- Plum Curculio
- Spotted Wing Drosophila
- Strawberry Clipper Weevil
- Strawberry Rootworm
- Swede Midge
- Tarnished Plant Bug
- Tentiform Leafminer
- Thrips

HOW IT WORKS:

CORMORAN® kills insect eggs by contact and larvae by ingestion. Containing two modes of action, CORMORAN® provides both rapid knockdown and residual control of insect pests.

REGISTERED AND SUPPORTED TANK MIXES:

- Acramite® 50 WS
- Luna Sensation™
- Polyram® DF WSP
- Pristine® WG

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

CLICK HERE FOR FULL PRODUCT DETAILS.





CORMORAN[®]

CROP STAGING AND RATES:

The need and timing of an application should be based on the presence of pests at vulnerable developmental stages and significant populations, as determined by local monitoring. Consult the label for specific crop and insect timing.

MIXING INSTRUCTIONS:

1. Fill sprayer tank ½ full of water.
2. Start sprayer tank agitation.
3. Add the required amount of CORMORAN[®].
4. If necessary, add any required adjuvants or surfactants for tank-mix partners.
5. Complete filling tank with water.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

RE-ENTRY INTERVALS:

Crop	Re-entry activities	Restricted entry interval
Alfalfa (grown for seed)	All activities	12 hours
Apples	Fruit thinning	7 days
	All other activities	12 hours
Bushberries (Crop sub-group 13-07B)	All activities	12 hours
Stone Fruit	Fruit thinning	6 days
	All other activities	12 hours
Strawberries	All activities	12 hours
Brassica (cole) leafy vegetables (Crop Group 5-13)	Hand weeding, scouting, tying and training	2 days
	All other activities	12 hours
Leafy vegetables-Brassica leafy greens (Crop sub-group 4-13B)	All activities	12 hours
Peppers (bell and non-bell)		
Potatoes		
Sweet Corn		

PRE-HARVEST INTERVALS:

- Strawberries: 1 day
- Brassica leafy vegetables (crop group 5 and subgroup 4), Peppers, Potatoes, Stone Fruits: 7 days
- Bushberries (crop group 13-07B): 8 days
- Sweet Corn: 10 days
- Alfalfa (grown for seed), Apples: 14 days

STORAGE:

- Store in original, tightly closed container.
- Do not ship or store near food, feed, seed and fertilizers.
- Store in cool, dry, locked, well-ventilated area without floor drain.
- Keep away from fire or open flame, or other sources of heat.

CROP ROTATIONS:

No restrictions.

QUICK TIPS:

Consider early applications (before petal fall) of CORMORAN[®] to allow beneficial insects to build up later in the season. To minimize the possibility of transient effects on honeybee brood development, do not use CORMORAN[®] on blooming crops when bees are actively foraging. If orchards have been historically infested with mites or aphids, be sure to scout regularly and use miticides to control their populations.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

CORMORAN[®] is a registered trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd.