

Active Ingredients:

Imazamox 80 g/L = SL Clethodim 120 g/L = EC

Application Rates and Acres Treated:

Rate: DAVAI[®] 80 SL: 100 ml/ac ARROW ALL IN[®]: 150 ml/ac

Acres Treated: 40 ac/case

Packaging:

Co-pack includes: 1 x 4 L jug of DAVAI[®] 80 SL 1 x 6 L jug of ARROW ALL IN[®]

Water Volume:

Ground: 40 L/ac (10 US gal/ac) **Aerial:** Do not apply by air.

Rainfastness:

3 hours

DAVAI[®] A PLUS DAVAI[®] 80 SL, is conveniently packaged with

ADAMA

ARROW ALL IN[®] to offer broad-spectrum control in dry beans, peas and soybeans.

GROUP 1 & 2

REGISTERED CROPS:

- Dry Beans
- Field Peas
- Soybeans

Stork's Bill

WEEDS CONTROLLED AND APPLICATION TIMING:

BROADLEAF WEEDS

Cotyledon - 4 leaf:

- Cleavers¹
- Cow Cockle
- Flixweed
- Green Smartweed

GRASSY WEEDS 1-6 true leaf:

- Barnyard Grass
- Crabgrass (smooth, large)³
- Fall Panicum
- Green Foxtail²
- Japanese Brome Grass¹

- Lamb's-Quarters
- Redroot Pigweed
- Shepherd's Purse
- Stinkweed
- Persian Darnel
- Proso Millet
- Quackgrass¹
- Volunteer Canary Grass
- Volunteer Cereals (barley, oats, wheat)

Volunteer Canola⁵

Wild Buckwheat¹

Wild Mustard

- Volunteer Corn³
- Wild Oats²
- Witchgrass
- Volunteer Canary Seed⁴
 Yellow Foxtail

¹Suppression ²Including Group 1-resistant weeds. DAVAI® A PLUS will NOT control weed biotypes that are multiple-resistant to both Group-1 and Group-2 herbicides.

³2- to 6-leaf

⁴1- to 4-leaf

⁵ Non imidazolinone-tolerant varieties

HOW IT WORKS:

DAVAI® A PLUS combines two (2) actives to tackle broadleaf and grassy weeds. See DAVAI® 80 SL and ARROW ALL IN® for more information.

DAVAI®A PLUS

CROP STAGING:

- Dry beans, Soybeans: Emergence to 3 expanded trifoliate leaves
- Field peas: 1-6 true leaf

MIXING INSTRUCTIONS:

- Fill clean tank ½ to ¾ full of clean water and turn agitation on.
- 2. Start sprayer tank agitation.
- 3. Add the required amount of DAVAI® 80 SL herbicide and continue to agitate.
- 4. Add the required amount of ARROW ALL IN[®] herbicide and continue to agitate.
- 5. If necessary, add any required adjuvants or surfactants for tank-mix partners.
- 6. Complete filling the tank with water as agitation continues.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

RECROPPING RESTRICTIONS:

There are several factors that effect the re-cropping following an Imidazolinone application. These include in order of importance:

- 1. Product: Imazethapyr, for example, is more persistent than imazamox.
- 2. Soil moisture: Requires more than 125 mm (5") of rain between herbicide application and August 31 in the year of application.
- **3. Organic matter:** Brown soil zones (< 3% organic matter) are more susceptible to carryover crop injury the year after application.
- 4. Rate: Depending on the crop and rates, soil residues can be an issue.
- 5. Soil pH: Product persists longer in a pH < 5.5-6.

Depending on the following crop the level of sensitivity will vary. Please contact your local ADAMA Area Business Manager for more details.

CROP ROTATIONS:

(Can be seeded the following spring)

- Barley
- Canary Seed
- Canola
- Corn
- Field Peas
- Imidazolinone-Tolerant Sunflowers
- Oats
- Soybeans
- Wheat (spring)

PRE-HARVEST INTERVALS:

- Dry Beans: 75 days
- Field Peas: 60 days
- Soybeans: 85 days

GRAZING RESTRICTIONS:

- Field Peas: 30 days
- All other crops: Do not graze.

ADJUVANT RATE:

No adjuvant required.

RE-ENTRY INTERVAL:

12 hours

STORAGE:

Do not freeze.

CLICK HERE FOR FULL PRODUCT DETAILS.



QUICK TIPS:

For best results, apply DAVAI® A PLUS to actively growing weeds. If the plants are under environmental stress, expect to see temporary yellowing or burning on the leaves.



Always read and follow label directions. Toll-free: 1.855.264.6262 | Website: ADAMA.COM

ARROW ALL IN® and DAVAI® are registered trademarks of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd.