



RUSH 24 ALL IN[®]

Controls a wide spectrum of broadleaf weeds, like cleavers, with excellent grassy weed herbicide compatibility for wheat and barley.



GROUP 4

Active Ingredients:

Fluroxypyr 90 g/L and 2,4-D Ester 360 g/L as an EC

Packaging:

Case: 2 x 8.9 L jug

Bulk: 106.8 L

Tote: 427.2 L

Application Rates & Acres Treated:

Rate: 445 ml/ac

Acres Treated: 20 ac/jug;
240 ac/drum;
960 ac/tote

Water Volume:

Ground: 12-40 L/ac
(3-10 US gal/ac)

Aerial: 12-20 L/ac
(5-10 US gal/ac)

Rainfastness:

1 hour

KEY BENEFITS:

- Control of Group 2-resistant biotypes of Kochia
- Suppression of Group 2-resistant biotypes of Chickweed
- Convenient and trusted formulation
- Easily tankmixes with the most popular grassy weed control herbicides
- Available in convenient bulk options

REGISTERED CROPS:

- Barley
- Wheat (spring, winter)

WEEDS CONTROLLED:

Control		Suppression
At labelled rate of 445 ml/ac		
<ul style="list-style-type: none"> • Bluebur • Burdock • Canola (Volunteer) • Cleavers (1-8 whorls) • Clovers (Sweet) • Cocklebur • Flixweed • Field Horsetail¹ • Goat's-Beard • Hemp-Nettle (2- to 6-leaf stage) • Hoary Cress¹ • Kochia² • Lamb's-Quarters 	<ul style="list-style-type: none"> • Mustards (except Green Tansy, Dog & Grey Tansy) • Plantain • Prickly Lettuce • Ragweed • Round-Leaved Mallow (1- to 6-leaf) • Shepherd's Purse • Stork's Bill (1- to 8-leaf) • Stinkweed • Sunflower (Annual) • Vetch • Wild Buckwheat (1-6 leaf) • Wild Radish • Volunteer Flax (1-12 cm) 	<ul style="list-style-type: none"> • Common Chickweed² (up to 8 cm) • Redroot Pigweed • Sow Thistle (Perennial)¹
Additional weeds controlled/suppressed with addition of 81 ml/ac (2 oz/ac) of 2,4-D ESTER 700:		
<ul style="list-style-type: none"> • Blue Lettuce¹ • Dandelion (Spring Rosettes) • Docks • Field Bindweed¹ • Field Peppergrass • Gumweed • Hairy Galinsoga • Hedge Bindweed 	<ul style="list-style-type: none"> • Lady's Thumb • Leafy Spurge¹ • Mustard (Dog, Tansy) • Oak-Leaved Goosefoot • Redroot Pigweed • Russian Thistle • Smartweed • Tartary Buckwheat 	<ul style="list-style-type: none"> • Common Chickweed² (up to 8 cm) • Canada Thistle¹ • Sow Thistle (Perennial¹, Annual)

¹Top growth control only

²Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

RUSH 24 ALL IN®

HOW IT WORKS:

Group 4 herbicides disrupt normal plant growth, resulting in twisting and cupping of leaves, epinasty and death of susceptible plants in 2–10 days.

CROP STAGING:

4-leaf to just prior to flag leaf emergence.

REGISTERED AND SUPPORTED TANK MIXES:

Wheat only:

- Simplicity™ GoDRI™¹
- Traxos®

Wheat and Barley:

- Puma® Advance

¹ Additional 2,4-D Ester is not recommended when mixing RUSH 24 ALL IN® and Simplicity™ GoDRI™.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

1. Fill sprayer tank ½ full of water.
2. Start sprayer tank agitation.
3. Add the required amount of RUSH 24 ALL IN®.
4. If necessary, add the required amount of additional 2,4-D Ester 700 herbicide.
5. If necessary, add any required adjuvants of surfactants for tank-mix partners.
6. Complete filling tank with water with agitation on.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

CROP ROTATIONS:

The following crops may be grown one (1) year after application:

- | | | |
|----------------|--------------|---------|
| • Barley | • Oats | • Rye |
| • Canola | • Field Peas | • Wheat |
| • Forage Grass | | |

GRAZING RESTRICTIONS:

- Do not permit lactating dairy animals to graze fields within seven (7) days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least three (3) days before slaughter.

PRE-HARVEST INTERVAL:

Barley, Wheat (spring): 60 days

RE-ENTRY INTERVAL:

12 hours

STORAGE:

Do not freeze.

[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



QUICK TIPS:

RUSH 24 ALL IN® activity is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, for example drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

RUSH ALL IN® are registered trademarks of an ADAMA Group Company.
All other products are trademarks of their respective companies.
© 2025 ADAMA Agricultural Solutions Canada Ltd.