



Active Ingredients:

Fluroxypyr 80 g/L Bromoxynil 200 g/L MCPA 200 g/L

Application Rates and Acres Treated:

Rate: 567 ml/ac

Acres Treated: 20 ac/jug

Packaging:

Case: 2 x 11.37 L jugs

Water Volume:

Ground: 20 – 40 L/ac (5 – 10 US gal/ac)

Aeriel: Do not apply by air.

Rainfastness:

1 hour

REGISTERED CROPS:

Barley

• Wheat (spring)

Cocklebur

WEEDS CONTROLLED:

Top growth control only:

- American Nightshade
- Canada Thistle

Up to 4-leaf:

- Bluebur
- Cow Cockle
- Flixweed
- Lady's Thumb
- Night-Flowering Catchfly

Up to 8-leaf:

- Common Groundsel
- Lamb's Quarters
- Stinkweed

- Perennial Sow Thistle
- Redroot Pigweed
- Shepherd's Purse
- Smartweed (green, pale)
- Volunteer Canola
- Volunteer Sunflower
- Wild Buckwheat
 - Wild Mustard¹
 - Wormseed Mustard

Other:

- Cleavers (up to 4 whorls)¹
- Kochia (up to 5 cm)²
- Russian Thistle (up to 5 cm)
- Wild Radish

¹Including Group 2-resistant biotypes

² Including Group 2- and glyphosate-resistant biotypes



FORCEFIGHTER ALL IN®

HOW IT WORKS:

Quickly causes plants to stop growing. This convenient all in one formulation controls a wide range of weeds, including glyphosate-resistant and Group 2 resistant Kochia, Group 2 resistant Cleavers and Group 2 resistant Wild Mustard.

CROP STAGING:

2-leaf to early flag leaf

REGISTERED AND SUPPORTED TANK MIXES:

Spring wheat:

Traxos[®] Herbicide

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

- 1. Fill spray tank ½ full with water.
- 2. Start sprayer tank agitation.
- 3. Add the required amount of FORCEFIGHTER ALL IN®
- 4. If necessary, add any required adjuvants or surfactants for tank-mix partners.
- 5. Fill the tank and agitate again before use.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

CROP ROTATIONS:

Can be seeded the following year to barley, canola, flax, forage grasses, lentils, mustard, oats, peas, rye and wheat or fields can be summerfallowed.

PRE-HARVEST INTERVAL:

60 days

GRAZING RESTRICTIONS:

30 days

RE-ENTRY INTERVAL:

24 hours

STORAGE:

SCAN THE QR CODE





Do not freeze.



Do not apply before the 2-leaf stage as crop

Use 40 L/ac application volume when there is a heavy canopy or when most weeds are at an advanced stage of growth.

Activity is influenced by weather conditions. Optimal application temperature is 12 – 24° C. Avoid application 3 days before or after frost.



Always read and follow label directions. Toll-free:1.855.264.6262 | Website: ADAMA.COM