



ADAMA

BROMOTRIL® 240 EC

Tough broadleaf weed control with tank-mix flexibility and excellent crop safety.



GROUP 6

Active Ingredient:

Bromoxynil Octanoate Ester 240 g/L = EC

Application Rates:

Pre-plant: 490 – 590 ml/ac

In-crop broadleaf: 490 – 570 ml/ac

Packaging and Acres Treated:

Case: 2 × 9.7 L jugs

Drum: 116.4 L

Acres Treated:

Pre-plant: 20 ac/jug;

200 – 240 ac/drum

In-crop broadleaf: 17 – 20 ac/jug;

200 – 240 ac/drum

Water Volume:

Ground: 20 – 80 L/ac

(5 – 20 US gal/ac)

Aerial (in-crop only)*: 8 – 16 L/ac

(3 – 5 US gal/ac)

*Wheat and Barley only

Rainfastness:

30 minutes

REGISTERED CROPS:

Crop	Timing
Barley, Canola, Oats, Wheat	Pre-plant burn-off with glyphosate

Crop	Crop Stage
Alfalfa (seedling)	2 – 6 trifoliolate
Alfalfa (established for seed production only)	Spring: before the crop begins to shield the weeds
Barley, Oats, Triticale, Wheat (spring)	2 leaf to early flag
Winter Wheat	Fall: 2 – 4 leaf Spring: first growth to early flag
Corn (field, sweet)	4 – 8 leaf (beyond 8-leaf requires drop pipes)
Fall Rye	Spring: from first growth to early flag
Forage Millet, Sorghum	4 leaf to 8 inches

WEEDS CONTROLLED:

Seedling up to 4-leaf stage:

- American Nightshade
- Bluebur
- Cocklebur
- Common Ragweed
- Cow Cockle¹
- Green Smartweed
- Kochia²
- Lady's Thumb
- Pale Smartweed
- Pigweed¹
- Russian Thistle²
- Stinkweed¹
- Velvetleaf³
- Wild Mustard¹

Seedling up to 8-leaf stage:

- Common Buckwheat
- Common Groundsel
- Lamb's Quarters
- Tartary Buckwheat
- Wild Buckwheat

¹ In normal conditions, it will be controlled up to 4-leaf stage. Plants beyond this stage are unlikely to be controlled; the higher rate generally gives better results.

² Spray before plants are 2 inches high.

³ Spray before plants are 3 inches high.



BROMOTRIL® 240 EC

HOW IT WORKS:

BROMOTRIL® 240 EC is a contact herbicide which controls Group 2 and Group 9 resistant biotypes. Leaves will yellow in 2–4 days with complete control in 7–14 days.

REGISTERED AND SUPPORTED POST-EMERGENT TANK MIXES:

Post-emergent herbicides:

- Barley, Wheat (spring, winter): 2,4-D Ester 700, BISON® 400 L MCPA Ester 600, Liquid Achieve™ SC
- Corn: Accent® 75 DF, Atrazine, Banvel® (dicamba), Banvel® II
- Oats, Fall Rye: MCPA Ester 600

Pre-plant herbicide:

- Glyphosate

Fungicide:

- BUMPER® 432 EC

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

1. Fill spray tank ½ full with water.
2. Add required amount of BROMOTRIL® 240 EC. Begin agitation.
3. If tank mixing, add any tank-mix partners to the spray tank first, agitate and then add BROMOTRIL® 240 EC (unless otherwise directed by the BROMOTRIL® 240 EC and tank-mix partner label).
4. Add the remaining amount of water while agitation continues.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

CROP ROTATIONS:

No restrictions.

PRE-HARVEST INTERVAL:

30 days

STORAGE:

Do not freeze.

RE-ENTRY INTERVAL:

24 hours

POST-EMERGENT GRAZING RESTRICTIONS:

- Do not use treated crops for grazing of livestock or green feed until 30 days after application.
- Do not cut treated crops for forage until 30 days after application.

[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



QUICK TIPS:

Avoid spraying if temperatures are above 25° C. Leaf scorching may occur in corn and flax if applied during or after adverse growing conditions, such as cool and wet or hot (above 27° C) and humid weather. For best results, spray when weeds are in the seedling stage.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

BISON®, BROMOTRIL®, and BUMPER® are registered trademarks of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2026 ADAMA Agricultural Solutions Canada Ltd.