



ADAMA

MAXENTIS®

Provides protection against White Mould in soybeans and Rust in lowbush blueberries. Features two distinct elite modes of action (Groups 3 & 11) for resistance management and long lasting systemic activity.



POWERED BY

Asorbital®

FORMULATION TECHNOLOGY

GROUP 3 & 11

Active Ingredients:

Azoxystrobin 120 g/L,
Prothioconazole 90 g/L = EC

Application Rates:

Rate: 253 – 422 ml/ac

Packaging and Acres Treated:

Case: 2 x 8.45 L jugs
Drum: 118.1 L

Acres Treated: 40 – 60 ac/case;
280 – 467 ac/drum

Water Volume:

Ground: 80 L/ac (20 US gal/ac)
Aerial: 20 L/ac (5 US gal/ac)

Rainfastness:

Avoid applying when rain is forecast.

KEY BENEFITS:

- Unique combination of prothioconazole and azoxystrobin
- Enhanced EC formulation – with bulk packaging capabilities
- Multi-mode activity for resistance management (Groups 3 & 11)
- Broad-spectrum disease control with protective and curative action
- Whole plant protection: translaminar and systemic movement
- Wide window of application

REGISTERED CROPS*:

- Soybeans
- Lowbush Blueberries

KEY DISEASES CONTROLLED*:

- Leaf Rust (suppression)
- White Mould (*Sclerotinia*)

*See label for the complete list of registered crops as well diseases controlled and rates for each crop.

HOW IT WORKS:

Two powerful active ingredients, combined for the first time in Canada, that provide protective and curative action on a wide range of diseases, across multiple crops.



MAXENTIS®

APPLICATION TIMING AND CROP STAGING:

Crop	Diseases	Rate	Timing
Lowbush Blueberries	Rust (suppression)	253 ml/ac	Apply at the first sign of disease in the sprout year. After initial application, one additional application may be made 1–14 days afterwards if conditions remain favourable for continued or increased disease development. Maximum applications per year: 2
Soybeans	White Mould (<i>Sclerotinia</i>)	422 ml/ac	Apply as a preventive foliar spray at the beginning of flowering or at first sign of disease. After the initial application, one additional application may be made 10–14 days afterwards if conditions remain favourable for continued or increased disease development. Apply the higher rate when conditions favour disease development, or when growing less disease resistant varieties. Maximum applications per year: 2

MIXING INSTRUCTIONS:

1. Fill sprayer tank $\frac{3}{4}$ full of water.
2. Start sprayer tank agitation.
3. Add the required amount of MAXENTIS®.
4. If applicable, add any required adjuvants or surfactants for tank-mix partners.
5. Complete filling tank with water.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

GRAZING RESTRICTIONS:

Forage, hay: 30 days

Grazing or green feed: 6 days

CROP ROTATIONS:

No restrictions.

PRE-HARVEST INTERVAL:

- Soybeans: 20 days
- Lowbush Blueberries: 30 days

NOTE: Refer to the label for pre-harvest intervals for all other registered crops.

REGISTERED AND SUPPORTED TANK MIXES:

None registered.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

RE-ENTRY INTERVAL:

24 hours

STORAGE:

Do not freeze.

[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



QUICK TIPS:

Environmental conditions are a major driver of disease pressure and severity. Under high disease pressure conditions, a rate of MAXENTIS® at the upper end of the rate range is recommended. When temperatures are above 25° C we recommend application in the evening.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

ASORBITAL® and MAXENTIS® are registered trademarks of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2026 ADAMA Agricultural Solutions Canada Ltd.