


NEW

SORADUO™ ALL IN

Get two proven modes of action for Fusarium and broad-spectrum cereal leaf diseases. Now in a convenient formulation and bulk packaging.

GROUP 3

Active Ingredients:

Prothioconazole 125 g/L
 Tebuconazole 125 g/L
 EC (Emulsifiable Concentrate)

Application Rates:

Rate: 324 ml/ac

Packaging and Acres Treated:

Drum: 104.0 L
 Acres Treated: 320 ac/drum

Water Volume:

Ground: 40 L/ac
 Aerial: 20 L/ac

Rainfastness:

Avoid application when heavy rain is forecast.

REGISTERED CROPS:

- Wheat (spring, winter)
- Barley
- Triticale (spring, winter)

DISEASES CONTROLLED:

- Fusarium Head Blight (suppression)
- Rust (leaf, stem and stripe)
- Leaf & Glume Blotch
- Tan Spot
- Powdery Mildew
- Net Blotch
- Scald
- Spot Blotch

HOW IT WORKS:

SORADUO™ ALL IN is a combination of a triazole fungicides which have broad-spectrum system activity plus long-lasting foliar protection.

CROP STAGING:

Crop	Disease	Rate	Timing
Wheat (spring, winter)	Suppression of: Fusarium Head Blight Control of: Rust (leaf, stem and stripe), Leaf & Glume Blotch, Tan Spot, Powdery Mildew	324 ml/ac	75% of the wheat heads on the main stem are fully emerged to when 50% of the heads on the main stem are in flower.
Barley	Suppression of: Fusarium Head Blight Control of: Net Blotch, Spot Blotch, Scald, Leaf Blotch, Leaf Stem and Stripe Rusts, Powdery Mildew	324 ml/ac	70 to 100% of the barley main stem heads are fully emerged, to 3 days after full head emergence
Triticale (spring and winter)	Suppression of: Fusarium Head Blight Control of: Rusts, Leaf & Glume Blotch, Tan Spot, Powdery Mildew, Spot Blotch	324 ml/ac	75% of the heads on the main stem are fully emerged to when 50% of the heads on the main stem are in flower



SORADUO™ ALL IN

REGISTERED AND SUPPORTED TANK MIXES:

None

MIXING INSTRUCTIONS:

1. When mixing SORADUO™ ALL IN alone add ½ of the required amount of water to the spray or mixing tank and start agitation.
2. Add the required quantity of SORADUO™ ALL IN to the water and complete filling with water to the required total volume.
3. Maintain agitation throughout mixing and spraying

CROP ROTATION RESTRICTIONS:

Wheat, Barley, Triticale, Soybean: None

A rotational plant-back interval of 120 days for all other crops.

RE-ENTRY INTERVAL (REI):

24 hours

PRE-HARVEST INTERVAL:

36 days

GRAZING RESTRICTIONS:

6 days

STORAGE:

Do not freeze.

Please note that there has been a re-evaluation of Tebuconazole by the PMRA; the total seasonal rate is limited to a maximum of 136 g a.i./ha/year, except for those crops where the yearly total rate is currently below 136 g a.i./ha.



[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

SORADUO™ ALL IN is a trademark of an ADAMA Group Company.
All other products are trademarks of their respective companies.
© 2026 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS:

Apply SORADUO™ ALL IN early as a preventative measure for Fusarium Head Blight as an additional benefit of controlling many key cereal leaf diseases on the flag and penultimate leaves.