




CUSTODIA®

A multi-mode of action fungicide offering preventative and curative protection of the flag leaf against Rust and other major leaf diseases.



GROUP 3 & 11

Active Ingredients:

Tebuconazole 200 g/L,
Azoxystrobin 120 g/L = SC

Application Rates:

Rate: 190 – 250 ml/ac

Packaging and Acres Treated:

Case: 2 x 10.08 L jugs

Acres Treated: 40 – 53 ac/jug

Water Volume:

Ground: 40 L/ac (10 US gal/ac)

Aerial: 20 L/ac (5 US gal/ac)

Rainfastness:

Avoid applying when heavy rainfall is in the forecast.

KEY BENEFITS:

- Multiple modes of action for resistance management
- Curative and preventative disease control
- Broad spectrum leaf disease control with superior Rust control compared to competitors
- Penetrates quickly into the leaf tissue
- Keeps the plant greener longer

KEY DISEASES CONTROLLED:

- | | |
|------------------------|---------------|
| • Leaf Rust | • Tan Spot |
| • Stem Rust | • Net Blotch |
| • Stripe Rust | • Spot Blotch |
| • Septoria Leaf Blotch | |

HOW IT WORKS:

CUSTODIA® is a broad-spectrum, preventative fungicide with systemic and curative properties recommended for the control of many important plant diseases. CUSTODIA® may be applied as a foliar spray in spray programs or in tank mixes with other crop protection products. All applications must be made according to the use directions that follow.

CLICK HERE
FOR FULL
PRODUCT
DETAILS.





CUSTODIA®

REGISTERED AND SUPPORTED TANK MIXES:

Manipulator™

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

REGISTERED CROPS AND STAGING:

Crop	Rate	Diseases	Application Timing
Wheat (spring, winter)	190 – 250 ml/ac	Leaf Rust, Stem Rust, Stripe Rust, Septoria Leaf Blotch, Tan Spot	Apply CUSTODIA® at the very early stages of disease development.
Barley		Net Blotch, Spot Blotch, Leaf Rust, Stem Rust, Stripe Rust, Septoria Leaf Blotch, Tan Spot	Use of the higher rate should be considered when weather conditions are conducive to heavy disease development or when heavy disease pressure is present.
Soybeans		Asian Soybean Rust, Frogeye Leaf Spot	
Oats	190 ml/ac	Crown Rust, Stem Rust, Septoria Leaf Blotch	Apply CUSTODIA® at the very early stages of disease development.

Please note that there has been a re-evaluation of Tebuconazole by the PMRA; the total seasonal rate is limited to a maximum of 136 g a.i./ha/year, except for those crops where the yearly total rate is currently below 136 g a.i./ha.

MIXING INSTRUCTIONS:

1. Fill sprayer tank ½ full of water.
2. Start sprayer tank agitation.
3. Add the required amount of CUSTODIA®.
4. If applicable, add any required adjuvants or surfactants for tank-mix partners.
5. Complete filling tank with water.

* Use a 50-mesh (or coarser) filter screen.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

CROP ROTATIONS:

No restrictions.

RE-ENTRY INTERVAL:

12 hours

PRE-HARVEST INTERVALS:

- Forage, hay: 6 days
- Soybeans: 20 days
- Mature grains: 36 days

GRAZING RESTRICTIONS:

Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment. Straw cut after harvest may be fed or used for bedding.

STORAGE:

Do not freeze.

QUICK TIPS:

CUSTODIA® should be applied at flag leaf for optimal leaf disease control. Pathogens coverage is key; do not use less than recommended water volumes.

For Fusarium control, we would recommend an application of SORATEL® fungicide at full head emergence after CUSTODIA® at flag leaf.

For resistance management, CUSTODIA® contains Group 3 and 11 fungicides. When possible, rotate the use of CUSTODIA® or other Group 3 and 11 fungicides with different groups that control the same pathogens.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

CUSTODIA® and SORATEL® are registered trademarks of an ADAMA Group Company. All other products are trademarks of their respective companies.
© 2026 ADAMA Agricultural Solutions Canada Ltd.