



ADAMA

NEW

EMPHASIS[®] MAX

EMPHASIS[®] MAX with its dual mode of action, has an increased rate of carfentrazone for enhanced control.



Active Ingredients:

Carfentrazone-ethyl at 240 g/L as an EC = EMPHASIS[®] A, Bromoxynil Octanoate Ester at 240 g/L as an EC = BROMOTRIL[®] 240 EC

Packaging:

Co-pack: 2 x 1.2 L EMPHASIS[®] A +
2 x 9.7 L BROMOTRIL[®] 240 EC

Water Volume:

Ground: 40 L/ac (10 US gal/ac)

Aerial: Do not apply by air.

Rainfastness:

6 hours

GROUP 6 & 14

REGISTERED CROPS:

- Wheat
- Barley
- Oats

STAGING:

Pre-seed burn-off with glyphosate.

HOW IT WORKS:

EMPHASIS[®] MAX is a multi mode contact herbicide that controls broadleaf weeds including Group 2 and 9 resistant biotypes such as Kochia. Within a few hours following application, the foliage of susceptible weeds shows signs of desiccation, and in subsequent days, necrosis and death of the plant occur.

MIXING INSTRUCTIONS:

1. Fill spray tank with ½ of the volume of clean water needed.
2. With agitator running add the required amount of EMPHASIS[®] A to spray tank.
3. Next add the required amount of BROMOTRIL[®] 240 EC.
4. Add more water, then add glyphosate.
5. Complete filling the tank to desired level.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

ADJUVANT RATE:

No adjuvant required.

CROP ROTATIONS:

No restrictions.

STORAGE:

Do not freeze.

EMPHASIS® MAX

WEEDS CONTROLLED

When used as directed, EMPHASIS® A will provide control of the listed weeds up to ten (10) cm in height, or as specified. See the individual labels of each tank-mix partner for specific rates and weed staging and always follow the directions for use of the most stringent label. Good spray coverage is essential for optimal weed control.

Weeds controlled by EMPHASIS® MAX alone	EMPHASIS® A Rate	BROMOTRIL® Rate
80 ac/case Pre-Seed Rate		
80 ac/case	30 ml/ac	236 ml/ac
Black Nightshade (up to 5 cm), Common Waterhemp, Eastern Black Nightshade (up to 5 cm), Kochia (group 2,4 & 9 resistant biotypes), Lamb’s Quarters (up to 7.5 cm), Morning Glory (up to 3 leaves), Redroot Pigweed, Tall Waterhemp (up to 5 cm), Velvetleaf, Volunteer Canola (including glyphosate-tolerant), Round-Leaved Mallow, Hairy Nightshade, Stinkweed, Prostrate Pigweed, Smooth Pigweed, Common Purslane, Smartweed (seedling), Tansy Mustard, Carpetweed, Cleavers, Cocklebur, Jimsonweed, Russian Thistle (up to 5 cm), Shepherd’s Purse, Flixweed, Tumble Pigweed		
Weeds controlled when tank mixed with glyphosate plus weeds listed above by glyphosate rate*		
Cow Cockle, Green Foxtail, Green Smartweed, Lady’s Thumb, Volunteer Barley, Volunteer Wheat, Wild Mustard, Wild Oats	0.5 REL/ac	180 g a.i./ac
Weeds Listed Above + Common Ragweed, Wild Buckwheat, Canada Fleabane**, Downy Brome, Giant Foxtail, Hempnettle, Persian Darnel, Russian Thistle, Volunteer Flax, Narrow-Leaved Hawk’s Beard	0.75 REL/ac	277 g a.i./ac
Weeds Listed Above + Annual Bluegrass, Annual Sow Thistle, Canada Thistle (rosette stage, summerfallow), Crabgrass, Dandelion (less than 15 cm), Narrow-Leaved Vetch, Prickly Lettuce, Quack Grass (light to moderate infestations, 3 – 4 green leaves or more)	1 REL/ac	360 g a.i./ac
40 ac/case Pre-Seed Rate for Wheat, Barley & Oats only		
40 ac/case	30 ml/ac	472 ml/ac
	* 1 jug of EMPHASIS® A remaining	
Weeds controlled by 80 ac EMPHASIS® MAX rate and weeds controlled at desired glyphosate rate plus: American Nightshade, Bluebur, Carpetweed, Common Buckwheat, Common Groundsel, Common Purslane, Common Waterhemp, Hairy Nightshade, Jimsonweed, Pale Smartweed, Tansy Mustard, Tartary Buckwheat, Tumble Pigweed		

* See glyphosate label for complete list of weeds controlled at each rate as the EMPHASIS® MAX + glyphosate combination controls ~70 weeds, not all are listed here.

** Glyphosate resistant biotypes will not be controlled.



[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

BROMOTRIL® and EMPHASIS® MAX are registered trademarks of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS:

Wait at least 1 day after application before seeding. Allow adequate time for weed control. Avoid overnight storage of spray mixtures when possible. Check weather conditions in advance. Premixing EMPHASIS® MAX spray solutions in nurse tanks is not recommended. Use a water volume of at least 40 L/ac (10 US gal/ac) for adequate coverage.