



ADAMA

NEW

CAZADO™

The ONLY solution that provides true dual-mode of action control of Wild Oats, giving wheat growers their first ever in-crop opportunity to proactively protect against grassy weed resistance.



GROUP 1 & 2

Active Ingredient:

Pinoxaden 100 g/L + Thien carbazone-methyl 8.33 g/L = OD

Application Rates:

Rate: 243 ml/ac

Packaging**& Acres Treated:**

Case: 2 × 9.72 L jug

Drum: 77.8 L with built-in paddle mixer

Acres Treated: 40 ac/jug;
320 ac/drum with
built-in paddle mixer

Water Volume:

Ground: 20–40 L/ac (5–10 US gal/ac)

Aerial: 12 L/ac (3 US gal/ac)

Rainfastness:

1 hour

KEY BENEFITS:

- Only dual-mode of action option
- Full rates of Pinoxaden (Group 1) & Thien carbazone-methyl (Group 2) in one formulation
- The best in-crop herbicide option for Wild Oat management in wheat
- Proactive approach to delay future onset of Wild Oat resistance
- Tank Mix Flexibility

REGISTERED CROPS:

Crop	Stage
Spring Wheat	1 to 6 leaves on main stem, plus 3 tillers but prior to jointing (presence of first node)
Winter Wheat	Apply either in the fall or spring when the majority of plants have one leaf to full tillering, but prior to jointing (presence of first node)

WEEDS CONTROLLED:**Grassy Weeds 1-6 Leaf:**

- Barnyard Grass
- Green Foxtail
- Persian Darnel⁵
- Proso Millet
- Volunteer Oats
- Wild Oats¹
- Yellow Foxtail

Broadleaf Weeds 1-6 Leaf:

- Cleavers
- Hemp-Nettle
- Lamb's-Quarters⁵
- Pale Smartweed
- Redroot Pigweed
- Round-Leaved Mallow⁵
- Shepherd's-Purse
- Stinkweed
- Volunteer Canola²
- Wild Buckwheat
- Wild Mustard

⁵Suppression only.

¹Including Wild Oat biotypes resistant to Group 1 or Group 2 herbicides. For better resistant populations control, a weed stage of 1 to 6 leaves, prior to 3rd tiller is recommended.

²Non ALS-tolerant.

CAZADO™ will not control weeds resistant to both Group 1 and Group 2 herbicides.

CAZADO™ will not control Group 2 (ALS resistant) broadleaf weeds.



CAZADO™

HOW IT WORKS:

CAZADO™ is absorbed by the leaves and is rapidly translocated to the growing points of leaves and stems. Actively growing susceptible grasses stop growing within 48 hours of treatment.

MIXING INSTRUCTIONS:

1. Shake CAZADO™ jugs to ensure product is fully mixed. If using CAZADO™ drum with paddle mixer- use the provided key on a cordless drill to engage the mixer at a low speed for a minimum of 2 minutes to ensure the drum is fully mixed. Do not mix on high and avoid over mixing.
2. Fill the spray tank ¼ to ½ full with clean water and begin agitation or bypass.
3. Add any water conditioners (AMS at a rate of 0.405 L/ac of 49% solution if using on **Spring Wheat only**)
4. Add tank mix partner if desired (WDG, WG, SL or SC formulations).
5. Add the appropriate rate of CAZADO™, directly to the spray tank. Maintain sufficient agitation during both mixing and application.
6. Add tank mix partner if formulation is an EC.
7. Add recommended adjuvant of the tank mix partner. If one is not required, you may use the CAZADO™ adjuvant recommendations.
8. Fill the spray tank with balance of water required.
9. Maintain sufficient agitation during both mixing and application of CAZADO™.

ADJUVANT RECOMMENDATION:

Under challenging conditions (heavy weed pressure, stressed environmental conditions, etc) an adjuvant may be added to enhance grassy weed control:

- MSO or COC (ex Cohere or Addit at 0.5% v/v Spring Wheat only)

Do not use additional adjuvant if tank mix partner does not support. Click the product image for more information or reach out to your local Area Business Manager.

PRE-HARVEST INTERVALS:

Spring Wheat for Grain or Straw: 60 days

Winter Wheat for Grain or Straw: 72 days

BEST PRACTICES:

- For best results apply when targeted pest is 2-3 leaf stage and ideal growing conditions.
- Best to apply on fields with no history of resistance to pinoxaden and thien carbazone.
- For best efficacy spray at 10 US gal/ac.
- For more effective control, especially in situations with high water hardness, ammonium sulphate at 202 g/ac (99%) or 0.405 L/ac (49% solution) may be added. If using an ammonium sulphate product with a different concentration, adjust the rate accordingly. Use ammonium sulphate as a water conditioner only in spring wheat applications.

CROP ROTATIONS:

Alfalfa, Spring Barley, Canola, Field Corn, Dry Bean, Flax (including low linolenic acid varieties), Mustard, Spring Oats, Field Peas, Soybeans, Sunflowers, Timothy, Wheat: 10 months

GRAZING RESTRICTIONS:

DO NOT graze the treated crops or cut for forage within 7 days or cut for hay within 30 days of application.

STORAGE:

Do not freeze. Shake jugs well before using. Mix drums utilizing attached drill adapter.



[CLICK HERE FOR FULL PRODUCT DETAILS.](#)

QUICK TIPS:

Do not spray CAZADO™ within three days of temperature that is 3° C or less prior to or following the temperature. Do not spray if time between seeding and spraying exceeds 35 days (as drought hastens crop development) – applicable to all applications.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

CAZADO™ is a trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2026 ADAMA Agricultural Solutions Canada Ltd.