



NEW

SAFACYN® ET

Pre-Harvest/Harvest Aid

A new standard in harvest aid and weed management. This advanced formulation delivers fast and complete crop and weed dry down, while also enabling flexibility to use the adjuvant type and rate that fits your field & agronomic needs.

HERBICIDE



GROUP 14

Active Ingredient:

Saflufenacil 160 g/L SL

Application Rates:

Rate: 156–312 ml/ha or
63–126 ml/ac

Packaging & Acres Treated:

Case: 4 x 2.53 L

Acres Treated: 20–40 acres per jug;
80–160 acres per case

Water Volume:

Ground: 100–200 L/ha (10–20 gal/ac)
Aerial: 50 L/ha (5 gal/acre)

Rainfastness:

Avoid application when heavy rain is in the forecast.

KEY BENEFITS:

- **Faster, more complete crop and weed dry down:** Delivers consistent dry down for a faster, more efficient harvest.
- **Flexible adjuvant options:** Choose the adjuvant type and rate that best fits your crop, weed pressure and environmental conditions.
- **Improved harvest efficiency:** Supports easier cutting, faster combining, reduced fuel use, and lower dockage.
- **Flexible fit across multiple cropping systems:** Fits with all canola herbicide systems and is registered across a wide range of crops including canola, cereals, peas, and soybeans.
- **Cleaner fields next spring:** Helps manage late-season weeds and sets fields up for a cleaner start next season.

REGISTERED CROPS – HARVEST AID & PRE-HARVEST WEED MANAGEMENT:

Improved dry down of Volunteer Canola (all types including Roundup Ready®), Common Ragweed, Canada Fleabane, Redroot Pigweed and Wild Buckwheat. Consult the glyphosate label regarding additional pre-harvest weed control.

SAFACYN® ET may be used as a harvest aid to accelerate the rate of dry down for crop and green weeds and improve crop uniformity to facilitate direct combining of soybeans, dry beans and canola. Early application may result in yield loss.

CROP	TIMING	Seed Production or Restrictions on Glyphosate Use	No Restrictions on Glyphosate Use
Canola	Apply when 80% of seeds have changed colour.	126 ml/ac of SAFACYN® ET + adjuvant	63–126 ml/ac of SAFACYN® ET + 360 g a.e./ac of glyphosate + adjuvant
Dry Common Beans, Soybeans	Apply when 90% of pods have changed colour from green to yellow or brown.		
Wheat, Barley & Triticale	Hard dough stage; a thumbnail impression remains on seed; less than 30% moisture.		



[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



SAFACYN® ET

HOW IT WORKS:

SAFACYN® ET is quickly absorbed into the plant and works by blocking an important enzyme called protoporphyrinogen oxidase (PPO), which is essential for normal plant function. When PPO is inhibited, the weed's cell membranes are damaged, causing the plant to die. Under active growing conditions, susceptible weeds can show symptoms within hours of application, with complete plant death typically occurring within 3–5 days depending on environmental conditions.

MIXING INSTRUCTIONS:

1. Fill clean tank $\frac{1}{2}$ to $\frac{3}{4}$ full of clean water and turn agitation on.
2. Add the required amount of SAFACYN® ET herbicide and continue to agitate.
3. If required, add the correct amount of tank-mix partner while agitating.
4. Add the required amount of adjuvant while agitating.
5. Continue agitating and fill the remainder of the spray tank with water.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS for proper mixing protocol.

ADJUVANT RATE:

Add 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge Adjuvant, Addit® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) at 0.25% v/v.

PRE-HARVEST INTERVAL FOR HARVEST AID AND PRE-HARVEST USE PATTERNS*:

Dry Beans: 2 days

Soybeans, Canola, Wheat, Barley & Triticale: 3 days

When using a tank-mix partner always refer to the PHI of that product.

* For tank mixtures with glyphosate make sure to follow the glyphosate's PHI.

GRAZING RESTRICTION:

- Dry Beans, Soybeans: Do not graze or feed treated hay or straw to livestock.
- Canola, Wheat, Barley & Triticale: Straw can be used as feed or grazed 3 days after pre-harvest application.

REGISTERED AND SUPPORTED TANK MIXES:

- Glyphosate

STORAGE:

Do not freeze.

QUICK TIPS:

SAFACYN® ET has to enter the plant in order to work. When conditions are such that targeted weeds are hardened off, with tougher cuticles, higher surfactant rates (ie. 1% v/v of an MSO or COC) can help improve efficacy.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

SAFACYN® ET is a registered trademark of an ADAMA Group Company.
All other products are trademarks of their respective companies.
© 2026 ADAMA Agricultural Solutions Canada Ltd.