

Pendigan®

Pendimethalin 30% EC

- ▶ **Pendigan** is a residual dinitroaniline herbicide for selective pre-emergence control of annual grass and broad-leaved weeds in many crops.
- ▶ **Pendigan** is a selective herbicide which is absorbed by both the roots and the leaves.
- ▶ **Pendigan** inhibits cell division and cell elongation, thus causing the death of sensitive weeds shortly after their germination or following their emergence from the soil.
- ▶ **Pendigan** must be activated by mechanical incorporation, rainfall or irrigation up to 7 days after spraying.



Available Pack sizes:
500ml, 1ltr, 5ltr

Antidote:

No specific antidote is know.
Treat symptomatically.



Recommendation for use:

Crop	Weeds	Dosage	
		Lt/Hectare	Lt/Acre
Wheat	<i>Phalaris minor, Chenopodium album, Melilotus alba, Portulaca oleracea, Anagallis arvensis, Fumaria parviflora, poa annua</i>	3.3 (Light soil) 4.2 (Medium soil) 5 (Heavy soil)	1.3 (Light soil) 1.7 (Medium soil) 2 (Heavy soil)
Rice (Transplanted & direct sown upland)	<i>Echinochloa colona, E. crusgalli, Fimbristylis miliacea, Marsilea quadrifolia, Alternanthera sessilis, Ammannia baccifera, Ludwigia parviflora, Eclipta alba, Cyperus difformis</i>	3.3-5	1.3-2

Pendigan®

Pendimethalin 30% EC

Recommendation for use:

Crop	Weeds	Dosage	
		Lt/Hectare	Lt/Acre
Cotton	<i>Echinochloa spp.</i> , <i>Euphorbia hirta</i> , <i>Amaranthus viridis</i> , <i>Portulaca oleracea</i> , <i>Trianthema spp.</i> , <i>Eleusine indica</i>	2.5-4.2	1-1.7
Soybean	<i>Echinochloa spp.</i> , <i>Euphorbia spp.</i> , <i>Amaranthus viridis</i> , <i>Portulaca oleracea</i> , <i>Trianthema spp.</i> , <i>Eleusine indica</i>	2.5-3.3	1-1.3