## **WARNING**



# OPTIMUS® 175 EC PLANT GROWTH REGULATOR

A plant growth regulator to reduce the risk of lodging in wheat, barley and oats and promote seed yield increases in ryegrass seed crops

## READ THE ATTACHED BOOKLET COMPLETELY BEFORE MIXING OR APPLYING

### **Active Ingredient:**

Contains 175 g/L TRINEXAPAC-ETHYL in the form of an emulsifiable concentrate

This product is NOT classified as a Dangerous Good for transport in NZ.

Batch number:
Date of Manufacture:
Barcode to be inserted

NET CONTENTS: 1 L, 5 L, 10 L, 20 L

#### REFER TO BOOKLET FOR FULL LABEL INSTRUCTIONS.



#### **WARNING**

Causes skin irritation. Causes serious eye irritation. May cause allergic skin reaction. Hazardous to soil organisms

#### HANDLING PRECAUTIONS

Use appropriate protective equipment as indicated in the personal protection section of this label. Avoid breathing spray mist or vapours. Wash exposed skin thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace.

#### **PCBU RESPONSIBILITES**

A PCBU with management or control of work using this substance must ensure that every worker who uses, handles, manufactures, or stores this hazardous substance (including hazardous waste) is, before the worker is allowed to carry out or supervise work involving those substances, provided with adequate information, training and instruction. This includes the use of personal protective equipment to be used to minimise risks to the health and safety of workers when carrying out work using this substance.

#### **EQUIPMENT**

A PCBU with management or control of work using this substance must ensure that equipment used to handle this substance retains the substance, without leakage, at all of the temperatures and pressures at which the equipment is to be used; and dispenses or applies the substance, without leakage, at a rate and in a manner that the equipment is designed for. The PCBU must ensure that the equipment is accompanied by documentation that is readily available and readily understandable about the use and maintenance of the equipment.

#### **STORAGE**

Store in the original, tightly-closed container in a cool, dry place, out of direct sunlight and away from stockfeed, seeds, fertilisers or foodstuffs. Store in accordance with NZS 8409 Management of Agrichemicals.

#### SHELF LIFE

When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

#### PERSONAL PROTECTION

When mixing or applying wear protective clothing, including overalls, gloves and eye/face protection. Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse.

#### **SPILLAGE**

Exclude all bystanders from the vicinity of the spill and prevent material from entering waterways. Wear appropriate protective clothing as in the personal protection section of this label to clean up spills. Absorb spills with an inert material (such as vermiculite, earth, sand or synthetic absorbent substance) and place in waste containers. Wash area with water and alkaline detergent. Then absorb any remaining liquid with further inert material. Dispose of the waste safely in an approved landfill.

#### **DISPOSAL**

**Container Disposal -** Triple rinse empty container and add rinsate to spray tank. Submit clean empty container to an Agrecovery<sup>®</sup> depot for recycling. Alternatively, puncture and bury in a suitable landfill. Avoid contamination of any water supply with product or empty container.

**Product Disposal -** Dispose of this product only by using according to the label, or through the Agrecovery<sup>®</sup> Chemical Recovery service or other approved facility.

#### **SAFETY DATA SHEET**

A safety data sheet for this product is available at www.adama.com.

#### **FIRST AID**

If medical advice is needed, have product container or label at hand.

**IF ON SKIN:** Wash with plenty of soap and water. If skin irritation or rash occurs, get medical attention. Wash contaminated clothing before reuse.

**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists, get medical attention.

#### **CONDITIONS OF SALE**

ADAMA New Zealand Limited will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than the responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by the Statutes applicable to the sale and supply of these goods. To the extent allowed by such Statutes the liability of ADAMA New Zealand Limited is limited to the replacement of the goods or (at the option of ADAMA New Zealand Limited) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to ADAMA New Zealand Limited within thirty days of delivery.



Registered to and distributed by: ADAMA New Zealand Limited Level 1, 93 Bolt Road Tahunanui, Nelson 7011 PO Box 1799, Nelson 7040 New Zealand Telephone 03 543 8275 www.adama.com

Registered pursuant to the ACVM Act 1997, No. P8562 See www.foodsafety.govt.nz for registration conditions

Approved pursuant to the HSNO Act 1996, Approval No. HSR100710 See www.epa.govt.nz for approval controls

Optimus is a registered trademark of an ADAMA Group Company.



#### **EMERGENCY RESPONSE**

(All hours)
Ring from anywhere in New
Zealand
0800 734 607

In a Transport Emergency
Phone 111
Police or Fire

#### PRODUCT INFORMATION

OPTIMUS 175 EC is absorbed by the leaves and stems of cereals and ryegrass. OPTIMUS 175 EC has no root or soil activity. OPTIMUS 175 EC is absorbed quickly into the plant tissue and rapidly translocated to the growing parts where the elongation of the stem internodes is inhibited, thereby reducing the straw height. This reduction in straw height reduces the risk of lodging in cereals and ryegrass. In ryegrass OPTIMUS 175 EC shortens the straw height, increases seed set and the time of lodging of the crop is delayed. OPTIMUS 175 EC is rainfast within 2 hours of application.

#### **DIRECTIONS FOR USE**

# TAKE ALL REASONABLE STEPS TO ENSURE THAT THE SUBSTANCE DOES NOT CAUSE ANY SIGNIFICANT ADVERSE EFFECTS TO THE ENVIRONMENT BEYOND THE APPLICATION AREA.

CROP	SITUATION	RATE	CRITICAL COMMENTS
Ryegrass seed crops (annual and perennial)	To promote seed yield by increasing seed set	575 mL to 2.3 L in 200-400 L of water/ha	GS31-32 growth stage, i.e. when the 2 <sup>nd</sup> node is visible on the main tiller and the majority of the other tillers have the 1 <sup>st</sup> node detectable above ground level. The higher rate is recommended on high fertility paddocks where there is no moisture stress e.g. under irrigation. OPTIMUS 175 EC should be applied to actively growing ryegrass NOT suffering from disease, drought stress or insect damage.
Barley Oats Wheat	To reduce the risk of lodging	575 mL in 200 L of water/ha	The addition of a surfactant is NOT necessary. Apply at the beginning of stem elongation, as the 1 <sup>st</sup> node becomes visible (GS31). OPTIMUS 175 EC should be applied to actively growing cereals NOT suffering from disease, insect damage, drought stress or nutrient deficiency.  Note: The recommended rate of OPTIMUS 175 EC may cause severe stunting in oats, however final yield should be unaffected.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

#### WITHHOLDING PERIOD

Wheat, barley, oats, ryegrass: Do not graze or cut for stock food for 42 days after application.

#### **EQUIPMENT**

OPTIMUS 175 EC can be applied with conventional aerial and ground spraying equipment.

#### COMPATIBILITY

OPTIMUS 175 EC is compatible with commonly used fungicides and herbicides in cereals. **Note:** An increase in effect may occur if applied in mixture with triazole fungicides under conditions of rapid vegetative growth. The application timing of herbicides mixed with OPTIMUS 175EC may not be optimal for weed control. As formulations of other manufacturer's products are beyond the control of ADAMA New Zealand Limited, all mixtures should be tested prior to use.