



ODEON[®] 720 SC

Reg. no. L7596 Act/Wet 36 of/van 1947
N-AR 0913

A suspension concentrate contact fungicide for the preventive control of diseases in crops and ornamentals as indicated. 'n Suspensiekonsentraat kontak swamdoder vir die voorkomende beheer van siektes in gewasse en sierplante soos aangedui..

FRAC FUNGICIDE GROUP CODE **M5** **FRAC SWAMDODER GROEPKODE**

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL

Chlorothalonil..... 720 g/LChlorothalonil

NET VOLUME/NETTO VOLUME

..... L

REGISTRATION HOLDER/REGISTRASIEHOUER

ADAMA South Africa (Pty.) Ltd.
Reg. no. 1992/001741/07
Ground Floor, Simeka House
The Vineyard Office Estate
99 Jip de Jager Drive
Belville, 7530
T: +27 21 982 1460
infocpt@adama.com

ODEON[®] is the registered trademark of a company of the ADAMA GROUP.
ODEON[®] is die geregistreerde handelsmerk van 'n maatskappy van die ADAMA GROEP.

Batch number
Date of Manufacture

Lotnommer
Datum van Vervaardiging

UN no.: 2996



**CAUTION
VERSIGTIG**



WARNINGS

Beans (seed).....	3 days
Beans (hay).....	42 days
Cucurbits.....	3 days
Groundnuts (hay).....	42 days
Potatoes.....	3 days
Tomatoes.....	3 days

- Harmful if spray mist is inhaled or if the concentrate is swallowed.
- **ODEON 720 SC** may irritate eyes, nose and throat.
- Toxic to fish and very toxic to other aquatic organisms.
- Store in a cool, dry place away from food and feed.
- Keep out of reach of children, uninformed persons and animals.
- **Re-entry interval:** Do not enter treated field until spray deposit has dried unless wearing protective clothing.
- **Aerial application: (Potatoes, beans and groundnuts only).** Inform all inhabitants in the immediate vicinity of the area to be sprayed and issue the necessary warnings. Do not spray over or allow spray drift to contaminate water or adjacent areas.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

Precautions

- Do not inhale spray mist.
- Wear a face shield and rubber gloves when mixing the concentrate.
- Wash with soap and water after accidental skin contact. Wash contaminated work clothes.
- Do not eat, drink or smoke while mixing or spraying or before hands and face are washed.
- Avoid drift of spray onto other crops, grazing, rivers, dams or any other area not under treatment.
- Prevent contamination of food, feed, drinking water and eating utensils.
- Clean applicator after use and dispose of wash water where it will not contaminate other crops, grazing, rivers or dams.
- Invert the empty container over the spray or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water of at least 10 % of that of the container. Add the rinsings to the contents of the spray tank before destroying the empty container in the prescribed manner.

- Do not re-use the empty container for any other purpose.

Resistance Warning

For resistance management, **ODEON® 720 SC** is a group code M5 fungicide. Any fungus population may contain individuals naturally resistant to **ODEON® 720 SC** and other group code M5 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly. These resistant fungi may not be controlled by **ODEON® 720 SC** or any other group code M5 fungicide. To delay fungicide resistance:

- Avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank-mix with products from different fungicide group codes.
- Integrate the control methods (chemical, cultural, biological) into fungicide control programmes.
- For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE:

Use only as indicated.

Compatibility

The compatibility of **ODEON® 720 SC** with other insecticides and fungicides is unknown. The compatibility must be established beforehand on a small scale. When used in tank mixes with other crop protection chemicals the conditions that appear on the labels of those chemicals should be studied and adhered to. If applied by means of a center pivot irrigation system, **ODEON® 720 SC** must not be mixed with any other agricultural remedy or fertilizer.

Mixing Instructions

Half fill the spray tank with clean water. Mix the required volume of **ODEON® 720 SC** separately in a small volume of water before adding to the spray tank while agitating. When **ODEON® 720 SC** is used in a tank mixture with other products, add the products in the following sequence: adjuvant/acidifier, wettable powder, suspension concentrate, emulsifiable concentrate. Fill the spray tank to the required volume with water while agitating the spray mixture. It is important to maintain agitation during mixing and application. Prepared spray mixtures must not be left for any length of time, e.g. overnight.

APPLICATION

ODEON® 720 SC must be applied as a preventive application. Good coverage on both upper and lower surfaces of leaves should be ensured. Repeat application if 12 mm or more rain or irrigation is received within 72 hours after application. If weather conditions favorable for the development of disease prevail for long periods, it should be considered to alternate **ODEON® 720 SC** with registered systemic fungicides.

Ground Application

ODEON® 720 SC can be applied by any suitable medium or high volume spraying apparatus provided it is correctly calibrated, has an efficient agitation system and is able to distribute the spray mixture evenly over the target area.

Aerial application: (Potatoes, beans and groundnuts only).

Aerial application of **ODEON® 720 SC** may only be done by a registered Aerial Application Operator using a correctly calibrated, registered aircraft according to the instructions of SANS 10118 (Aerial Application of Agricultural Pesticides). Ensure that the spray mixture is distributed evenly over the

target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria be met:

- **Volume:** A spray mixture volume of 30-35 l/ha is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy, or be held responsible for any adverse effects if this product is applied aurally at a lower volume rate than recommended above.
- **Droplet coverage:** 50-70 droplets/cm² must be recovered at the target area.
- **Droplet size:** A droplet spectrum with a VMD of 220-250 micron is recommended. Limit the production of fine droplets less than 150 micron (high drift and evaporation potential) to a minimum.
- **Flying height:** Maintain the height of the spray boom at 3-4 m above the target. Do not spray when aircraft is diving, climbing or when banking
- Use suitable atomising equipment that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product. The spraying system must produce a droplet spectrum with the lowest possible Relative Span.
- Position all the atomisers within the inner 60-75 % of the wingspan to prevent droplets from entering the wingtip vortices.
- The difference in temperature between the wet and dry bulb thermometers, of a whirling hygrometer, should not exceed 8 °C.
- Stop spraying if the wind speed exceeds 15 km/h.
- Stop spraying under turbulent, unstable and dry conditions during the heat of the day.
- Spraying under temperature inversion conditions (spraying in or above the inversion layer) and/or high humidity conditions (relative humidity 80 % and above) may lead to the following:
 - a) reduced efficacy due to suspension and evaporation of small droplets in the air (inadequate coverage).
 - b) damage to other sensitive crops and/or non-target areas through drifting of the suspended spray cloud away from the target field.
- Ensure that the Aerial Spray Operator knows exactly which fields to spray.
- Obtain an assurance from the Aerial Spray Operator that the above requirements will be met and that relevant data will be compiled in a logbook and kept for future reference.
- **Note:** For potatoes and beans, apply in not less than 30 l water/ha and for groundnuts in 50 l water/ha.

Centre Pivot Application

ODEON® 720 SC may be applied through a centre pivot irrigation system. The supply tank and connecting pipes should be leak free and fitted with an efficient agitation mechanism. The spray mixture must be injected into the irrigation system by means of a positive displacement metering pump. Non- return valves in the system must be in place to prevent a back flow of the spray mixture to the water sources. If a chemigation system is not fitted to the centre pivot and the spray mixture is injected directly into the irrigation system the centre pivot should be set at maximum speed and minimum water delivery. The spray mixture must be applied evenly over the target area in one continuous operation. Run-off must be prevented in order to eliminate the risk of pollution or phytotoxicity. Flush the whole system with clean water after use. Ensure that the flush water does not contaminate water sources.

APPLICATION RATES

CROP/TARGET	DOSAGE	REMARKS
Beans Rust <i>(Uromyces appendiculatus)</i>	Dry land Ground application 0,7–1,4 l/ha Aerial application 0,7–1,4 l/ha Irrigated Ground application 1,4 l/ha Aerial application 1,4 l/ha	Commence application as a preventive spray when weather conditions favour the development of the disease (periods of high humidity of 8–10 hours). Use the higher dosage rate on bigger plants or when climatic conditions favouring the development of the disease prevail. Repeat application every 7–14 days. Apply as a full cover spray in 250–500 l water/ha depending on plant size. Apply as above in at least 30 l water/ha. Apply as above as a full cover spray in 250–500 l water/ha depending on plant size. Apply as above in least 30 l water/ha.
Cucurbits Anthracnose <i>(Colletotrichum orbiculare)</i> Downy mildew <i>(Pseudoperonospora cubensis)</i>	1,0–2,0 l/ha	Apply as a full cover preventive spray or when weather conditions favour the development of the disease. Apply in at least 500 l water/ha. Use the higher dosage rate when weather conditions favour the development of the disease. Repeat application at 7-10 day intervals. Ensure good coverage of all plant parts.
Groundnuts Leafspot <i>(Cercospora spp., Cercosporidium spp., Phoma spp., Botrytis rot (Botrytis spp.))</i>	Dry land Ground application 0,7–1,55 l/ha Aerial application 1,55 l/ha Irrigation 1,55 l/ha	Apply as a full cover preventive application in not less than 500 l water/ha. Repeat every 10–14 days at the lower rate for low infections. Commence application 6–10 weeks after planting or earlier if weather conditions favour the development of the disease. Use the higher rate for more severe infections and increase the water volume from 500 l to 1,500 l/ha depending on plant size. Repeat application every 7-10 days for high infections. Apply in at least 50 l water/ha when tractor applications are impractical due to very wet conditions. Apply as above in up to 1500 l water/ha. Repeat application every 7-14 days

CROP/TARGET	DOSAGE	REMARKS
Ornamentals Flowers and Lawns Gladioli Leaf spot Lawns and Turf Dollar spot Roses Black spot	190 ml/100 ℓ water Preventive 350 ml/100 ℓ water Corrective 520 ml/100 ℓ water 190 ml/100 ℓ water	Apply as a full cover preventive spray every 7–14 days. Do not apply during flowering. Apply at least 10 ℓ spray mixture/100 m ² in a spray programme at 7–10 day intervals. Apply twice within 7 days and thereafter use the preventive programme as above. Apply as a preventive full cover spray programme at intervals of 7–14 days.
Potatoes Early blight (Alternaria solani) Late blight (<i>Phytophthora infestans</i>)	Ground application 0,7–2,0 ℓ/ha Centre application pivot 2,0 ℓ/ha Aerial application 1,0–2,0 ℓ/ha	Commence application as a preventive treatment before the disease appear or if weather conditions favour the development of the disease. Low infection: Apply at a rate of 0,7 ℓ/ha at 7 day intervals or at 1,0 ℓ/ha at 10 day intervals under dry land cultivation. Apply in 500 ℓ water/ha. High infection: Apply at a rate of 2,0 ℓ/ha and repeat application at 5–7 day intervals in not less than 500 ℓ and up to 1500 ℓ water/ha depending on the density of the foliage and the size of the plants. Use this rate also for potatoes under irrigation. If necessary alternate ODEON® 720 SC with registered systemic products. Refer to ground application for usage directions. Apply as a spray programme starting 2 weeks after emergence at intervals of 7–10 days. Do not exceed a 10 day interval between applications. Refer to section under centre pivot application under APPLICATION. It is important to ensure that full cover, even wetting is obtained on all parts of potato plants. NOTE: Up-to-date potatoes cultivated under centre pivot irrigation are extremely susceptible to late blight infection from January - April especially in the Limpopo Province and ODEON® 720 SC might not give adequate control under such high disease pressure during this period. It is suggested to apply ODEON® 720 SC at 2,5 ℓ/ha at spray intervals of 4-5 days during this period on this cultivar. Use as for ground application. Apply the lower rate for lower infections and the higher rate for more severe infections. Apply in not less than 30 ℓ water/ha.

CROP/TARGET	DOSAGE	REMARKS
<p>Tomatoes Early blight (<i>Alternaria solani</i>) Late blight (<i>Phytophthora infestans</i>) Botrytis rot (<i>Botrytis cinerea</i>)</p>	<p>015 ml/100 l water 190 ml/100 l water 210 ml / 100 l water 380 ml / 100 l water</p>	<p>Apply as a preventive application from flowering before any symptoms appear or when weather conditions favour the development of the disease. Apply as a full cover spray using up to 2,000 l spray mixture/ha depending on plant size or whether plants are trellised or not. When Botrytis rot threatens apply the higher rate at 7 day intervals for both high and low volume applications. Ensure good coverage of both upper and lower leaf surfaces.</p> <p>High volume: Apply at 1,000-2,000 l spray mixture/ha on mature plants.</p> <p>Low infection: Apply every 10–14 days. Do not use less than 1,05 l ODEON® 720 SC/ha on mature plants.</p> <p>High infection: Apply every 7–10 days. Do not use less than 1,9 l ODEON® 720 SC/ha on mature plants.</p> <p>Low volume: Apply 500 l spray mixture/ha on mature plants.</p> <p>Low infection: Apply every 10–14 days. Do not apply less than 1,05 l ODEON® 720 SC/ha on mature plants.</p> <p>High infection: Apply every 7–10 days. Do not apply less than 1,9 l ODEON® 720 SC/ha on mature plants.</p>

WAARSKUWINGS

Onthoudingsperiodes (Minimum aantal dae tussen laaste toediening en oes):

Aartappels	3 dae
Bone (saad).....	3 dae
Bone (hooi).....	42 dae
Grondbone (hooi).....	42 dae
Pampoengewasse	3 dae
Tamaties	3 dae

- Skadelik indien spuitnewel ingeasem of die konsentraat ingesluk word.
- **ODEON® 720 SC** mag die oë, neus en keel irriteer.
- Giftig vir visse en baie giftig vir ander waterorganismes.
- Berg in 'n koel droë plek weg van voedsel en voer.
- Hou buite bereik van kinders, oningeligte persone en diere.
- **Herbetredingsinterval:** Moet nie behandelde gebied binnegaan voordat spuitnewel droog is nie tensy beskermende klere gedra word.
- **Lugtoediening: (Slegs aartappels, bone en grondbone).** Stel alle inwoners in die onmiddellike omgewing van die gebied wat bespuit gaan word in kennis en reik die nodige waarkuwings uit. Moet nie toedien oor of toelaat dat spuitnewel dryf oor omliggende gebiede of water nie.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale grond, klimaats- en bergingstoestande, kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die siekte teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in geval van enige onsekerheid.

Voorsorgmaatreëls

- Moet nie die spuitnewel inasem nie.
- Dra 'n gesigskerm en rubberhandskoene wanneer die konsentraat vermeng word.
- Was met water en seep as dit toevallig met die vel in aanraking kom. Was besoedelde klere daaglik.
- Moet nie eet, drink of rook tydens vermenging en bespuiting of voordat hande en gesig gewas is nie.
- Verhoed dat die spuitnewel wegdryf na ander gewasse, weiding, riviere, damme of enige ander gebied wat nie behandel moet word nie.
- Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.

- Maak toedieningsapparaat skoon na gebruik en raak van waswater ontslae waar dit nie ander gewasse, weiding, riviere of damme sal besoedel nie.
- Keer die leë houer om oor die spuit- of mengtenk en laat vir 30 sekondes dreineer nadat die vloei tot 'n gedrup verminder het. Spoel daarna die leë houer drie keer uit met 'n volume water gelykstaande aan minstens 10 % van die volume van die houer. Gooi die spoelwater in die spuittenk voor die leë houer op die voorgeskrewe manier vernietig word.
- Moet nie die leë houer vir enige ander doel gebruik nie.

Weerstandswaarskuwing

ODEON® 720 SC is 'n groepkode M5 swamdoder. Enige populasie van 'n spesifieke swam mag individue insluit wat 'n natuurlike weerstand teen **ODEON® 720 SC** of enige ander groepkode M5 swamdoder het. Indien hierdie swamdoders herhaaldelik aangewend word, kan die weerstandbiedende individue uiteindelik die swampopulasie oorheers. Hierdie weerstandbiedende swamme sal waarskynlik nie deur **ODEON® 720 SC** of enige ander groepkode M5 swamdoder beheer word nie. Om weerstand teen swamdoders te vertraag:

- Vermyn die eksklusiewe herhaaldelike gebruik van swamdoders met dieselfde groepkode. Wissel af met, of gebruik tenkmengsels van produkte in verskillende swamdodergroepkodes.
- Integreer ander beheermaatreëls (chemies, verbouing, biologies) in swamdoderprogramme.
- Vir spesifieke inligting oor weerstandsbestuur kontak die registrasiehouer van hierdie produk.

GEBRUIKSAANWYSINGS:

Gebruik slegs soos aangedui.

Verenigbaarheid

Die verenigbaarheid van **ODEON® 720 SC** met ander insek- en swamdoders is onbekend. Verenigbaarheid moet vooraf op 'n klein skaal bepaal word. Indien dit egter in tenkmengsels met ander lanboumiddels gebruik word, moet daardie middels se etiketvoorskrifte bestudeer en nagekom word. Indien **ODEON® 720 SC** deur 'n spulpuntbesproeiingstelsel toegedien word, moet dit nie met ander landboumiddels of kunsmis gemeng word nie.

Meng Instruksies

Maak die spuittenk halfvol met skoon water. Meng die verlangde hoeveelheid **ODEON® 720 SC** apart in water en voeg in die tenk terwyl geroer word. Indien **ODEON® 720 SC** in 'n tenkmengsel met ander produkte gebruik wil word, voeg die produkte in die volgende volgorde in die spuittenk: bevordering-/versuringsmiddel, benatbare poeier, suspensiekonsentraat, emulgeerbarekonsentraat. Vul die spuittenk tot die verlangde volume met water terwyl geroer word. Dit is belangrik om roering gedurigdeur te handhaaf tydens vermenging en toediening. Aangemaakte mengsels moet nie toegelaat word om vir lang periodes in die tenk te staan nie, bv. oornag.

TOEDIENING

Algemeen

ODEON® 720 SC moet as 'n voorkomende bespuiting toegedien word. 'n Goeie bedekking moet op die bo- en onderkante van blare verseker word. Die bespuiting moet herhaal word indien 12 mm of meer reën of besproeiing binne 72 uur na toediening ontvang is. Indien weerstoestande wat bevorderlik is vir die ontwikkeling van siekte langdurig voorkom, moet oorweeg word om **ODEON® 720 SC** met sistemiese middels af te wissel.

Grondtoediening

ODEON® 720 SC kan deur middel van enige geskikte medium- of hoëvolume spuitapparaat toegedien word mits dit korrek gekalibreer is, oor 'n doeltreffende roermeganisme beskik en in staat is om die spuitmengsel eweredig oor die teikengebied te versprei.

Lugtoediening: (Slegs aartappels, bone en grondbone)

ODEON® 720 SC kan slegs deur 'n geregistreerde Lugbespuitingsoperateur met 'n korrek gekalibreerde, geregistreerde vliegtuig volgens die instruksies van SANS 10118 (Aerial Application of Agricultural Pesticides) uit die lug bespuit word. Verseker dat die spuitmengsel eweredig oor die teikenarea versprei word, en die verlies aan spuitmengsel tydens toediening tot 'n minimum beperk word. Dit is daarom belangrik om aan die volgende vereistes te voldoen:

- **Volume:** 'n Spuitmengsel volume van 30-35 l/ha word aanbeveel. Hierdie produk is nie teen 'n verlaagde volume getoets nie. Die registrasiehouer kan nie effektiwiteit waarborg, of verantwoordelik gehou word vir enige nadelige effekte indien hierdie produk teen 'n laer volume, as hierbo aanbeveel, toegedien word nie.
- **Druppelbedekking:** 50-70 druppels/cm² moet op die teikenarea herwin word.
- **Druppelgrootte:** 'n Druppelspektrum met 'n VMD van 220-250 mikron word aanbeveel. Beperk die produksie van druppels kleiner as 150 mikron (hoë drywing en verdampingspotensiaal) tot 'n minimum.
- **Vlieghoogte:** Handhaaf die hoogte van die spuitbalk bo die teiken op 3-4 m. Moet nie spuit wanneer die vliegtuig duik, klim of draai nie.
- Gebruik geskikte atomiseringsapparaat wat die vereiste druppelgrootte en bedekking sal produseer, maar die minste verlies van produk verseker. Die spuitstelsel moet 'n druppelspektrum met die kleinste moontlike Relatiewe Span produseer.
- Plaas al die atomiseerders in die binneste 60-75 % van die vlerkspan om te verhoed dat druppels binne-in die vlerkpuntvorteks beweeg.
- Die verskil in temperatuur tussen die nat- en droëboltermometer van 'n swaaihygrometer, moet nie 8 °C oorskry nie.
- Stop bespuiting indien die windspoed 15 km/u oorskry.
- Stop bespuiting tydens turbulente, onstabiele en droë toestande gedurende die hitte van die dag.
- Bespuiting onder temperatuur inversie toestand (deur bo of binne die inversie laag te spuit) en/of hoë lugvog toestand (relatiewe humiditeit 80 % en meer) mag tot volgende probleme aanleiding gee:
 - a) Verlaagde doeltreffendheid aangesien die druppels as 'n wolk in die lug bly hang en moontlik verdamp (onvoldoende bedekking op teiken).
 - b) Skade aan nie-teiken gewasse of sensitiewe areas as gevolg van wegdrywing van die spuitwolk na nie-teiken area.
- Verseker dat die Lugbespuitingsoperateur presies weet watter lande bespuit moet word.
- Dit is noodsaaklik om 'n versekering van die Lugbespuitingsoperateur te verkry dat aan al die bogenoemde vereistes voldoen sal word en dat data van belang in 'n logboek saamgevat is vir toekomstige verwysing.
- **LET WEL:** Dien toe in nie minder nie as 30 l water/ha vir aartappels en bone en in 50 l water vir grondbone.

Spilpunt Toediening

ODEON® 720 SC mag deur 'n spilpunt besproeiingsstelsel toegedien word. Die toevoertenk en alle pypleidinge moet sonder enige lekkasies wees en moet met 'n doeltreffende roermeganisme toegerus wees. Die spuitmengsel moet met 'n positiewe verplasingmeetpomp in die stelsel ingevoer word. Terugslagkleppe moet in die stelsel wees om te verhoed dat die spuitmengsel na die waterbron kan terugvloei. Indien die spilpunt nie voorsien is van 'n "Chemigation" stelsel nie en die spuitmengsel direk in die besproeiingsstelsel ingespuet word, behoort die spilpunt teen maksimum spoed en minimum lewering ingestel te word. Die spuitmengsel moet eweredig oor die teikengebied in een deurlopende aksie aangewend word. Afloop van spuitmengsel moet vermy word om besoedeling en plantskade te voorkom. Spoel die hele stelsel met skoon water uit na gebruik en sorg dat die spoelwater nie waterbronne besoedel nie.

TOEDIENINGSHOEVEELHEDE

GEWAS/TEIKEN	DOSIS	OPMERKINGS
Aartappels Vroeëroes (<i>Alternaria solani</i>) Laatroes (<i>Phytophthora infestans</i>)	<p>Grondtoediening 0,7-2,0 l/ha</p> <p>Spilpuntoediening 2,0 l/ha</p> <p>Lugtoediening 1,0-2,0 l/ha</p>	<p>Dien voorkomend toe voor die siekte verskyn of as weerstoestande die ontwikkeling van die siekte bevoordeel.</p> <p>Lae infeksie: Dien toe teen 'n dosis van 0,7 l/ha elke 7 dae of teen 'n dosis van 1,0 l/ha elke 10 dae onder droëlandverbouing. Dien toe in 500 water/ha.</p> <p>Hoë infeksie: Dien toe teen 'n dosis van 2,0 l/ha en herhaal toediening met 5-7 dae tussenposes in nie minder nie as 500 l tot 1,500 l water/ha afhangende van plantgrootte en digtheid van loof. Gebruik hierdie dosis ook op aartappels onder besproeiing. Indien nodig moet ODEON® 720 SC afgewissel word met geregistreerde sistemiese middels.</p> <p>Verwys na grondtoediening vir gebruiksaanwysings asook die gedeelte by spilpuntoediening onder TOEDIENING. Dien in 'n spuitprogram beginnende 2 weke na opkom met 7–10 dae tussenposes toe. Dit is belangrik om te verseker dat 'n volle dekhoediening egalig oor alle dele van die plante verkry word. Moet nie 'n 10 dae tussenpose oorskry nie.</p> <p>LET WEL: “Up-to-date” aartappels verbou onder spilpuntbesproeiing is baie vatbaar vir laatroes infeksies gedurende Januarie - April veral in die Limpopo Provinsie en ODEON® 720 SC mag moontlik nie voldoende beheer gee onder sulke hoë besmettingsdruk nie. Daar word aanbeveel dat 2,5 l elke 4-5 dae toegedien word gedurende dié tydperk in die geval van hierdie kultivar.</p> <p>Gebruik soos vir grondtoediening. Dien die laer toedieningshoeveelheid toe vir lae infeksies en die hoër toedieningshoeveelheid vir swaarder infeksies. Dien toe in nie minder nie as 30 l water/ha.</p>
Bone Bruinroes (<i>Uromyces appendiculatus</i>)		<p>Dien as 'n voorkomende bespuiting toe as weerstoestande die ontwikkeling van die siekte bevoordeel (tydperke van 8–10 uur van hoë humiditeit). Gebruik die hoër toedieningshoeveelheid op groter plante of as weerstoestande wat die siekte bevoordeel langdurig voortduur. Herhaal toediening elke 7–14 dae.</p>

GEWAS/TEIKEN	DOSIS	OPMERKINGS
	Droëland Grondtoediening 0,7–1,4 l/ha Lugtoediening 0,7–1,4 l/ha Besproeiing Grondtoediening 1,4 l/ha Lugtoediening 1,4 l/ha	Dien as 'n volle dekhoediening in 250–500 l water/ha toe afhange van plantgrootte. Dien toe soos bo in ten minste 30 l water/ha. Dien toe soos bo as 'n volle dekhoediening in 250–500 l water/ha afhange van plantgrootte. Dien toe in ten minste 30 l water/ha.
Grondbone Blaarvlek (<i>Cercospora</i> spp., <i>Cercosporidium</i> spp., <i>Phoma</i> spp.) Botrytis vrot (<i>Botrytis</i> spp.)	Droë land Grondtoediening 0,7–1,55 l/ha Lugtoediening 1,55 l/ha Besproeiing 1,55 l/ha	Dien voorkomend toe as 'n volle dekhoediening in nie minder nie as 500 l water/ha. Herhaal elke 10–14 dae teen die laer dosis vir lae infeksies. Begin met toediening ongeveer 6–10 weke na plant of vroeër indien weerstoestande die ontwikkeling van die siekte bevorder. Gebruik die hoër toedieningshoeveelheid vir swaarder infeksies en vermeerder die watervolume van 500 l na 1500 l/ha afhange van plantgrootte. Herhaal toediening elke 7-10 dae vir hoër infeksies. Dien toe in ten minste 50 l water/ha wanneer trekkertoedienings nie prakties is nie a.g.v. baie nat toestande. Dien toe soos hierbo in tot 1500 l water/ha. Herhaal toediening elke 7-14 dae
Pampoengewasse Antraknose (<i>Colletotrichum orbiculare</i>) Donsige skimmel (<i>Pseudoperonospora cubensis</i>)	1,0–2,0 l/ha	Dien toe as 'n voorkomende volle dekhoediening wanneer weerstoestande die ontwikkeling van die siekte bevorder. Dien toe in ten minste 500 l water/ha. Gebruik die hoër toedieningshoeveelheid as weerstoestande die ontwikkeling van die siekte bevoordeel. Herhaal toediening met 7-10 dae tussenposes. Verseker deeglike bedekking van alle plantdele.

GEWAS/TEIKEN	DOSIS	OPMERKINGS
Sierplante, Blomme en Grasperke Swaardlelies Blaarvlek Grasperke Dollarvlek Rose Swartvlek	 190 ml/100 l water Voorkomend 350 ml/100 l water Korrektief 520 ml/100 l water 190 ml/100 l water	 Dien toe as 'n volle voorkomende dekhoediening en herhaal elke 7–14 dae. Moet nie tydens blom toedien nie. Dien ten minste 10 l spuitmengsel/100 m ² toe in 'n spuitprogram. Herhaal elke 7–10 dae. Dien twee keer toe binne 7 dae en gebruik daarna die voorkomende program soos hierbo aangedui. Dien toe as 'n voorkomende volle dekhoediening. Herhaal toediening elke 7–14 dae.
Tamaties Vroeëroes (<i>Alternaria solani</i>) Laatroes (<i>Phytophthora infestans</i>) Botrytis vrot (<i>Botrytis cinerea</i>)	 105 ml/100 l water 190 ml/100 l water 210 ml/100 l water 380 ml/100 l water	Dien toe as 'n voorkomende toediening vanaf blom voor enige simptome verskyn of as weerstoestande die ontwikkeling van die siekte bevorder. Dien toe as 'n volle dekhoediening en gebruik tot 2,000 l spuitmengsel/ha afhangende van plantgrootte en of die plante opgelei is of nie. Indien Botrytis dreig, dien die hoër dosis toe met 7 dae tussenposes vir beide hoë- en laevolume toediening. Verseker deeglike benatting van die boonste sowel as die onderste blaaroppervlakke. Hoëvolume: Dien toe teen 1,000–2,000 l spuitmengsel/ha op volwasse plante. Lae infeksie: Dien elke 10–14 dae toe. Moet nie minder as 1,05 l ODEON® 720 SC /ha op volwasse plante toedien nie. Hoë infeksie: Dien elke 7–10 dae toe. Moet nie minder as 1,9 l ODEON® 720 SC /ha toedien op volwasse plante nie. Laevolume: Dien 500 l spuitmengsel/ha op volwasse plante toe. Lae infeksie: Dien elke 10–14 dae toe. Moet nie minder as 1,05 l ODEON® 720 SC /ha toedien op volwasse plante nie. Hoë infeksie: Dien elke 7–10 dae toe. Moet nie minder as 1,9 l ODEON® 720 SC /ha op volwasse plante toedien nie.

