



FOLPAN[®] 80 WDG

Reg. no. L5333 Act/Wet 36 of/van 1947

A water dispersible granule organic-based fungicide for the preventive control of anthracnose, downy mildew, dead-arm disease and Botrytis rot in wine grapes, dead-arm disease, anthracnose and downy mildew pre-blossom in table grapes as well as late blight and early blight in potatoes and tomatoes.

'n Waterdispergeerbare korrel organies-gebaseerde swamdoder vir die voorkomende beheer van antraknose, donsige skimmel, streepvleksiekte en botritisvrot in wyndruiwe, streepvleksiekte, antraknose en donsige skimmel voorblom in tafeldruiwe asook laat- en vroeëroes in aartappels en tamaties.

FRAC FUNGICIDE GROUP CODE

M4

FRAC SWAMDODER GROEPKODE

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL

folpet (phtalimide) 800 g/kg folpet (ptalimied)

NET MASS/NETTO MASSA

..... kg

REGISTRATION HOLDER/REGISTRASIEHOUER

ADAMA South Africa (Pty.) Ltd.

Reg. no. 1992/001741/07

21 Viben Street

7560 Brackenfell

T: +27 21 982 1460 F: +27 21 982 5810

infocpt@adama.com

**FOLPAN[®] is the registered trademark of a company of the ADAMA GROUP.
FOLPAN[®] is die geregistreerde handelsmerk van 'n maatskappy van die ADAMA GROEP.**

CONTACT IN EMERGENCY/KONTAK IN NOOD

Griffon Poison Information Centre: +27 82 446 8946

Tygerberg Poison Information Centre: +27 21 931 6129

Batch number

Lotnommer

Date of manufacture

Datum van vervaardiging

UN no: 3077



**CAUTION
VERSIGTIG**



WARNINGS

Withholding periods (minimum number of days between last application and harvest):

Tomatoes	3 days
Table grapes	Not later than 5 mm berry diameter
Wine grapes	14 days

- Irritating to skin and eyes and may cause sensitization by skin contact.
- Toxic to fish.
- Store in a cool place away from food and feed.
- Keep out of reach of children, uninformed persons and animals.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment, or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

- Do not eat, drink or smoke whilst handling this product.
- Prevent contamination of food, feed, drinking water and eating utensils.
- Wear rubber gloves and face shield when handling the concentrate. If you get it in your eyes, flush it out at once. If you get it on your skin, wash it off at once. If you spill it on your clothes, change and wash them.
- Containers and packages must be completely emptied before being disposed of. Shake out thoroughly into the mixing or spray tank before destroying the empty container.
- While spraying, avoid contact with the spray as much as you can. Avoid spray drift onto other crops, grazing, rivers or dams.
- Clean applicator thoroughly after use and dispose of wash water where it will not contaminate crops, grazing, rivers or dams.
- Destroy empty container and do not use for any other purpose. Change and wash your work clothes. Wash yourself.

DIRECTIONS FOR USE

Use only as indicated.

Compatibility

FOLPAN® 80 WDG is compatible with most insecticides and fungicides commonly used with the exception of lime, lime sulphur and Bordeaux mixture.

Mixing Instructions

Mix the required quantity of **FOLPAN® 80 WDG** separately in a small amount of water to make a smooth cream before adding it to the recommended amount of water in the spray tank. Start agitation whilst adding **FOLPAN® 80 WDG** into the spray tank and maintain agitation throughout the application operation. Spray thoroughly to ensure complete coverage of the plant, including the under-surface of the leaves. **FOLPAN® 80 WDG** is recommended as a preventive treatment.

APPLICATION RATES

CROP AND TARGET	DOSAGE	REMARKS
Wine and table grapes Anthracnose Dead-arm disease Botrytis rot Downy mildew	125 g/100 L water 125 g/100 L water	Apply every two weeks or weekly when weather conditions are favourable for the development of the disease. Pre-blossom period: Apply the first application when shoots are 10 cm long. Repeat at 7- to 10-day intervals depending on weather conditions which will favour the development of the disease. Blossom to 80 % petal drop: Only apply when the flowering period lasts for more than two weeks. Post-blossom period: Apply twice or more at 7- to 14-day intervals, depending on weather conditions.
Potatoes Late blight (<i>Phytophthora infestans</i>) Early blight (<i>Alternaria solani</i>)	1.25 kg/ha	Preventive treatment: Apply from emergence at 7- to 10-day intervals in not less than 500 L water/ha in the pre-blossom stage and in 1000 L water/ha in the post-blossom stage. For the control of early and late blight, particularly after flowering, it is important to ensure thorough coverage of the plants. Add a suitable adjuvant at the recommended rate.

CROP AND TARGET	DOSAGE	REMARKS
<p>Tomatoes</p> <p>Late blight (<i>Phytophthora infestans</i>)</p> <p>Early blight (<i>Alternaria solani</i>)</p>	1.25 kg/ha	<p>Apply as a preventive application shortly after transplant and before symptoms of the disease appear. Depending on plant size use 300–1500 L water/ha. Ensure thorough coverage of both upper and lower leaf surfaces. Repeat application at 7- to 10-day intervals.</p>

WAARSKUWINGS

Onthoudingsperiodes (minimum aantal dae tussen laaste toediening en oes):

Tamaties	3 dae
Tafeldruiwe	Nie na 5 mm korreldeursnee nie
Wyndruiwe	14 dae

- Irriterend vir die vel en oë en mag sensitisering veroorsaak deur velkontak.
- Giftig vir visse.
- Stoor in 'n koel plek weg van voedsel en voer.
- Hou buite bereik van kinders, oningeligte persone en diere.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale klimaats- en bergingstoestande, kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die siekte teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

VOORSORGMAATREËLS

- Moet nie eet, drink of rook terwyl die produk hanteer word nie.
- Vermoed besoedeling van voedsel, voer, drinkwater en eetgerei.
- Dra rubberhandskoene en gesigskerm wanneer die konsentraat hanteer word. Indien dit in u oë kom, spoel onmiddellik uit. Indien dit op u vel kom, was dit dadelik af. Indien dit op u klere mors, trek uit en was dit.
- Houers en verpakkings moet heeltemal leegemaak word voordat hulle vernietig word. Skud die houer deeglik uit in die meng- of spuitenk voordat leë houer vernietig word.
- Vermoed kontak met die spuitstof sover moontlik tydens toediening. Vermoed oorwaai van spuitnewel na ander gewasse, weiding, riviere of damme.
- Maak toediener deeglik skoon na gebruik en gooi waswater waar dit nie gewasse, weiding, riviere of damme sal besoedel nie.
- Vernietig leë houer en moet dit nie vir enige ander doel gebruik nie. Trek werksklere uit en was dit. Was uself.

GEBRUIKSAANWYSINGS

Gebruik slegs soos aangedui.

Verenigbaarheid

FOLPAN® 80 WDG is verenigbaar met die meeste insek- en swamdoders wat algemeen gebruik word met die uitsondering van kalk, kalkswael en Bordeaux-mengsel.

Menginstruksies

Meng die vereiste hoeveelheid **FOLPAN® 80 WDG** apart met 'n klein hoeveelheid water tot 'n gladde room voordat dit by die aanbeveelde hoeveelheid water in die spuitenk gevoeg word. Roer voortdurend tydens die byvoeging van **FOLPAN® 80 WDG** in die spuitenk asook tydens toediening. Dien deeglik toe om algehele bedekking van die plant te verseker, insluitend die onderkant van die blare. **FOLPAN® 80 WDG** word as 'n voorkomende behandeling aanbeveel.

TOEDIENINGSHOEVEELHEDE

GEWAS EN TEIKEN	DOSIS	OPMERKINGS
Wyn- en tafeldruive Anthraknose Streepvleksierte Botritis-vrot Donsige skimmel	125 g/100 L water 125 g/100 L water	Dien elke twee weke of weekliks toe wanneer weerstoestande gunstig is vir die ontwikkeling van die siekte. Voorblomperiode: Dien toe wanneer die lote 10 cm lank is. Herhaal met 7- tot 10-dae tussenposes afhangend van weerstoestande wat gunstig is vir die ontwikkeling van die siekte. Blom tot 80 % blomblaarval: Dien alleenlik toe wanneer die blomperiode langer as twee weke is. Nablomperiode: Dien twee keer of meer met tussenposes van 7 tot 14 dae toe, afhangend van weerstoestande.
Aartappels Laatroes (<i>Phytophthora infestans</i>) Vroeëroes (<i>Alternaria solani</i>)	1.25 kg/ha	Voorkomende behandeling: Dien vanaf opkom met tussenposes van 7 tot 10 dae toe in nie minder nie as 500 L water/ha in die voorblomstadium en 1000 L water/ha in die nablomstadium. Vir doeltreffende beheer van vroeë- en laatroes, veral na blom, is dit belangrik dat die plante deeglik bedek moet word. Voeg 'n geskikte benatter teen die aanbevole dosis by.

GEWAS EN TEIKEN	DOSIS	OPMERKINGS
<p>Tamaties</p> <p>Laatroes (<i>Phytophthora infestans</i>)</p> <p>Vroeëroes (<i>Alternaria solani</i>)</p>	1.25 kg/ha	<p>Dien as 'n voorkomende bespuiting net na uitplant toe en voordat die siektesimptome verskyn. Gebruik 300–1500 L water/ha, afhangend van die plantgrootte. Verseker deeglike bedekking van die boonste en onderste blaaroppervlaktes. Herhaal die toediening met tussenposes van 7 tot 10 dae.</p>