

# CITRUS

## PEST MANAGEMENT CHEAT SHEET

For more detailed information on each product, including safe usage, type the product name into the search bar on [our website](#).



PEST	PRODUCT TO USE	ACTIVE INGREDIENT	PRODUCT CATEGORY	RECOMMENDED SPRAY TIME	PRE-HARVEST INTERVAL (PHI) SA MRL
Citrus Thrips, Psylla, Red Scale, Leaf Miner, Aphids	<b>Kohinor® 350 SC</b> Reg. no. L8447 act 36 of 1947 (Harmful)	Imidacloprid	IRAC 4A	This is a soil-drench product. Use before the end of September from the green-tip to white-bud stage	<b>212</b> days PHI as registered
Citrus Thrips, Psylla, Red Scale, Mussel Scale, Circular Purple Scale, African Bollworm, Mealy bug, Orange dog, Psylla, Aphids	<b>Pyrinex® 250 CS</b> Reg. no. L6515 act 36 of 1947 (Caution) <b>Pyrinex® 480 EC</b> Reg. no. L4673 act 36 of 1947 (Harmful)	Chlorpyrifos	IRAC 1B	Apply during the crop's dormant period, before bud swell. Winter spray application only	<b>60</b> days PHI for spray as registered
Red Scale, Soft Green Scale Waxy Scale, African Bollworm, Mealy bug, Orange Dog, Aphids	<b>Methomex® 900 SP</b> Reg. no. L5254 act 36 of 1947 (Toxic)	Methomyl	IRAC 1A	Use early in the season, after harvest of previous crop, and spray at the first signs of pests	<b>60</b> days PHI as for all registered usages
Citrus Thrips, Psylla, Red Scale, Mealy bug, Leaf Miner	<b>Mulan® 20 SP</b> Reg. no. L9167 act 36 of 1947 (Caution)	Acetamiprid	IRAC 4A	Apply from bud swell to 90% pedal fall	<b>150</b> days PHI as registered
Citrus Thrips, Red Scale, Mealy bug	<b>Trivor®</b> Reg. no. L10879 act 36 of 1947 (Caution)	Acetamiprid + Pyriproxyfen	IRAC 4A + 7C	Apply at 90% pedal fall and repeat only once, 14 days later, if needed	<b>84</b> days PHI as registered



**Emergency contacts:**

Griffon Poison Information Centre: +27 82 446 8946  
 Tygerberg Poison Information Centre: +27 861 555 777

**ADAMA offices:**

Cape Town: +27 21 982 1460

Johannesburg +27 11 635 4300  
 99 Jip de Jager Drive, The Vineyards Oce Estate,  
 Simeka House, De Bron, 7530, South Africa

**Disclaimer:** Please read the label information and safety data sheets (SDS) available for download on our [website](#). The information in the SDS was obtained from sources which we believe are reliable. However, the information is provided without any warranty, express or implied, regarding its correctness. The conditions or methods of handling, storage, use or disposal of the product are beyond our control and may be beyond our knowledge.

For this and other reasons, we do not assume responsibility and expressly disclaim liability for loss, damage or expense arising out of or in any way connected with the handling, storage, use or disposal of the product. Each SDS was prepared and is to be used only for the product it pertains



PEST	PRODUCT TO USE	ACTIVE INGREDIENT	PRODUCT CATEGORY	RECOMMENDED SPRAY TIME	PRE-HARVEST INTERVAL (PHI) SA MRL
<b>African Bollworm, False Codling Moth (FCM)</b>	<b>Warlock™ 19,2 EC</b> Reg. no. L9872 act 36 of 1947 (Caution)	Emamectin	IRAC 6	African Bollworm: Apply at first signs of pest during flowering. FCM: Apply in a structured programme towards the end of the season	<b>14</b> days PHI as registered
<b>Citrus Thrips</b>	<b>Klartan® 240 EW</b> Reg. no. L5096 act 36 of 1947 (Caution)	<b>Tau-fluvalinate</b>	IRAC 3A	Apply at first signs of flowering	<b>43</b> days PHI as registered
<b>Citrus Thrips</b>	<b>Makhromectin®</b> Reg. no. L7941 act 36 of 1947 (Toxic)	Abamectin	IRAC 6	Apply between December and April	<b>3</b> days PHI as registered
<b>Fruit Fly (Natal and Mediterranean)</b>	<b>Devipan® 100</b> Reg. no. L4669 act 36 of 1947 (Toxic)	Dichlorvos	IRAC 1B	Apply as soon as the pest is noticed. Should not be used within 21 days of harvesting	<b>21</b> days PHI