



# TOPIK<sup>®</sup>



ADAMA



HERBICIDE

Emulsifiable  
concentrate

MAPP  
18568

H



## Ingredients

- Clodinafop-propargyl:  
240 g/l (22.2% w/w)
- Cloquintocet-mexyl:  
60 g/l (5.6% w/w)



## Crops

A herbicide for the control of wild oats, rough meadowgrass and moderate control of Italian rye-grass in wheat, durum wheat, rye and triticale.



## Scan for Safety Data Sheet

Or use the weblink :  
<https://bit.ly/3gpDJfQ>  
Alternatively, contact your  
supplier.



TRACEABIL-  
ITY PRINTING  
51/58 x 27 mm

1Le

**Danger**

May be fatal if swallowed and enters airways.  
May cause damage to the nervous system through prolonged or repeated exposure.

Very toxic to aquatic life with long lasting effects.  
Repeated exposure may cause skin dryness or cracking.

Contains clodinafop-propargyl and cloquintocet-mexyl. May produce an allergic reaction.

Keep out of reach of children.

Do not breathe vapours/spray

IF SWALLOWED: Immediately call a POISON CENTRE or doctor/physician.

Do NOT induce vomiting.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment comply with the instructions for use

**SAFETY PRECAUTIONS****Operator protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

**WEAR SUITABLE PROTECTIVE GLOVES**

when handling the concentrate or handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

**WASH ALL PROTECTIVE CLOTHING** thoroughly after use, especially the inside of gloves.

**WASH SPLASHES** from skin immediately.

**WASH HANDS AND EXPOSED SKIN** before meals and after work.

**IF YOU FEEL UNWELL**, seek medical advice (show the label where possible).

**Environmental protection**

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from

L12189IPWB\_01

**IMPORTANT INFORMATION**

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE

<b>Crops:</b>	Wheat, durum wheat, rye and triticale
<b>Maximum individual dose:</b>	0.25 litres per ha
<b>Maximum number of applications:</b>	One per crop
<b>Latest time of application:</b>	Wheat – before flag leaf sheath extending stage (GS 41) Durum wheat, rye and triticale – before second node detectable stage (GS 32)

**Other specific restrictions:**

To avoid the build up of resistance, do not apply this product or any other product containing an ACCase inhibitor herbicide more than twice per crop. In addition, do not use this product in mixture or sequence with any other product containing clodinafop-propargyl.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

**This leaflet is part of the approved Product Label.**

---

**TOPIK® (MAPP 18568)** is an emulsifiable concentrate containing 240 g/l (22.2% w/w) clodinafop-propargyl and 60 g/l (5.6% w/w) cloquintocet-mexyl. It is a herbicide for the control of wild oats, rough meadowgrass and moderate control of Italian rye-grass in wheat, durum wheat, rye and triticale.

---

### **Environmental protection**

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards or roads.

### **Storage and disposal**

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

*The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.*

**For advice on medical emergencies, fires or major spills telephone the National Chemical Emergency Centre on 01865 407333.**

---

## **DIRECTIONS FOR USE**

*IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.*

### **RESTRICTIONS**

Do not use on barley or oats.

Do not spray crops under stress or to crops suffering from waterlogging, pest attack, disease or frost.

Do not spray crops undersown with grass mixtures.

Rain within one hour after application may reduce grass-weed control.

Avoid the use of hormone-containing herbicides in mixture or sequence with TOPIK®.

When TOPIK is applied first, leave 7 days before applying hormone herbicides. If mecoprop-p or 2,4-DB containing products are applied first, leave 14 days before TOPIK is applied. If MCPA or 2,4-D containing products are applied first, leave 21 days before TOPIK is applied.

Crops should not be re-entered until spray residues are dry.

## RESISTANCE MANAGEMENT

- Always follow WRAG guidelines for preventing and managing herbicide resistant grass-weeds.
- Do not use TOPIK or any other ACCase inhibitor as the sole means of grass-weed control in successive crops.
- Use grass-weed herbicides with different modes of action throughout the cropping rotation.
- To reduce the risk of developing resistance, applications should be made to young, actively growing weeds.
- Use crop rotation and other cultural control measures to prevent and manage herbicide resistant grass-weeds.
- Use tank/product mixes or sequences of herbicides with different modes of action within individual crops, or successive crops.
- Only apply TOPIK once per crop and do not use in mixture or sequence with any product containing clodinafop-propargyl.
- Monitor weed control effectiveness and investigate any odd patches of poor grass-weed control. If unexplained, contact your agronomist who may consider a resistance test appropriate.

This product contains clodinafop-propargyl which is an ACCase inhibitor, also classified by the Herbicide Resistance Action Committee as 'Group 1'.

Use only as part of a resistance management strategy that includes cultural methods of control and does not use ACCase inhibitors as the sole chemical method of grass-weed control.

Applying a second product containing an ACCase inhibitor to a crop will increase the risk of resistance development; only use a second ACCase inhibitor to control different weeds at a different timing.

Strains of some annual grasses (e.g. black-grass, wild oats and Italian rye-grass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the AHDB, Croplife UK, your distributor, crop adviser or product manufacturer.

## WEEDS CONTROLLED

TOPIK is a foliar-acting grass-weed killer for the control of wild oats and rough meadowgrass which can be used post-emergence on all varieties of wheat, durum wheat, autumn-sown spring wheat, rye and triticale. TOPIK can be mixed with many broad-leaved weed and residual grass-weed herbicides to widen the spectrum of weeds controlled.

TOPIK may also be used for the control of black-grass as part of a black-grass management strategy.

## CROP SPECIFIC INFORMATION

### Weed control

Activity is faster under good growing conditions and may appear relatively slow under cold or dry weather conditions.

TOPIK has no residual activity. Optimum weed control will only be achieved when all grass-weeds have

emerged.

TOPIK does not control broad-leaved weeds and if these are present a specific broad-leaved weed herbicide will be required.

TOPIK can be used on all varieties of winter and spring wheat, autumn-sown spring wheat, durum wheat, rye and triticale.

### **Timing**

Spray in the autumn, winter or spring from one true leaf to before flag leaf sheath extending stage of the crop (wheat) or before second node detectable stage of the crop (rye and triticale). Spraying should be done when the majority of weeds have germinated but before weed competition reduces yield. Dry or cloddy seedbeds favour late germination of grass-weeds and spraying should be delayed. A tank mix with a residual herbicide will control weeds germinating after autumn treatment (see tank mix recommendations).

### **RATE OF USE**

Apply TOPIK at 0.25 litres per hectare alone OR 0.125 litres per hectare when used in tank mix with a suitable adjuvant (see 'Compatibility' section below).

### **FOLLOWING CROPS**

The activity of TOPIK is not affected by soil type, organic matter or straw residues. In the event of crop failure, any broad-leaved crop may be sown, or after an interval of 3 weeks, any cereal may be sown. Following normal harvest of a treated crop, any broad-leaved or cereal crop may be sown.

### **MIXING AND SPRAYING**

#### **Spray volume**

Spray TOPIK in 100 - 200 litres of water per hectare. Where there are dense weed populations or advanced growth stages the water volume should be 200 litres per hectare.

#### **Mixing procedure**

Make sure the sprayer is set to give an even application at the correct volume. Fill the spray tank with half the required volume of water and begin agitation. Add the required amount of TOPIK to the spray tank and allow to disperse before adding any other product. Add the rest of the water and continue to agitate the mixture thoroughly. Always agitate during spraying.

#### **Spray quality**

Apply TOPIK using a conventional fan nozzle producing a spray quality at the finer end of the medium range as defined by the British Crop Protection Council. Do not use pre-orifice and air induction nozzles as these may give reduced control, which in high weed populations can prove unacceptable. A spray pressure of 2-3 bars is recommended.

## COMPATIBILITY

### Use of additives

The maximum dose rate of TOPIK is 0.125 litres per hectare when used in mixture with a suitable additive.

TOPIK must always be applied with an authorised adjuvant from the methylated seed oil, mineral oil or non ionic surfactant classes. Use with approved methylated seed oils is preferred. Always use adjuvants at their recommended rates. Please refer to ADAMA literature for lists of approved adjuvants.

TOPIK +/- Additives - Grass-weed susceptibility

Product(s)	Doses (l/ha)	Weed Growth Stages Controlled		
		Wild oats	Rough meadow-grass	Italian rye-grass**
TOPIK	0.25	1st leaf unfolded to flag leaf ligule visible	1st leaf unfolded to main shoot and 3 tillers	Moderately susceptible 1st leaf unfolded to main shoot and 3 tillers
TOPIK + Additive	0.125*	1st leaf unfolded to flag leaf ligule visible	1st leaf unfolded to main shoot and 3 tillers	Moderately susceptible 1st leaf unfolded to main shoot and 3 tillers

\*For dose see text

\*\*Always use as part of a weed control programme including other products active against ryegrasses

TOPIK can contribute to the control of black-grass as part of a herbicide resistance management strategy, involving mixtures and sequences with herbicides of alternative modes of action.

TOPIK may be tank mixed with a variety of plant protection products providing that the application timing is correct for both TOPIK and the partner(s) in the mixture.

*The products should be added separately to the bulk of water in the spray tank. Continuous agitation should be maintained and the products used immediately after mixing.*

For further information on the approval status of mixture partners, consult ADAMA Agricultural Solutions UK Ltd.

Always read the label of the partner product carefully before use.

---

## COMPANY ADVISORY SECTION

*This section is not part of the Product Label under the Plant Protection Products Regulations and provides additional advice on product use at the discretion of ADAMA Agricultural Solutions UK Ltd.*

### DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Marketed by:

ADAMA Agricultural Solutions UK Ltd

Third Floor East

1410 Arlington Business Park

Theale

Reading

RG7 4SA

Tel: 01635 860555

Technical Helpline: 01635 876622

[www.adama.com](http://www.adama.com)

Email: [ukenquiries@adama.com](mailto:ukenquiries@adama.com)

TOPIK® is a registered trademark of a company of the ADAMA Group.