

STAVENTO[®]

GROUP M4 FUNGICIDE

F

PCS 06685

A fungicide for protection against leaf blotch (*Zymoseptoria tritici*) on wheat and triticale and leaf blotch (*Rhynchosporium secalis*) on barley and triticale.

A suspension concentrate formulation containing 500 g/l (39.5% w/w) folpet. Contains folpet and hexamethyl tetramine: may produce an allergic reaction.



ADAMA



Scan for Safety Data Sheet or use weblink:
<https://bit.ly/32UrxKY>

FOR PROFESSIONAL USE ONLY UFI: P3TJ-WFJ4-9Q09-FXP6

FR R3.0.1 JUL21



5Le

SAFETY INFORMATION

Warning

May cause an allergic skin reaction.
Suspected of causing cancer.
Very toxic to aquatic life with long lasting effects.

Wash hands thoroughly after use.
Wear protective gloves/protective clothing.
Avoid breathing vapours/spray.
Obtain special instructions before use.
IF ON SKIN: wash with plenty of soap and water.
If skin irritation or rash occurs: get medical advice/attention.
Contaminated work clothing should not be allowed out of the workplace.
If exposed or if you feel unwell: call a POISON CENTRE or doctor/physician.

Do not handle until all safety precautions have been read and understood.
Store locked up.
Collect spillage.
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.
To protect aquatic organisms, respect an unsprayed buffer of 5 metres to surface water bodies.
To avoid risks to human health and the environment, comply with the instructions for use.

PCS 06685

L20282IPFR_02

STAVENTO®**PCS 06685****PROTECT FROM FROST
SHAKE WELL BEFORE USE****DIRECTIONS FOR USE:**
full instructions in this leaflet.**Authorised to & marketed by:**

Adama Agricultural Solutions UK Ltd

Third Floor East, 1410 Arlington Business Park, Theale, Reading RG7 4SA

Telephone: +44 (0)1635 860555 | **Technical Helpline:** +44 (0)1635 876622**www.adama.com** | **ukenquiries@adama.com****For advice on medical emergencies, fires or major spills telephone the
UK National Chemical Emergency Centre on +44 (0)1865 407333**

Batch No.: see packaging

SAFETY PRECAUTIONS**Operator Protection**WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS)
AND SUITABLE PROTECTIVE GLOVES when handling
the concentrate or handling contaminated surfaces.

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating or
drinking and after work.IF YOU FEEL UNWELL, seek medical advice immediately
(show the label where possible).**Environmental Protection**Do not contaminate water with the product or its
container. Do not clean application equipment near
surface water. Avoid contamination via drains from
farmyards and roads.**Storage and Disposal**KEEP IN ORIGINAL CONTAINER, tightly closed, in a
safe place.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose (litres of product/ha)	Maximum total dose (litres of product/ ha/crop)	Latest time of application
Winter wheat, spring wheat, winter barley, spring barley and triticale	1.5	3	Before end of heading/ inflorescence fully emerged (GS 59)

BK.R3.0.1.JUL21

L202821PBB_02

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT
WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT
PROTECTION PRODUCTS.**

This leaflet is part of the approved label.

For advice on medical emergencies, fires or major spills telephone
the UK National Chemical Emergency Centre on +44 (0)1865 407333

A fungicide for protection against leaf blotch (*Zymoseptoria tritici*) on wheat and triticale and leaf blotch (*Rhynchosporium secalis*) on barley and triticale.

A suspension concentrate formulation containing 500 g/l (39.5% w/w) folpet.
Contains folpet and hexamethyl tetramine: may produce an allergic reaction.

SAFETY INFORMATION

Warning

May cause an allergic skin reaction.

Suspected of causing cancer.

Very toxic to aquatic life with long lasting effects.

Wash hands thoroughly after use.

Wear protective gloves/protective clothing.

Avoid breathing vapours/spray.

Obtain special instructions before use.

IF ON SKIN: wash with plenty of soap and water.

If skin irritation or rash occurs: get medical advice/attention.

Contaminated work clothing should not be allowed out of the workplace.

If exposed or if you feel unwell: call a POISON CENTRE or doctor/physician.

Do not handle until all safety precautions have been read and understood.

Store locked up.

Collect spillage.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

To protect aquatic organisms, respect an unsprayed buffer of 5 metres to surface water bodies.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS 06685



IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose (litres of product/ha)	Maximum total dose (litres of product/ha/crop)	Latest time of application
Winter wheat, spring wheat, winter barley, spring barley and triticale	1.5	3	Before end of heading/ inflorescence fully emerged (GS 59)

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

SAFETY PRECAUTIONS

Operator Protection

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces. WASH CONCENTRATE from skin or eyes immediately. WASH HANDS AND EXPOSED SKIN before eating or drinking and after work. IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.
DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Resistance

STAVENTO® contains folpet, a protectant fungicide from the phthalimide chemical group with multisite contact activity. For resistance management purposes, its mode of action is classified as M4 under the FRAC* code which indicates low-risk without any signs of resistance or cross-resistance developing.

For current advice on resistance management contact your agronomist or specialist advisor and visit the FRAG-UK website.

*FRAC: Fungicide Resistance Action Committee

Disease control in wheat, barley and triticale

For the reduction of leaf spot (*Zymoseptoria tritici*) in winter and spring wheat and triticale and leaf blotch (*Rhynchosporium secalis*) in winter and spring barley and triticale. STAVENTO may be applied in a programme of 2 suitably timed sprays.

Dose rate

1.5 L/ha per application.

Septoria leaf spot (*Zymoseptoria tritici*) in winter and spring wheat and triticale

As STAVENTO is a protectant fungicide with contact activity, it is important to apply the first spray before the disease becomes established in the crop (normally GS 30-33). A second spray should be timed to protect new growth from disease and therefore needs to be applied prior to further infection. This is normally 3 weeks after the first application but may be sooner under conditions of rapid growth and high disease pressure. Where disease infection or re-infection has occurred prior to application, STAVENTO should be mixed with alternative chemistry to provide curative activity (see Compatibility section).

Leaf blotch (*Rhynchosporium secalis*) in winter and spring barley and triticale

As STAVENTO is a protectant fungicide with contact activity, it is important to apply the first spray before the disease becomes established in the crop (normally GS 30-33). A second spray should be timed to protect new growth from disease infection. This is normally 3 weeks after the first application but may be sooner under conditions of rapid growth and high disease pressure. Where disease infection or re-infection has occurred prior to application, STAVENTO should be mixed with alternative chemistry to provide curative activity (see Compatibility section).

Water volume

Apply STAVENTO in a water volume of 200-400 L/ha. As STAVENTO has contact activity, it is important to achieve a good coverage of the crop when spraying.

Mixing and spraying

For use by tractor mounted/trailed sprayer only.

Before spraying it is important to check all hoses, filters and nozzles, and to ensure that the sprayer is clean and correctly calibrated to give an accurate application at the correct volume. Half-fill sprayer tank with clean water. Add the required amount of STAVENTO to the water and commence agitation. On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely. Add the remainder of the water whilst continuing to agitate the mixture until spraying is complete.

Apply STAVENTO as a medium spray (as defined by the BCPC) in cereals.

Thoroughly clean the sprayer and measuring equipment after using STAVENTO.

Crops should not be re-entered until spray residues are dry.

Compatibility

Consult Adama Agricultural Solutions UK Ltd or your distributor for details on the products which may be mixed with STAVENTO.

Maintain agitation during the mixing process and continue until the spraying is complete. Do not leave in spray tank for long periods without using.

Caution

After spraying a herbicide, sprayer tanks and equipment should be thoroughly cleaned before using STAVENTO.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Authorised to and marketed by:

Adama Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park
Theale, Reading RG7 4SA
Telephone: + 44 (0)1635 860555
Technical Helpline: + 44 (0)1635 876622
www.adama.com
Email: ukenquiries@adama.com

Stavento® is a registered trademark of a company of the Adama Group.
© Adama Agricultural Solutions UK Ltd