

Enzo™ IRON

MAPP 19501

A ready-to-use, pelleted bait containing 2.94% w/w anhydrous ferric phosphate for the control of slug damage in a range of agricultural and horticultural crops.

Rainproof formulation.



ADAMA

For advice on medical emergencies, fires or major spills telephone the National Chemical Emergency Centre on 01865 407333

Adama Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park
Theale, Reading RG7 4SA
Telephone: (01635) 860555
Technical Helpline: (01635) 876622
www.adama.com
Email: ukenquiries@adama.com

To access the Safety Data Sheet for this product scan the QR code or use the weblink:
<https://bit.ly/3dvRO5s>
Alternatively, contact your supplier.



Batch number: see packaging



LDPE
Please recycle this packaging.



Enzo IRON is packaged in high quality MINrinse bags

- Fully recyclable - Accepted by all of the major recycling and waste disposal companies.
- Not classified as hazardous waste - Empty packaging can be treated as a triple rinsed container.
- Easy to store - Packs can be stacked safely due to the unique antislip technology.

20 kg

THE (COSHH) CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH REGULATIONS
MAY APPLY TO THE USE OF THIS PRODUCT AT WORK.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL MOLLUSCICIDE

Crop/situations	Maximum individual dose (kg product/ha)	Maximum total dose
Borage, linseed, mustard, oilseed rape, poppy, soya bean, sunflower	7	28 kg product/ha/year
Celeriac, radish, swede, turnip	7	28 kg product/ha/year
Barley, durum wheat, oats, rye, triticale, wheat	7	28 kg product/ha/year
Forage maize, grain maize, sweetcorn	7	28 kg product/ha/year
Fodder beet, red beet, sugar beet	7	28 kg product/ha/year
Beans without pods - fresh, combining pea, vining pea, dwarf French bean, field bean, runner bean, broad bean - fresh.	7	28 kg product/ha/year
Baby leaf crops, chicory - witloof, endive, lamb's lettuce, land cress, lettuce, purslane, red mustard, rocket, spinach beet	7	28 kg product/ha/year
Broccoli/calabrese, Brussels sprout, cabbage, cauliflower, kale	7	28 kg product/ha/year
Carrot	7	28 kg product/ha/year
Celery	7	28 kg product/ha/year
Leek	7	28 kg product/ha/year
Bulb onion, garlic	7	28 kg product/ha/year
Fennel leaves, globe artichoke, salad onion	7	28 kg product/ha/year
Potato	7	28 kg product/ha/year

Time of application: a one day pre-harvest interval must be observed for all crops. A minimum period of 7 days must elapse between applications for all crops.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

SAFETY PRECAUTIONS

Operator Protection

WEAR SUITABLE PROTECTIVE GLOVES when handling the product or applying by hand-held equipment.

DO NOT EAT, DRINK OR SMOKE, when using this product.

WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental Protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads. Do not empty in to drains, dispose of this material and its container in a safe way.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

Do not use original packaging or empty container for any other purpose.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

DO NOT RE-USE container for any purpose.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Enzo™ IRON is a blue mini pellet containing ferric phosphate which occurs naturally in the environment.

Enzo IRON controls all species of slugs including *Deroceras reticulatum* (grey field slug), *Arion* spp and *Arion vulgaris* (Spanish slug). After the slug has eaten the pellet, feeding will cease to continue resulting in protection of the crop from feeding damage.

RESTRICTIONS OR WARNINGS

Enzo IRON is attractive to slug species that damage plants. Due to the mode of action of Enzo IRON, no excessive slime secretions will be evident on, or around, the crop and the slugs retreat underground to die and dead slugs will not be visible on the soil surface.

Effectiveness should therefore be measured by the decreased feeding damage to the crop.

Controls slugs feeding on the soil surface in crops of vegetables grown outdoors, reducing damage during plant emergence and lower leaf damage. Damage caused by sub-surface activity to underground plant parts such as potatoes, carrots and other roots, tubers, bulbs, corms and rhizomes etc. may not be reduced. Similarly, damage caused to aerial plant parts such as Brussels sprout buds etc. may not be reduced.

PESTS CONTROLLED

For the control of slugs in the crops listed above.

CROP SPECIFIC INFORMATION

Enzo IRON can be applied with a maximum individual application dose of 7 kg/ha up to a maximum total dose of 28 kg/ha. Enzo IRON provides protection during establishment, applying at later growth stages may not provide any benefit.

Time of application

Apply as a broadcast treatment as soon as damage is seen or indicated by slug trapping. For some crops e.g. cereals and oilseed rape, where slug activity is present prior to crop emergence, it is advised to treat the crop before emergence and after seed bed preparation is complete.

Application timing should be based on likely pest presence and the part of the crop attacked by the pest e.g. seeds, seedling plants or harvestable produce.

A repeat application may be required if pest pressure remains high. A further dose of 7 kg/ha may be carried out if there are no pellets remaining on the soil surface from the previous application (typically 1 week later). The maximum total dose that may be applied per year is 28 kg/ha.

RATE OF USE

Maximum individual dose: 7 kg/ha

Maximum total dose: 28 kg/ha of product per year

APPLICATION

This product is suitable for application through the following spreaders (or other comparable machines):

Stocks FanJet Pro 65

Logic TFS

Do not apply using other machinery.

Calibrate all equipment before use.

Take care to ensure granules do not remain lodged within foliage, florets or other parts of the plants.

Apply evenly over the plants to be protected.

Slug Trapping

To establish the need for pellet application on winter wheat or winter oilseed rape, monitor for slug activity. Where slug traps are used, use a foodstuff attractive to slugs e.g. chicken layers' mash which has proven to be particularly effective.

For further information on slug control and damage risk assessment, please refer to the AHDB Information Sheet 04 Winter 2018 (Integrated Slug Control) which can be found at www.ahdb.org.uk/knowledge-library/integrated-slug-control or use the weblink: <http://bit.ly/sluginfo>

Important

- Complete control of slugs is difficult to achieve with any type of molluscicide, so some crop loss may occur especially in wet seasons. Best control is achieved by a programme of good husbandry and properly timed applications of a pellet bait such as Enzo IRON.
- The first application of Enzo IRON is usually made as a preventative measure as specified. Follow-up applications may be required, especially in wet seasons. Assessment by slug trapping will indicate when slugs are active at the soil surface and if conditions are suitable for further Enzo IRON treatment.
- Late application of pellets to broad-leaved plants may result in lodging of the pellets in foliage. Care should be taken to avoid this when making applications to edible crops, e.g. lettuce and cabbage. In cabbage and cauliflower apply prior to hearting or curd formation.
- AVOID application if heavy rain is expected, as this may reduce effectiveness of the pellets.
- Best results are obtained when slugs are actively feeding, generally following a period of light rain in mild conditions. Where a heavy attack is likely to occur, re-apply 2-3 weeks later or whenever slug trapping shows it to be necessary.

Adama Agricultural Solutions UK Ltd

Third Floor East, 1410 Arlington Business Park

Theale, Reading RG7 4SA

Telephone: 01635 860555

Technical Helpline: 01635 876622

www.adama.com

Email: ukenquiries@adama.com

Enzo™ is a trademark of a company of the Adama Group.

Other brand names referred to on this label are trademarks of other manufacturers in which proprietary rights may exist.

© Adama Agricultural Solutions UK Ltd

COMPANY ADVISORY INFORMATION

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.