



TITHE®



ADAMA



HERBICIDES

Emulsifiable concentrate

MAPP 19515

H



Ingredients

- Propaquizafop 100 g/L (9.7% w/w)
- Also contains: Also contains propylene carbonate and hydrocarbons, C10-C13, aromatics, <1% naphthalene.



Crops

A systemic foliar applied herbicide for the post-emergence control of annual and perennial grass weeds in oilseed rape, swedes, turnips, linseed, combining peas, field beans, French dwarf beans, Navy beans, broad beans, potatoes, sugar beet, fodder beet, carrots and bulb onions.



Danger

May be fatal if swallowed and enters airways.
 Causes serious eye irritation.
 Toxic to aquatic life with long lasting effects.
 Repeated exposure may cause skin dryness or cracking.
 Contains propaquizafop. May produce an allergic reaction.
 Keep out of reach of children.
 Avoid breathing vapours/spray.
 Wear protective gloves/protective clothing/eye protection/face protection.
 IF IN EYES: rinse cautiously with water for several minutes.
 Remove contact lenses, if present and easy to do. Continue rinsing.
 IF SWALLOWED: immediately call a poison centre or doctor/physician.
 Do NOT induce vomiting.
 Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.
 To avoid risks to human health and the environment, comply with the instructions for use.



Scan for Safety Data Sheet

Or use the weblink : <https://bit.ly/3UeJsuQ>
Alternatively, contact your supplier.



R 3.0.2 Jan 23

This label is compliant with the Voluntary Initiative Guidance.

NO VARNISH AREA

L1115161PFR_03

5Le



TITHE®



ADAMA



HERBICIDES

Emulsifiable concentrate

MAPP19515



GROUP 1 HERBICIDES

SHAKE WELL BEFORE USE
SAFETY PRECAUTIONS

PROTECT FROM FROST

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable, in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES, RUBBER BOOTS AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND RUBBER BOOTS when applying by hand-held equipment.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

TAKE OFF IMMEDIATELY all contaminated clothing.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

WHEN USING, DO NOT EAT, DRINK OR SMOKE.

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE

Crops: oilseed rape (winter & spring), field bean, broad bean, French dwarf bean, Navy bean, linseed, sugar beet, fodder beet, swede, turnip, potato (maincrop & early), carrot, bulb onion, combining pea.

Maximum individual dose; Maximum number of applications; Latest time of application; Aquatic buffer zone distance and other specific restrictions: full details are given in the information box within the leaflet attached to the product.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

EO_BBDT9TSTT7

Marketed by: Adama Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park,
Theale, Reading, RG7 4SA
Tel: 01635 860555

Technical Helpline: 01635 876622

www.adama.com

Email: ukenquiries@adama.com

To protect aquatic organisms, respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1m of the top of a ditch which is dry at the time of application. DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1m of the top of the bank of a static or flowing waterbody. Aim spray away from water.

This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with HSE's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

For advice on medical emergencies, fires or major spills telephone the National Chemical Emergency Centre on 01865 407333.

R 3.0.2 Jan 23



5 056325 501864

Danger

May be fatal if swallowed and enters airways.

Causes serious eye irritation.

Toxic to aquatic life with long lasting effects.

Repeated exposure may cause skin dryness or cracking.

Contains propaquizafop. May produce an allergic reaction.

Keep out of reach of children.

Avoid breathing vapours/spray.

Wear protective gloves/protective clothing/eye protection/face protection.

IF IN EYES: rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do.

Continue rinsing.

IF SWALLOWED: immediately call a poison centre or doctor/physician.

Do NOT induce vomiting.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



This leaflet is part of the approved label.

TITHE® is an emulsifiable concentrate containing 100 g/l (9.7% w/w) propaquizafop. It also contains propylene carbonate and hydrocarbons, C10-C13, aromatics, <1% naphthalene.

It is a systemic foliar applied herbicide for the post-emergence control of annual and perennial grass weeds in oilseed rape, swedes, turnips, linseed, combining peas, field beans, French dwarf beans, Navy beans, broad beans, potatoes, sugar beet, fodder beet, carrots and bulb onions.

| IMPORTANT INFORMATION | | | | |
|--|---|--------------------------------|---|---------------------------------------|
| FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE | | | | |
| Crops/situations | Maximum individual dose (litres product/ha) | Maximum number of applications | Latest time of application | Aquatic buffer zone distance (metres) |
| Oilseed rape (winter) | 1.5 | One per crop | Before flower buds visible stage and 90 days before harvest | 5 |
| Oilseed rape (spring) | 1.5 | One per crop | Before eight fully expanded leaves stage and 90 days before harvest | 5 |
| Field bean | 1.5 | One per crop | Before flower buds visible stage and 7 weeks before harvest | 5 |
| Broad bean, French dwarf bean, Navy bean | 1.5 | One per crop | Before flower buds visible stage and 7 weeks before harvest | 5 |

| IMPORTANT INFORMATION | | | | |
|--|---|--------------------------------|--|---------------------------------------|
| FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE | | | | |
| Crops/situations | Maximum individual dose (litres product/ha) | Maximum number of applications | Latest time of application | Aquatic buffer zone distance (metres) |
| Linseed | 1.5 | One per crop | Before flower buds visible stage and 16 weeks before harvest | 5 |
| Sugar beet, fodder beet | 1.5 | One per crop | 60 days before harvest | 5 |
| Swede, turnip | 1.5 | One per crop | 30 days before harvest | 5 |
| Potato (maincrop), potato (early) | 1.5 | One per crop | 30 days before harvest | 5 |
| Carrot | 1.5 | One per crop | 30 days before harvest | 5 |
| Bulb onion | 1.5 | One per crop | 30 days before harvest | 5 |
| Combining pea | 1.5 | One per crop | 7 weeks before harvest | 5 |
| <p>Other specific restrictions: To avoid the build-up of resistance do not apply products containing an ACCase inhibitor herbicide more than twice to any crop. In addition, do not use this product in mixture or sequence with any other product containing propanilaprop.</p> <p>READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.</p> | | | | |

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

TITHE is a systemic foliar applied herbicide for the post-emergence control of annual and perennial grass weeds in oilseed rape, swedes, turnips, linseed, combining peas, field beans, dwarf French bean and Navy beans, broad beans, potatoes, sugar beet, fodder beet, carrots and bulb onions.

Weeds must be emerged at the time of application.

RESTRICTIONS OR WARNINGS

TITHE is foliar acting and the dose is therefore independent of soil type.

Avoid overlaps.

Avoid spray drift onto neighbouring crops, especially cereal crops.

Peas and Beans

If TITHE is applied during periods of high temperatures and/or low soil moisture content chlorotic spotting of the crop may result, particularly on combining peas and field beans, but there is no adverse effect on subsequent growth or yield of combining peas.

Carrots and Onions

Crop effects can occur when the couch dose is applied at early growth stages in carrots and onions.

Potatoes

TITHE must not be applied to seed crops.

Crops suffering from frost damage should not be treated.

Qualified Minor Use Recommendation.

TITHE can also be used on dwarf French beans, broad beans and Navy beans as a qualified minor use recommendation as crop safety on these crops is based on limited evidence.

Resistance Warning

This product contains propaquizafop which is an ACCase inhibitor, also classified by the Herbicide Resistance Action Committee as 'Group 1'. Use only as part of a resistance management strategy that includes cultural methods of control and does not use ACCase inhibitors as the sole chemical method of grass-weed control. Applying a second product containing an ACCase inhibitor to a crop will increase the risk of resistance development, only use a second ACCase inhibitor to control different weeds at a different timing.

Strains of some annual grasses (e.g. Wild oats and Italian rye-grass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action group and copies are available from the HGCA, CPA, your distributor, crop advisor or product manufacturer.

Specifically:

- To reduce the risk of developing resistance applications should be made to young, actively growing weeds.
- Use crop rotation and other cultural control measures to prevent and manage herbicide resistant grass weeds.
- Always follow WRAG guidelines for preventing and managing herbicide resistant grass weeds.
- Do not use 'TITHE' or any other ACCase inhibitor as the sole means of grass weed control in successive crops.
- Use grass weed herbicides with different modes of action throughout the cropping rotation.
- Use tank/product mixes or sequences of herbicides with different modes of action within individual crops, or successive crops.
- Monitor weed control effectiveness and investigate any odd patches of poor grass weed control. If unexplained contact your agronomist who may consider a resistance test appropriate.

WEED CONTROL

Speed of kill will be more rapid when weeds are growing actively under warm conditions with adequate moisture. Treatment under cool conditions will give slower activity. In poor conditions use the higher dose.

Weeds germinating after application will NOT be controlled.

Broad-leaved weeds will NOT be controlled.

The following weeds are controlled up to growth stages indicated at the dose indicated.

| Weed | Weed Growth Stage | Dose ⁽¹⁾ Litre product/ha |
|-------------------------|---|---|
| Volunteer barley | Optimum: 2 leaves unfolded to end of tillering. Latest: Stem erect | 0.5 – 1.0 |
| Volunteer wheat and rye | Optimum: 2 leaves unfolded to end of tillering. Latest: Stem erect | 0.7 – 1.0 |
| Wild oats | Optimum: 2 leaves unfolded to early tillering Latest: Stem erect | 0.7 – 1.0 |

| Weed | Weed Growth Stage | Dose ⁽¹⁾ Litre product/ha |
|---|---|---|
| Barley cover crops ⁽²⁾ | Optimum: 2 leaves unfolded to stem erect Latest: 2 nd node detectable | 1.0 – 1.2 |
| Ryegrass (from seed) | Optimum: 2 leaves unfolded to early tillering Latest: Before stem erect | 1.2 – 1.5 |
| Common couch | 3 leaves unfolded (when majority of shoots have emerged and are approx. 15 cm tall) | 1.5 |
| Sterile brome (<i>Bromus sterilis</i>) | Optimum: 2 expanded leaves to fully tillered | 0.7 – 1.0 |

Notes:

(1) Use highest dose specified if weeds are beyond optimum growth stage or under the following conditions:

- Poor growing conditions, e.g. cool temperatures, dry soil
- Overwintered weeds
- Severe weed infestations especially in non-competitive crops e.g. sugar beet, onions and thin crops of oilseed rape

(2) Barley cover crops: spray when risk of wind blow has passed and before there is serious competition with the crop. Use the higher dose of 1.2 L/ha if spraying is late.

Annual Meadowgrass

Growth will be checked at doses of 0.7 – 1 L/ha and severely checked at 1.5 L/ha. These effects will be reduced if annual meadowgrass is beyond 3 leaves unfolded stage at spraying.

Black-grass

TITHE can contribute to the control of black-grass as part of a herbicide resistance management strategy, involving mixtures and sequences with herbicides of alternative modes of action. Where resistant biotypes are present control from TITHE will be unacceptable.

CROP SPECIFIC INFORMATION

TITHE may be applied to the following crops as indicated:

| Crop | Optimum time to commence spraying when crop is at the following GS | Latest application timing (PHI – Pre-Harvest Interval) |
|----------------------|--|--|
| Oilseed Rape, winter | Expanded cotyledons* | Before flower buds visible stage (PHI = 90 days) |
| Oilseed Rape, spring | Expanded cotyledons* | Before 8 fully expanded leaves stage (PHI = 90 days) |
| Swedes, turnips | 2 pairs of leaves | Before weeds are covered by the crop (PHI = 30 days) |
| Linseed | 3 leaves | Before flower buds visible stage (PHI = 16 weeks) |
| Peas, combining | 2 pairs of leaves (3 rd node) | At flower buds visible stage (PHI = 7 weeks) |

| Crop | Optimum time to commence spraying when crop is at the following GS | Latest application timing (PHI – Pre-Harvest Interval) |
|---|--|--|
| Field beans, winter and spring | 2 pairs of leaves | Before flower bud visible stage (PHI = 7 weeks) |
| Broad bean, French dwarf bean and Navy bean | 2 pairs of leaves | Before flower bud visible stage (PHI = 7 weeks) |
| Potatoes (Do not treat seed crops) | 15-20cm high | Before weeds are covered by the crop (PHI = 30 days) |
| Sugar beet, fodder beet | 2 true leaves | Before weeds are covered by the crop (PHI = 60 days) |
| Carrots | 1 true leaf | PHI = 30 days |
| Onions, bulb | Immediately post crook | PHI = 30 days |

*1.5 L/ha (the Common couch dose) must not be applied to crops of winter and spring oilseed rape before the 5 leaf stage of the crop.

FOLLOWING CROPS

If a crop treated with TITHE should fail for any reason, or after normal harvest, the minimum intervals listed must be observed before replanting any of the following crops:

| | |
|--|----------|
| Winter wheat, winter barley | 2 weeks |
| Peas, field beans, maize and winter oilseed rape | 4 weeks |
| Winter oats | 16 weeks |
| Ryegrass | 8 weeks |

MIXING AND SPRAYING

TITHE should be applied as a FINE or MEDIUM spray (as defined by BCPC) in 100-200 litres water per hectare.

Good spray cover is essential for good results.

Use the higher spray volume in dense crop or weed situations.

Half fill the spray tank with clean water and begin agitation. Add required quantity of TITHE to the tank and complete the filling. Continue agitation until spraying is completed. Spray immediately after mixing.

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

Sprayers should be thoroughly cleaned before use and filters and nozzles checked for damage and blockage.

TITHE may be applied by tractor mounted/drawn sprayer. The sprayer should be calibrated to apply 200-250 L/ha as a FINE or MEDIUM spray (as defined by BCPC).

Crops should not be re-entered until spray residues are dry.

For knapsack sprayers:

Half fill the sprayer tank with clean water. Add the measured amount of product, with rinsings, to the sprayer tank and fit the tank lid. Gently shake the sprayer, by rocking, to ensure thorough mixing. Top up the tank with water to the correct level. Re-fit the tank lid and agitate gently before the spray is used. To ensure thorough

mixing.

When used at a walking speed of 1 m/sec to apply a swath of 1 m width, most knapsack sprayers fitted with a Lurmark AN 2.0 or similar nozzle deliver approximately 200 L/ha spray volume (or 10 L per 500 m²).

Table for product amounts applied through a knapsack sprayer

| Rate of product L/ha | Amount product (ml) per L water applied at 200 L/ha water volume |
|----------------------|--|
| 0.5 | 2.5 |
| 0.7 | 3.5 |
| 1.0 | 5.0 |
| 1.2 | 6.0 |
| 1.5 | 7.5 |

COMPANY ADVISORY SECTION

This section is not part of the Product Label under the Plant Protection Products Regulations and provides additional advice on product use at the discretion of ADAMA Agricultural Solutions UK Ltd.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Authorised to & marketed by:

ADAMA Agricultural Solutions UK Ltd

Third Floor East

1410 Arlington Business Park

Theale

Reading

RG7 4SA

Tel: 01635 860555

Technical Helpline: 01635 876622

www.adama.com

Email: ukenquiries@adama.com

TITHE® is a registered trademark of a company of the ADAMA Group.

© ADAMA Agricultural Solutions UK Ltd