



FOX[®]



ADAMA

H



HERBICIDE

Suspension concentrate

MAPP 20626



Ingredients

● Bifenox 480 g/l (40.6% w/w)



Crops

A herbicide recommended for the post-emergence control of a wide range of broad-leaved weeds in winter wheat, winter barley, winter rye, winter triticale and winter oilseed rape.



Warning


Very toxic to aquatic life with long lasting effects.
Contains 1,2-benzisothiazolin-3(2H)-one. May produce an allergic reaction.

Keep out of reach of children.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



L12334IPFR_02



Scan for Safety Data Sheet

Or use the weblink : <https://bit.ly/3M8Rrj5>

Alternatively, contact your supplier.



R 3.1.2 Jul 24

This label is compliant with the Voluntary Initiative Guidance.

NO VARNISH
AREA

5Le





FOX[®]



ADAMA



HERBICIDE

Suspension concentrate

MAPP 20626



SAFETY PRECAUTIONS

Operator Protection

WASH CONCENTRATE from skin and eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

When using do not eat, drink or smoke.

Keep away from food, drink and animal feeding stuffs.

KEEP OUT OF REACH OF CHILDREN.

If you feel unwell, seek medical advice (show label where possible).

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms, respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within the distance specified for the crop to the top of the bank of a static or flowing water body, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water. ALL CROPS WITH A BUFFER ZONE GREATER THAN 5M ARE NOT ELIGIBLE FOR BUFFER ZONE REDUCTION UNDER THE LERAP HORIZONTAL BOOM

GROUP 14 HERBICIDE

SPRAYERS SCHEME.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with HSE's published guidance or the statutory buffer zone as appropriate to the crop must be maintained. NOTE ALL BUFFER ZONES OF MORE THAN 5M ARE NOT REDUCIBLE. The results of the LERAP must be recorded and kept available for three years.

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place. DO NOT RE-USE CONTAINER for any purpose.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

For advice on medical emergencies, fires or major spills telephone the National Chemical Emergency Centre on 01865 407333.

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE

| | | |
|--|---|---|
| Crops: | Wheat (winter) Barley (winter) Rye (winter) Triticale (winter) | Oilseed rape (winter) |
| Maximum individual dose: | 1.5 litres per hectare | 1 litre per hectare |
| Maximum number of applications: | 1 per crop | 1 per crop |
| Latest time of application: | GS 29 (up to the end of tillering) | GS 19 (up to the end of leaf development) |
| Aquatic buffer zone distance | 10 metres | 6 metres |
| Other specific restriction | This product must not be applied by hand-held equipment. | |
| READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS. | | |

ADAMA Agricultural Solutions UK Ltd
Third Floor East,
1410 Arlington Business Park,
Theale, Reading, RG7 4SA
Tel: 01635 860555

Technical Helpline: 01635 876622

www.adama.com Email: ukenquiries@adama.com

SHAKE WELL BEFORE USE
PROTECT FROM FROST



5 056325 505688

R 3.1.2 Jul 24

20_UBDTHE271

Warning

Very toxic to aquatic life with long lasting effects.


Contains 1,2-benzisothiazolin-3(2H)-one. May produce an allergic reaction.

Keep out of reach of children.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



This leaflet is part of the approved label.

FOX® (MAPP 20626) is a suspension concentrate formulation containing 480 g/l (40.6% w/w) bifenox. It is a herbicide recommended for the post-emergence control of a wide range of broad-leaved weeds in winter wheat, winter barley, winter rye, winter triticale and winter oilseed rape.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

GENERAL INFORMATION

FOX® is a selective herbicide absorbed by the foliage, emerging shoots and roots of susceptible weeds. It is to be used to control a wide range of weeds in winter wheat, winter barley, winter rye, winter triticale and winter oilseed rape.

RESTRICTIONS

DO NOT APPLY FOX to any crop suffering stress at the time of application from factors such as herbicide damage, physical stress, nutrient deficiency, waterlogging, cold weather, periods of drought, if rain or frost is expected. Only apply to a dry crop.

Avoid drift onto broad-leaved plants outside the target area.

Do not use on crops intended for use as seed.

Avoid overlapping spray swaths.

Crops should not be re-entered until spray residues are dry.

WEED CONTROL

Cereals:

| Weed | | Weed susceptibility 1.5 L/ha | Maximum weed growth stage |
|------------------------|--------------------------------|---------------------------------|--|
| English name | Latin name | | |
| Cleavers | <i>Galium aparine</i> | MR | Up to 3 whorls, 5 cm diameter. |
| Common field-speedwell | <i>Veronica persica</i> | S | Plants up to 50 mm across/ high or before buds visible, whichever is the earliest. |
| Field pansy | <i>Viola arvensis</i> | S | Plants up to 50 mm across/ high or before buds visible, whichever is the earliest. |
| Field poppy | <i>Papaver rhoeas</i> | S | Plants up to 50 mm across/ high or before buds visible, whichever is the earliest. |
| Forget-me-not | <i>Myosotis arvensis</i> | MS | Plants up to 50 mm across/ high or before flower buds visible, whichever is the earliest. |
| Ivy-leaved speedwell | <i>Veronica hederifolia</i> | S | Plants up to 50 mm across/ high or before buds visible, whichever is the earliest. |
| Red dead-nettle | <i>Lamium purpureum</i> | S | Plants up to 50 mm across/ high or before buds visible, whichever is the earliest. |
| Shepherd's purse | <i>Capsella bursa-pastoris</i> | MS | Plants up to 50 mm across/ high or before buds visible, whichever is the earliest. |

Oilseed rape:

| Weed | | Weed susceptibility 1.0 L/ha | Maximum weed growth stage |
|------------------------|-------------------------|---------------------------------|--|
| English name | Latin name | | |
| Charlock | <i>Sinapis arvensis</i> | MS | Plants up to 6 leaf stage |
| Common field-speedwell | <i>Veronica persica</i> | S | Plants up to 50 mm across/ high or before buds visible, whichever is the earliest. |
| Field pansy | <i>Viola arvensis</i> | MS | Plants up to 6 true leaves. |
| Field poppy | <i>Papaver rhoeas</i> | MS | Plants up to 50 mm across/ high or before buds visible, whichever is the earliest. |

S = Susceptible, complete or near complete kill

MS =Moderately susceptible, effective suppression with or without mortality

MR =Moderately resistant, temporary suppression, the duration and degree depending on the vigour of the crop and weather

SPECIFIC CROP INFORMATION

FOX can safely be used on winter wheat, winter barley, winter rye, winter triticale and winter oilseed rape.

This product is authorised in winter sown crops. Growers choosing to apply this product to winter sown spring varieties should note that crop safety has not been demonstrated in spring varieties. As a result, application of this product to winter sown spring varieties is done so at the growers own commercial risk (and this also applies to unclassified varieties).

RATE OF USE AND VOLUME

Winter cereals:

Apply 1.5 litres per hectare in 200-400 litres water per ha. Use the higher volume where crop or weeds are dense to ensure good coverage. Pressure 1.5-3.5 bar as a MEDIUM quality spray defined by BCPC.

Winter oilseed rape:

Apply 1.0 litres per ha in 200-400 litres water per ha. Use the higher volume where crop or weeds are dense to ensure good coverage. Pressure 1.5-3.5 bar as a MEDIUM quality spray defined by BCPC.

TIMING

Winter cereals:

Make one application from 5 true leaves to before the start of stem elongation (from GS 15 to before GS 30).

Winter oilseed rape:

Only apply between 5-6 true leaves (GS 15-16) and up to the end of leaf development (GS 19). Weeds should be emerged at the time of application.

Ensure that the crop is well waxed and the leaves are dry, when FOX is applied.

If the crop is suffering from stress at the time of application or FOX is applied in adverse weather conditions e.g. frost, high temperatures, wide temperature fluctuations, chlorosis and mottling may occur, these crop effects are normally outgrown and have no effect on the yield.

FOLLOWING CROPS

Following normal harvest of a crop treated with FOX, the following crops may be sown after cultivation to 20cm: cereals, sugar beet, rye-grass, oilseed rape and peas. In the event of crop failure for any reason, only spring cereals, spring oilseed rape or rye-grass may be sown after cultivation to 20cm.

MIXING

Before spraying it is important to check all hoses, filters and nozzles, and to ensure that the sprayer is clean and correctly set to give an even application at the correct volume.

Half fill the spray tank with clean water. Add the required amount of FOX and begin agitation. Add the remainder of the water and continue agitation until spraying is completed.

Wash equipment thoroughly after use. Fill the tank with clean water and leave overnight. Spray out before storage or using other products. Traces of the product may cause damage to susceptible crops sprayed later.

COMPANY ADVISORY INFORMATION

This section is not part of the Product Label under the Plant Protection Products Regulations and provides additional advice on product use at the discretion of ADAMA Agricultural Solutions UK Ltd.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Authorised to & marketed by:

ADAMA Agricultural Solutions UK Ltd

Third Floor East

1410 Arlington Business Park

Theale

Reading

RG7 4SA

Tel: 01635 860555

Technical Helpline: 01635 876622

www.adama.com

Email: ukenquiries@adama.com

FOX® is a registered trademark of a company of the ADAMA Group.

© ADAMA Agricultural Solutions UK Ltd