



CLEAN  
CROP

# INDIGO



ADAMA

H



**HERBICIDE**

**Suspension concentrate**

**MAPP 21185**

SHAKE WELL BEFORE USE

PROTECT FROM FROST



## Ingredients

- Phenmedipham ..... 320 g/l (29.1 % w/w)  
Also contains alcohol C13-iso, ethoxylated.



## Crops

A selective herbicide for post-emergence use on listed annual broad-leaved weeds in sugar beet, fodder beet, red beet, mangels and strawberries.




## Danger

**Causes serious eye damage.**  
**Very toxic to aquatic life with long lasting effects.**  
**Contains 1,2-benzisothiazolin-3-one. May cause an allergic reaction.**

Keep out of reach of children.  
Wear protective gloves, eye protection & face protection.  
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.  
Immediately call a POISON CENTRE or doctor/physician  
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



## Scan for Safety Data Sheet

Or use the weblink : <https://bit.ly/4h7NNEI>  
Alternatively, contact your supplier.



R 3.0 Feb 25

This label is  
compliant with  
the Voluntary  
Initiative  
Guidance.

**NO VARNISH  
AREA**

**5Le**

L125J&IPFR\_01





# INDIGO



ADAMA



## HERBICIDE

Suspension  
concentrate

MAPP 21185



### SAFETY PRECAUTIONS

#### Operator Protection

*Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.*

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when applying by hand-held equipment.

*However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.*

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking, and after work.

DO NOT BREATHE SPRAY.

#### Environmental Protection

Do not contaminate water with product or its container. Do

**GROUP 5 HERBICIDE**

not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

#### Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

*The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.*

For advice on medical emergencies, fires or major spills telephone the National Emergency Centre on 01865 407333.

### IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL/HORTICULTURAL HERBICIDE

Crops:	Sugar beet, fodder beet, red beet, mangels	Strawberries
Maximum individual dose (litres per hectare):	3.0	3.0
Maximum total dose (litres per hectare):	3.0	3.0
Latest time of application:	Before the beet leaves meet between the rows	Before flowering

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

Marketed by: Masstock Arable (UK) Limited,  
Station Road, Andoversford, Cheltenham,  
Gloucestershire, GL54 4LZ  
Tel: 01242 821100

Authorised to:  
ADAMA Agricultural Solutions UK Ltd  
Third Floor East, 1410 Arlington Business Park,  
Theale, Reading, RG7 4SA

UFI: FYWF-42K7-900N-AA7G



5 056325 506340

R 3.0 Feb 25

12531PBB 01\_BBD19T52T7

## Danger

Causes serious eye damage.  
Very toxic to aquatic life with long lasting effects.  
Contains 1,2-benzisothiazolin-3-one. May cause an allergic reaction.

Keep out of reach of children.

Wear protective gloves, eye protection & face protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Immediately call a POISON CENTRE or doctor/physician

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.



To avoid risks to human health and the environment, comply with the instructions for use.

### PROTECT FROM FROST

This leaflet is part of the approved label.

### SHAKE WELL BEFORE USE

**CLEANCROP® INDIGO (MAPP 21185)** is a suspension concentrate formulation containing 320 g/l phenmedipham. It also contains alcohol C13-iso, ethoxylated.

It is a selective herbicide for post-emergence use on listed annual broad-leaved weeds in sugar beet, fodder beet, red beet, mangels and strawberries.

### DIRECTIONS FOR USE

*IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.*

#### RESTRICTIONS

Do not apply to crops suffering from stress due to lime or trace element deficiency, or drought, frost or pest damage.

Do not apply to crops suffering from damage due to an earlier herbicide treatment.

Do not apply where rolling or harrowing has been or will be carried out within 7 days.

Do not apply when temperatures exceed 21°C or in strong sunlight. If treatment is essential, delay spraying until the temperature falls and light intensity reduces.

Do not apply when rain is imminent.

Consult processors before using on crops grown for processing.

Do not apply to strawberries grown under polythene tunnels or cloches.

Avoid overlaps.

Crops should not be re-entered until spray residues are dry.

#### CROP SAFETY

All varieties of sugar beet, fodder beet, red beet and mangels may be safely treated, so long as the crop is not suffering any stress.

See 'Crop Specific Information' section for further details.

## RESISTANCE

This product contains phenmedipham, a Group 5 herbicide, based on the mode of action classification system of the Herbicide Resistance Action Committee. Repeated use of herbicides with the same mode of action can increase the risk of strains of weeds developing resistance to these compounds, leading to poor control. In order to minimise the risk, a strategy for preventing and managing such resistance should be adopted.

Growers are advised to apply products containing herbicides with different modes of action in sequence or tank mix where two or more components are active against the target weeds.

Use the recommended rate of CLEANCROP INDIGO and the correct application timing for the hardest to control weed species present in the field.

The above should be used in conjunction with effective cropping rotation and cultivation techniques e.g. stale seedbed, cultivation and use of non-selective herbicide prior to drilling.

Further guidance on weed resistance management is available from the Herbicide Resistance Action Committee (HRAC) and Weed Resistance Action Group (WRAG). Follow WRAG Guidelines.

## WEED CONTROL

CLEANCROP INDIGO is a contact herbicide absorbed by the leaves of weeds. An even application that wets all weeds with a MEDIUM spray under the right weather conditions and at the correct stage of weed growth is necessary to give good control of susceptible weeds.

As CLEANCROP INDIGO is only recommended post-emergence it is advised to use CLEANCROP INDIGO as part of a weed control programme following a pre-emergence herbicide. Weeds should be actively growing at the time of application. Susceptible weeds will show leaf discoloration and die within 3 to 5 days if growing conditions are good. Less susceptible weeds will take longer to die.

CLEANCROP INDIGO should be applied before susceptible weeds are beyond their recommended timing of control.

	Stage of Development		
	Cotyledons	2 true leaves	4 true leaves
Black-bindweed	R	R	R
Chickweed, Common	S	S	S
Dead-nettle, Red	S	S	MS
Fat-hen	S	S	MS
Groundsel	MS	MS	-
Orache, Common	MR	MR	-
Pansy, Field	MR	MR	MR
Poppy, Common	MR	MR	-

CLEANCROP INDIGO will not control perennial broad-leaved weeds, or grass weeds.

S = Susceptible, complete or near complete kill.

MS = Moderately susceptible, effective suppression, with or without mortality.

MR = Moderately resistant, temporary suppression, the duration and degree depending on the vigour of the crop and weather.

R = Resistant, no useful effect.

### **Tank Mix**

For improved control of less susceptible weeds it is advised to apply CLEANCROP INDIGO in mixture with authorised adjuvant Phase II at 1.0 litre per hectare.

### **CROP SPECIFIC INFORMATION**

#### **Sugar beet, fodder beet, red beet and mangels**

All varieties of sugar beet, fodder beet, red beet and mangels may be treated.

CLEANCROP INDIGO can be applied alone post-emergence of the crop and weed, as a conventional overall application. Apply once the crop has fully expanded cotyledons. Avoid overlapping spray swaths.

#### **Strawberries**

All varieties of strawberries may be treated.

Only apply as a conventional overall application to strawberries (not as a band application).

Apply at any time except from the beginning of flowering to picking. Some slight chlorosis may appear on the leaves which will be quickly outgrown with no effect on fruit.

### **OVERALL AND BAND APPLICATION**

**Rate of use:** 3.0 litres per hectare of CLEANCROP INDIGO

The following doses apply to OVERALL AND BAND APPLICATION. For band application, the doses relate to the area of the sprayed band. The area of field treated by these doses depends on the row spacing and the width of the band.

#### **APPLICATION**

**Volume:** 200 – 240 litres water / hectare

#### **Spray Quality – Overall Application**

CLEANCROP INDIGO should be applied as a MEDIUM quality spray (BCPC definition). Do NOT use a coarse quality spray such as produced by low pressure of “reduced-drift” nozzles, as this may give poor weed control.

#### **Spray Quality – Band Application**

Flat fan nozzles, including Evenspray nozzles, with a fan angle of 80° and having an output of 0.4 litres per minute at 3 bar pressure are particularly suitable for application to an 18 cm band at 240 litres per hectare. Apply the recommended rate and volume of CLEANCROP INDIGO per actual hectare band sprayed, NOT per hectare of crop.

#### **LOW DOSE PROGRAMME – strawberries only**

CLEANCROP INDIGO can be used in a repeat low dose programme of sprays starting when the weeds reach the cotyledon stage.

**Rate of use:** 1.5 litres per hectare

**Volume:** 200-300 litres water per hectare

#### **Timing**

Apply as soon as the largest weeds have reached the two-leaf stage. Apply at intervals of 5-7 days.

#### **Succeeding crops**

Beet crops may be sown at any time following the use of CLEANCROP INDIGO. Three months must elapse before any other crop is sown, and mould-board ploughing to a depth of 15 cm must be carried out first.

## MIXING AND APPLICATION

Half fill the spray tank with clean water. Add the required amount of CLEANCROP INDIGO and begin agitation. Add the remaining volume of water. Continue agitation during spraying.

Ensure that all spraying equipment is thoroughly washed out immediately after use with water and a recognized tank cleaner. Traces of CLEANCROP INDIGO left in the sprayer can cause harm to susceptible crops sprayed later.

---

## COMPANY ADVISORY INFORMATION

*This section is not part of the Product Label under the Plant Protection Products Regulations and provides additional advice on product use at the discretion of ADAMA Agricultural Solutions UK Ltd.*

## DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Marketed by:

Masstock Arable (UK) Limited

Station Road

Andoversford

Cheltenham

Gloucestershire

GL54 4LZ

Tel: 01242 821100

Authorised to:

ADAMA Agricultural Solutions UK Ltd

Third Floor East

1410 Arlington Business Park

Theale

Reading

RG7 4SA

CLEANCROP® is a registered trademark of Origin Enterprises plc.