

# BOGOTA<sup>®</sup>

MAPP 19486

**A growth regulator for the reduction of lodging in cereals.**

A soluble concentrate containing 305 g/l (27.2% w/w) chlormequat and 155 g/l ethephon (13.8% w/w).

Ethephon (2-chloroethylphosphonic acid) is an anticholinesterase organophosphate. Handle with care.



ADAMA



Scan for Safety Data Sheet or use weblink:  
<https://bit.ly/2ALzHdq>

FR R3.1 JUL20



## Warning

**Harmful if swallowed.**

**May cause respiratory irritation.**

Avoid breathing vapours or spray.

Wash hands thoroughly after handling.

Do not eat, drink or smoke when using the product.

Use only outdoors or in a well-ventilated area.

IF SWALLOWED: call a POISON CENTRE or doctor/physician if you feel unwell.

IF INHALED: remove person to fresh air and keep comfortable for breathing.

Call a POISON CENTRE or doctor/physician if you feel unwell.  
Rinse mouth.

Store in a well-ventilated place. Keep container tightly closed.

Store locked up.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

5Le

L70013IPFR\_01

# BOGOTA®

MAPP 19486

**SHAKE WELL BEFORE USE  
PROTECT FROM FROST  
DIRECTIONS FOR USE:**

full instructions in this leaflet.

Batch No.: see packaging

**Adama Agricultural Solutions UK Ltd**

Third Floor East, 1410 Arlington Business Park, Theale, Reading RG7 4SA

**Telephone:** 01635 860555 | **Technical Helpline:** 01635 876622

**www.adama.com | ukenquiries@adama.com**

**For advice on medical emergencies, fires or major spills telephone  
the National Chemical Emergency Centre on 01865 407333**

**The (COSHH) Control of Substances Hazardous to Health  
Regulations may apply to the use of this product at work.**

OPEN HERE ►

## SAFETY PRECAUTIONS

### Operator Protection

This product contains ethephon (2-chloroethylphosphonic acid) an anticholinesterase organophosphorus compound. DO NOT USE if under medical advice NOT to work with such compounds. Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces. However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection. WASH CONCENTRATE and diluted product from skin or eyes immediately. DO NOT BREATHE SPRAY. WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before meals and after work.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

### Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

### Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.  
KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.  
KEEP OUT OF REACH OF CHILDREN.  
DO NOT RE-USE CONTAINER for any purpose.  
WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

## IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS A PLANT GROWTH REGULATOR

| Crop          | Maximum individual dose L/ha | Maximum total dose per crop L/ha | Latest time of application    |
|---------------|------------------------------|----------------------------------|-------------------------------|
| Winter wheat  | 2                            | 2                                | Before flag leaf just visible |
| Winter barley | 2                            | 2                                | Before flag leaf just visible |
| Spring barley | 1.5                          | 1.5                              | Before flag leaf just visible |

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

BK R3.1 JUL20

L70013IPBB\_01

## **This leaflet is part of the approved label.**

For advice on medical emergencies, fires or major spills telephone  
the National Chemical Emergency Centre on 01865 407333

---

### **A growth regulator for the reduction of lodging in cereals.**

A soluble concentrate containing 305 g/l (27.2% w/w) chlormequat and 155 g/l ethephon (13.8% w/w).  
Ethephon (2-chloroethylphosphonic acid) is an anticholinesterase organophosphate. Handle with care.

---

## **Warning**

**Harmful if swallowed.**

**May cause respiratory irritation.**

Avoid breathing vapours or spray.

Wash hands thoroughly after handling.

Do not eat, drink or smoke when using the product.

Use only outdoors or in a well-ventilated area.

IF SWALLOWED: call a POISON CENTRE or doctor/physician if you feel unwell.

IF INHALED: remove person to fresh air and keep comfortable for breathing.

Call a POISON CENTRE or doctor/physician if you feel unwell.

Rinse mouth.

Store in a well-ventilated place. Keep container tightly closed.

Store locked up.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**



# IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS A PLANT GROWTH REGULATOR

| Crop          | Maximum individual dose L/ha | Maximum total dose per crop L/ha | Latest time of application    |
|---------------|------------------------------|----------------------------------|-------------------------------|
| Winter wheat  | 2                            | 2                                | Before flag leaf just visible |
| Winter barley | 2                            | 2                                | Before flag leaf just visible |
| Spring barley | 1.5                          | 1.5                              | Before flag leaf just visible |

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

## SAFETY PRECAUTIONS

### Operator Protection

This product contains ethephon (2-chloroethylphosphonic acid) an anticholinesterase organophosphorus compound. DO NOT USE if under medical advice NOT to work with such compounds.

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

**WEAR SUITABLE PROTECTIVE GLOVES** when handling the concentrate or handling contaminated surfaces. However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

**WASH CONCENTRATE** and diluted product from skin or eyes immediately.

**DO NOT BREATHE SPRAY.**

**WHEN USING DO NOT EAT, DRINK OR SMOKE.**

**WASH HANDS AND EXPOSED SKIN** before meals and after work.

**IF YOU FEEL UNWELL**, seek medical advice (show the label where possible).

### Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

### Storage and Disposal

**KEEP IN ORIGINAL CONTAINER**, tightly closed, in a safe place.

**KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.**

**KEEP OUT OF REACH OF CHILDREN.**

**DO NOT RE-USE CONTAINER** for any purpose.

**WASH OUT CONTAINER THOROUGHLY**, empty washings into spray tank and dispose of safely.

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

BOGOTA® shortens and stiffens the straw of winter barley, winter wheat and spring barley by reducing the length of internodes.

## RESTRICTIONS OR WARNINGS

Do not apply during periods of cold weather, frost or snow or if cold weather has been forecast.

Do not apply to crops under stress during hot weather.

Do not apply in periods of high temperature (more than 21°C) occurring just before, during or shortly after spraying.

Do not apply to wet foliage or if rain is forecast for that day.

Do not apply to crops under any growing stress such as drought, waterlogging, physical damage, pesticide toxicity, pest or disease attack (particularly take-all), poor nutrition or other disorder until the stresses have been fully relieved.

Do not apply within 10 days of a herbicide treatment.

Do not use on spring-sown winter varieties or late-sown crops.

Straw from treated crops must not be used in the growing of horticultural crops.

BOGOTA must not be used in a programme with other products containing ethephon (2-chloroethylphosphonic acid).

Extreme care must be taken to avoid drift onto neighbouring crops.

Care should be taken to avoid overlap of spray swaths.

Crops should not be re-entered until spray residues are dry.

## CROP SPECIFIC INFORMATION

Use BOGOTA as recommended in the table on crops likely to be at risk to lodging i.e. crops growing under fertile conditions. Treat only vigorous crops, adequately fertilised, growing under good soil and weather conditions. Winter wheat and winter barley will normally have been treated with an approved treatment of chlormequat before the first node.

### Cautionary note

Ear emergence may be slightly delayed as a result of straw shortening and some lodging may occur at a late stage. Latent stresses, particularly moisture stress, upon a crop may reduce crop growth and/or lead to secondary tillering. This may be accentuated by BOGOTA.

Seed merchants must be consulted before treating crops intended for seed.

### **Time of application and rate of use**

For best results crops must be actively growing under warm moist conditions.

| Crop   | Application timing<br>The application timings relate to the specified growth stage being found on the majority of tillers | Dose     | Further directions                      |
|--|---|----------|---|
| <b>Winter wheat</b><br>Vigorously growing crops not previously treated with growth regulator, or after chlormequat, but where a high risk of severe early lodging still remains. | During stem elongation from 2nd node to before flag leaf just visible, GS 32-36 inclusive.                                | 2 L/ha   | Apply well before any signs of lodging. |
| <b>Winter barley</b><br>Vigorously growing crops not previously treated with growth regulator or where further protection against lodging is required.                           | During stem elongation from 2nd node to before flag leaf just visible, GS 32-36 inclusive.                                | 2 L/ha   | Apply well before any signs of lodging. |
| <b>Spring barley</b><br>Vigorously growing crops requiring protection against lodging.   | During stem elongation from 2nd node to before flag leaf just visible, GS 32-36 inclusive.                                | 1.5 L/ha | Apply well before any signs of lodging. |

## MIXING AND SPRAYING

### Water volume

Apply the recommended dose as a MEDIUM spray (BCPC) in 225-340 litres of water per hectare using the highest volume in the densest crops.

Apply via established tramlines to minimise production of green side tillers.

### Mixing

Invert the product container several times and shake thoroughly before pouring.

Part-fill the spray tank with clean water and begin agitation. Mix in the required volume of BOGOTA through the top filter or filling device. Top up the tank with water and keep under agitation until sprayed out. Spray immediately after mixing.

### Spray tank cleaning

Thoroughly wash all spraying equipment and measuring equipment with water and a wetting agent using an integrated pressure rinsing device or manually rinsing three times immediately after use.

## DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

### Adama Agricultural Solutions UK Ltd

Third Floor East, 1410 Arlington Business Park

Theale, Reading RG7 4SA

Telephone: 01635 860555

Technical Helpline: 01635 876622

[www.adama.com](http://www.adama.com)

Email: [ukenquiries@adama.com](mailto:ukenquiries@adama.com)

BOGOTA® is a registered trademark of a company of the Adama Group.  
© Adama Agricultural Solutions UK Ltd