CAPTAN 80 WDG

MAPP 16293

For moderate control of apple and pear scab and the reduction of *Gloeosporium* rots of apple.

A water dispersible granule formulation containing 80% w/w captan.



ADAMA



Warning

May cause an allergic skin reaction.
Causes serious eye irritation.
Suspected of causing cancer.

Very toxic to aquatic life.

Harmful to aquatic life with long lasting effects.

Keep out of reach of children.

Wear protective gloves/protective clothing/ eye protection/face protection.

IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

BXFR R3.2.2 NOV19





CAPTAN 80 WDG MAPP 16293 PROTECT FROM FROST **DIRECTIONS FOR USE:**

full instructions in this leaflet Batch No.: see packaging

Adama Agricultural Solutions UK Ltd

Third Floor East, 1410 Arlington Business Park, Theale, Reading RG7 4SA

Telephone: 01635 860555 | **Technical Helpline:** 01635 876622 www.adama.com | ukenquiries@adama.com

For advice on medical emergencies, fires or major spills telephone the National Chemical Emergency Centre on 01865 407333

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

OPEN HERE

SAFETY PRECAUTIONS

Operator Protection

Spray equipment must only be used where the operator's normal working position is within a closed cab on a tractor or a self-propelled sprayer when making airassisted applications to top fruit. Other engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment. WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the product. WEAR SUITABLE PROTECTIVE GLOVES

when handling contaminated surfaces. WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when applying by hand-held equipment.

However, engineering controls in addition to those specified above may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH ANY CONTAMINATION FROM SKIN OR FYES IMMEDIATELY

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

TAKE OFF IMMEDIATELY all contaminated clothina.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible). WHEN USING DO NOT EAT, DRINK OR SMOKE. WASH HANDS before eating and drinking and after work.

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads. To protect aquatic organisms and

non-target plants do not apply on impermeable surfaces such as asphalt, concrete, cobblestones, railway tracks and other situations with a high risk of run-off.

DO NOT ALLOW DIRECT SPRAY from broadcast-air assisted sprayers to fall within 30 m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 5 m of the top of a ditch which is dry at the time of application.



Aim spray away from water. This product aualifies for inclusion within the Local **Environmental Risk** Assessment For Pesticides (LERAP) scheme.

Before each spraying operation from a broadcast air-assisted sprayer, either a LERAP must be carried out in accordance with HSE's published guidance or the statutory buffer zone must be maintained.

The results of the LFRAP must be recorded and kept available for three years.

Storage and Disposal

KEEP IN ORIGINAL CONTAINER. tightly closed, in a safe place. DO NOT RE-USE CONTAINER for any purpose. KEÉP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

97050438C/201910

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS A HORTICULTURAL FUNGICIDE

Crop: Apples and Pears

Maximum individual dose (kg product/ha) Maximum number of applications Latest time of application Other specific restrictions:

Full details are given in the leaflet attached to this product.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.



This leaflet is part of the approved label.

For advice on medical emergencies, fires or major spills telephone the UK National Chemical Emergency Centre on +44 (0)1865 407333

For moderate control of apple and pear scab and the reduction of Gloeosporium rots of apple.

A water dispersible granule formulation containing 80% w/w captan.

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS A HORTICULTURAL FUNGICIDE

Crop Maximum individual dose (kg product/ha) Maximum number of applications of application

Apples and Pears 2.0 10 per year 31 days before harvest

Other specific restrictions:

A minimum interval of 10 days must be observed between applications.

The maximum concentration must not exceed 1.7 g of product per 1 litre water when applying by hand held equipment.

Spray equipment must only be used where the operator's normal working position is within a closed cab on a tractor or on a self-propelled sprayer when making air-assisted applications to top fruit.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.



Warning

May cause an allergic skin reaction.

Causes serious eye irritation.

Suspected of causing cancer. Very toxic to aquatic life.

Harmful to aquatic life with long lasting effects.

Marmitul to aquatic life with long lasting effects

Keep out of reach of children.

Wear protective gloves/protective clothing/eye protection/face protection.

IF ON SKIN: Wash with plenty of soap and water.
IF IN EYES: Rinse cautiously with water for several

minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Dispess of continue rinsing.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS OR WARNINGS

After spraying with alkaline or oil containing products, sprayer tanks and equipment should be thoroughly cleaned before using CAPTAN 80 WDG.

We recommend that you do not spray Bramley's Seedling, Monarch, King Edward (Edward VII), Kidds Late Orange, Red Delicious, Winston Apple or D'Anjou pear as these may be sensitive. However, CAPTAN 80 WDG has been successfully used at the recommended doses on Bramley without crop damage. However, if you choose to use it on Bramley we strongly recommend that you check the susceptibility under your growing conditions by treating a small number of plants in the first instance.

Safety to parasites and predators used in integrated crop management has not been tested and safety cannot be assumed.

RESISTANCE

CAPTAN 80 WDG is a multi-site protectant fungicide. For resistance management purposes, its mode of action is classified as M4 under the FRAC code which indicates low-risk without any signs of resistance or cross-resistance developing.

CAPTAN 80 WDG is well-suited for use in antiresistance fungicide management spraying programmes.

DISEASES CONTROLLED

Apples

Scab (Venturia inaequalis) – moderate control Gloeosporium storage rots – reduction

Pears

Scab/Black spot of pears (Venturia pyrina) – moderate control

CROP SPECIFIC INFORMATION

Time of application

Apples and Pears

For control of apple or pear scab, CAPTAN 80 WDG is most effective when used preventatively as part of a programme of routine sprays beginning at the start of flowering. Commence spraying at bud burst and repeat at 10 to 14 day intervals until early July if scab conditions prevail.

Treatments should be made at 10 day intervals when trees are making rapid growth particularly during rainy conditions.

Apples

For reduction of *Gloeosporium* storage rots, spray in mid to late July and repeat at 2-3 week intervals for as long as orchard conditions permit or until 31 days before picking. Harvest before mid-October. When CAPTAN 80 WDG has been used for post-blossom scab control and heavy infection is not expected, spray in mid-August and repeat 3 weeks later.

We recommend that you do not spray Bramley's Seedling, Monarch, King Edward (Edward VII), Kidds Late Orange, Red Delicious, Winston Apple, or D'Anjou pear as these may be sensitive. However CAPTAN 80 WDG has been successfully used at the recommended doses on Bramley without crop damage. However, if you choose to use it on Bramley we strongly recommend that you check the susceptibility under your growing conditions by treating a small number of plants in the first instance.

RATE OF USE

Apples and Pears

Apply at a rate of 2.0 kg/ha in 500-1,000 L water/ha. A maximum of 10 applications per year may be made. Latest timing is 31 days before harvest.

Apply the recommended dose of 2.0 kg/ha to full height trees typical of UK orchards at full canopy density. Where tree height and/or canopy density is reduced, the dose (and water volume) should be adjusted in accordance with an appropriate dose adjustment scheme. Consult your specialist advisor for further information. Further information on the PACE scheme is available from AHDB Horticulture (formerly HDC), or see the AHDB Horticulture leaflet (Orchard Spraying: Opportunities to reduce rates) available on the AHDB Horticulture website at www.horticulture.ahdb.org.uk

FOLLOWING CROPS

There are no following crops restrictions.

MIXING AND SPRAYING

Before spraying it is important to check all hoses, filters and nozzles, and to ensure that the sprayer is clean and correctly set to give an even application at the correct volume.

Half-fill the sprayer tank with water. Commence agitating the water. Add the required amount of CAPTAN 80 WDG to the water and add the remainder of the water, continuing to agitate the mixture during filling and use. Do not leave mixed spray standing overnight.

Thoroughly clean the sprayer and measuring equipment after using CAPTAN 80 WDG.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Authorisation Holder and Marketed by

Adama Agricultural Solutions UK Ltd Third Floor East, 1410 Arlington Business Park

Theale, Reading RG7 4SA Telephone: 01635 860555 Technical Helpline: 01635 876622

www.adama.com

Email: ukenauiries@adama.com

Other brand names referred to on this label are trademarks of other manufacturers in which proprietary rights may exist.

© Adama Agricultural Solutions UK Ltd