



ADAMA

TOPSAIL®

TOPSAIL™ is for the control of annual grass weeds and annual broad-leaved weeds in winter barley, winter wheat and early and maincrop potatoes.



Crop Information

Crops	Max individual dose (L/ha)	Number of applications	Latest timing
Winter wheat, winter barley	5.0 OR 3.0	1	Pre-emergence Early tillering (GS21)
Potatoes	5.0	1	At emergence (soil rising over emerging potato shoots)

Product Information


Active ingredients:
800g/L prosulfocarb

Formulation:
Emulsifiable Concentrate

Pack size: 10 litres

TOPSAIL™ use recommendations:

4.0-5.0L/ha pre- or 3.0L/ha post-emergence
up until early tillering (BBCH 21)



Application Information

Water volume: 200-400L/ha

Spray quality: Medium

LERAP: B

TOPSAIL™ supported mixes:

	Winter Wheat		Winter Barley	
	Pre-emergence	Peri-emergence	Pre-emergence	Peri-emergence
TOPSAIL™ 2.0-4.0 + Sunfire 0.48		*(max. 2.0 TOPSAIL™)		
TOPSAIL™ 2.0-4.0 + Liberator 0.6		*(max. 2.0 TOPSAIL™)		
TOPSAIL™ 5.0 + Liberator 0.6	**			
TOPSAIL™ 2.0-4.0 + Trooper 4.0		*(max. 2.0 TOPSAIL™)		
TOPSAIL™ 2.0 + Trooper 4.0				**
TOPSAIL™ 2.0-4.0 + FFCT/PICO 1.0		*(max. 2.0 TOPSAIL™)		
TOPSAIL™ 4.0 + Liberator 0.6 + PDM (max.1200g)	**		(max. 2.0 TOPSAIL™)	
TOPSAIL™ 2.0-4.0 + Trooper 4.0 + DFF 60g	**		(max. 2.0 TOPSAIL™)	
AVADEX may be applied in sequence with any of the above if seedbed conditions are as required for triallate. Best advice: apply Topsail mixes pre-em and follow 5 days later with Avadex pre-em (no statutory interval)~				
TOPSAIL™ 3.0 + Liberator 0.6 + Avadex Factor 3.6				

- Not supported
- Supported with caveats
- Supported for efficacy & crop safety

* Transient crop effect may occur post-em where crops are not sufficiently hardened off.

** Seedbed conditions should be good with seed drilled to 30 mm
~ Information inclusive of Avadex Factor