



ADAMA TOPIK®

Controls wild-oats, rough meadow-grass and gives moderate control of Italian rye-grass in wheat, durum wheat, rye and triticale.



Key Benefits

- Wild oats are the most competitive grass weed – more than twice as severe as black-grass
- 1 plant/sqm can reduce winter cereal yields by 1.0 tonne/ha
- Most wild oats germinate in the spring
- Early wild oat control delivers significant yield increase

Crop Information

Wild oats are predominantly spring germinating. Winter cereal crops drilled later may eliminate many of the early autumn germinating wild oats, but the flush of spring germinators must still be controlled.

Smaller spring germinating wild oats offer a greater opportunity for highly effective control.

1 Litre Pack



*Always apply with approved adjuvant oil.

Smaller target weeds can be more effectively controlled using **TOPIK®**, potentially at a lower application rate.

TOPIK® can be tank mixed with SU herbicides when used at the full rate of 125ml/ha plus oil.

DO NOT mix reduced rate **TOPIK®** with an SU. **TOPIK®** should not be mixed with hormone herbicides and must be used in sequence; wait 14 days from application of a hormone herbicide before using **TOPIK®**, or seven days after **TOPIK®** application before using a hormone herbicide.

Product Information

Active ingredients:

240g/L clodinafop-propargyl

Formulation:

Emulsifiable Concentrate

Pack size: 1 litre

Application Information

Water volume: 100-200L/ha

Spray quality: Fine or medium

Aquatic buffer: None

Rate: 0.25L/ha
(or 0.125 with adjuvant)