

## **Key Benefits**

- MIRADOR XTRA® has both contact and systemic activity with protectant properties
- Can be used in all commercial varieties of wheat, barley, oats and rye
- Control a wide range of cereal diseases

## **Crop Information**

Crops	Max individual dose (L/ha)	Max number of applications	Latest time of application
Winter wheat, spring wheat, winter rye, spring rye	1.0	2	Up to and including anthesis complete (GS 69)
Winter barley, spring barley, winter oats, spring oats	1.0	2	Up to and including emergence of ear complete (GS 59)

### **Product Information**

#### **Active ingredients:**

200g/L azoxystrobin 80g/L cyproconazole

### Formulation:

Suspension Concentrate

Pack size: 5 litres

# **Application Information**

Water volume: 200L/ha
Spray quality: Medium
Rainfastness: 1 hour

### **Targets**

### Wheat

- Leaf Spot (Septoria tritici)
- Glume Blotch (Septoria nodorum)
- Yellow Rust (Puccinia striiformis)
- Brown Rust (Puccinia recondita)
- Eyespot (Tapesia yallundae & Tapesia acuformis) - useful reduction
- Can reduce the severity of Take-all (Gaeumannomyces graminis var. Tritici)
- When used to control other diseases, MIRADOR XTRA® will also control wheat powdery mildew

### Oats

- Crown Rust (Puccinia coronata)
- Powdery Mildew (Erysiphe graminis f.sp. avenae)

#### Rve

- Brown Rust (Puccinia recondita)
- Leaf Blotch (Rhynchosporium secalis) moderate control
- Powdery Mildew (Erysiphe graminis)

### Barley

- Leaf Blotch (Rhynchosporium secalis) - moderate control
- Net Blotch (Pyrenophora teres)
- Powdery Mildew (Erysiphe graminis hordei)
- Brown Rust (Puccinia hordei)
- Eyespot (Tapesia yallundae & Tapesia acuformis) - useful reduction
- Can reduce the severity of Take-all (Gaeumannomyces graminis var. Tritici)