



BLANCO™



ADAMA

H



HERBICIDES

Capsule suspension

MAPP
16704



Ingredients

- Clomazone:
360 g/l (30.5% w/w)



Crops

A residual herbicide for the control of broad-leaved weeds in winter and spring oilseed rape, winter and spring field beans, combining peas, vining peas and potatoes.



Scan for Safety Data Sheet

Or use the weblink:
<https://bit.ly/3C9ZyaC>
Alternatively, contact your supplier.



TRACEABILITY PRINTING
51/58 × 27 mm

1Le

GROUP 13 HERBICIDES

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

TO PROTECT NON-TARGET PLANTS, RESPECT AN UNTREATED BUFFER ZONE OF 5 METRES TO NON-CROP LAND.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS

KEEP IN ORIGINAL CONTAINER, TIGHTLY CLOSED, IN A SAFE PLACE.

EMPTY CONTAINER COMPLETELY and dispose of safely.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

For advice on medical emergencies, fires or major spills telephone the National Chemical Emergency Centre on 01865 407333.

ADAMA Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park,
Theale, Reading RG7 4SA
Tel: 01635 860555

Technical Helpline: 01635 876622
www.adama.com
Email: ukenquiries@adama.com

This label is compliant with the
Voluntary Initiative Guidance.





Warning

Very toxic to aquatic life with long lasting effects.

Keep out of reach of children.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

**SHAKE WELL BEFORE USE
PROTECT FROM FROST**

Open here



SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate or contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH CONCENTRATE from skin or eyes immediately.

DO NOT BREATHE SPRAY.

WHEN USING DO NOT EAT DRINK OR SMOKE

Wash all protective clothing after use.

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE

Crop	Maximum individual dose (L product/ha)	Maximum number of treatments (per crop)	Latest time of application
Winter oilseed rape Winter field beans	0.33	1	Pre-crop emergence
Spring combining peas & vining peas Spring field beans Spring oilseed rape	0.25	1	Pre-crop emergence
Potato	0.25	1	Pre-crop emergence after last ridging ± 7 days before cracking.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE.
FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

L11356IPWB_06

This leaflet is part of the approved label.

BLANCO™ is a capsule suspension containing 360 g/L (30.5% w/w) clomazone. It is a residual herbicide for the control of broad-leaved weeds in winter and spring oilseed rape, winter and spring field beans, combining peas, vining peas and potatoes.

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

TO PROTECT NON-TARGET PLANTS, RESPECT AN UNTREATED BUFFER ZONE OF 5 METRES TO NON-CROP LAND.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS

KEEP IN ORIGINAL CONTAINER, TIGHTLY CLOSED, IN A SAFE PLACE.

EMPTY CONTAINER COMPLETELY and dispose of safely.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

For advice on medical emergencies, fires or major spills telephone the National Chemical Emergency Centre on 01865 407333.

DIRECTIONS FOR USE

***IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.*

Blanco™ is a capsule suspension formulation containing clomazone for use as a residual herbicide in all varieties of winter and spring oilseed rape, winter and spring field beans, combining peas, vining peas and potatoes.

Blanco is recommended as a pre-emergence herbicide. Any plants emerged at the time of application may be damaged.

Blanco acts by inhibition of the biosynthesis of photosynthetic pigments leading to bleaching symptoms on susceptible weed species. Blanco is absorbed by the roots and shoots of emerging susceptible species. Susceptible weeds emerge but are lacking pigmentation and plant death occurs shortly afterwards.

RESTRICTIONS OR WARNINGS

Do not apply to broadcast crops.

Do not use on compacted soils or soils with poor structure which may then be liable to water logging.

Do not use Blanco on sands or very light soils. Do not use on soils with more than 10% organic matter.

Ensure seed is covered by a minimum of 20mm of soil.

Some transient crop bleaching can occur under certain climatic conditions and can be severe if heavy rainfall follows application. This transient crop effect is rapidly outgrown and usually has no effect on crop yield.

In prolonged, dry periods following application or if applications are made to dry, cloddy seed beds, reduced levels of weed control may occur.

Do not overlap spray swaths. Severe crop damage may occur on field beans. Where damage does occur, this is generally transient and should not affect crop yield.

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

Spring and winter oilseed rape – chlorotic symptoms may be evident at the early establishment phase of the crop from the cotyledon stage up to early stem extension (GS 10-32) with no permanent detrimental effect on crop health or development.

The safety of Blanco to potato daughter tubers has not been tested – consult the manufacturer or your advisor before using on seed crops.

CONSULT PROCESSORS BEFORE USING ON CROPS FOR PROCESSING.

Contract agents should be consulted before using on crops grown for seed.

There is a risk to adjacent crops and non-crop vegetation from the volatilisation of Blanco. This may result in damage/bleaching to neighbouring vegetation. If effects are seen, they must be reported to your crop protection distributor or adviser.

The risk emanating from volatilisation has been addressed via a stewardship campaign and a user education program, the details of which have been reviewed and agreed by CRD.

RESISTANCE

Blanco contains clomazone, a Group 13 herbicide based on the mode of action classification system of the Herbicide Resistance Action Committee. Repeated use of herbicides with the same mode of action can increase the risk of strains of weeds developing resistance to these compounds, leading to poor control. In order to minimise the risk, a strategy for preventing and managing such resistance should be adopted

Growers are advised to apply products containing herbicides with different modes of action in sequence or tank mix where two or more components are active against the target weeds.

Use the recommended rate of Blanco and the correct application timing. The above should be used in conjunction with effective cropping rotation and cultivation techniques e.g. stale seedbed, cultivation and use of non-selective herbicide prior to drilling.

Further guidance on weed resistance management is available from the Herbicide Resistance Action Committee (HRAC) and Weed Resistance Action Group (WRAG). Follow WRAG Guidelines.

WEEDS CONTROLLED

For optimum efficacy seedbeds should be firm, level and free of clods, loose and puffy seedbeds should be consolidated before spraying. The following weeds are controlled when Blanco is applied pre-emergence to a moist, firm seedbed free from clods.

Winter oilseed rape and winter field beans

- 0.33 L/ha

Pre-emergence application - susceptible

Cleavers

Common chickweed#

Common field-speedwell

Field forget-me-not

Red dead-nettle

Scentless mayweed#

Shepherd's purse

moderately susceptible in winter field beans

Season-long control of weeds may not occur.

Spring oilseed rape, spring field beans, combining peas, vining peas and potatoes

- 0.25 L/ha

Pre-emergence application - susceptible

Cleavers ~

Common chickweed ~

Common field speed-well

Red dead-nettle

Redshank

Scarlet pimpernel

Shepherd's purse

~ Moderately susceptible in peas

Pre-emergence application - moderately susceptible

Black bind-weed

Pre-emergence application - moderately resistant

Annual meadowgrass

Fat-hen

CROP SPECIFIC INFORMATION

Blanco can be used on all varieties of winter and spring oilseed rape, winter and spring beans, combining peas, vining peas and potatoes.

It is important for crop safety, that the seed is covered by a minimum of 20mm of soil. Direct drilled crops should be harrowed across slits to cover the seed before spraying. Where direct drilled crops are treated, then ploughing and cultivating to a depth of 15cm must take place before the planting of any subsequent crop.

TIMING

Winter and spring oilseed rape, winter and spring beans, combining peas and vining peas

Blanco should be applied as soon as possible following sowing of the crop and before emergence of both the crop and target weeds.

Potatoes

Blanco should be applied following the last ridging and before soil cracking (i.e. approximately 7 days before expected potato emergence).

RATE OF USE

Winter oilseed rape and winter field beans - 0.33 L/ha

Spring oilseed rape, spring field beans, combining peas, vining peas and potatoes - 0.25 L/ha.

WATER VOLUME AND SPRAY QUALITY

Apply at in 150-300 L water/ha as a COARSE spray as defined by BCPC.

FOLLOWING CROPS

Autumn treated crops

Following normal harvest of an autumn treated crop, cereals, oilseed rape, field beans, combining peas, potatoes, maize, turnip, linseed and sugar beet may be sown. Prior to planting, soil should be ploughed and cultivated to a minimum depth of 15cm.

Spring treated crops

Following normal harvest of a spring treated crop, cereals, oilseed rape, field beans, combining peas, potatoes, maize, turnip, linseed and sugar beet may be sown. Prior to planting, soil should be ploughed and cultivated to a minimum depth of 15cm.

CROP FAILURE

Autumn treated crops

In the autumn following crop failure, for any reason, only winter cereals and winter beans may be sown, provided at least six weeks has elapsed since application. In the spring following crop failure only spring cereals, combining peas, field beans, potatoes, maize, turnip, onion, carrots and linseed may be sown.

Prior to re-sowing spring cereals, maize, turnip, onion, carrots and linseed, a period of at least 7 months should have elapsed since application.

Prior to re-sowing combining peas, field beans and potatoes a period of at least 6 weeks should have elapsed since application.

Prior to re-sowing any of the listed replacement crops, soil should be ploughed and cultivated to a minimum depth of 25cm.

Spring treated crops

In the spring following crop failure, only combining peas, field beans, potatoes, maize, dwarf French beans, carrots, turnip, sugar beet, onion, and linseed may be sown.

Prior to re-sowing combining peas, field beans, potatoes, carrots, turnip, sugar beet, onion and linseed a period of at least six weeks should have elapsed since application. Prior to re-sowing maize and dwarf French beans a period of at least nine weeks should have elapsed since application.

For all spring replacement crops, soil should be ploughed and cultivated to a minimum depth of 25cm.

MIXING AND SPRAYING

Shake well before use.

Half fill the spray tank with clean water and begin agitation.

Add the required quantity of Blanco and add the remainder of the water.

Rinse empty containers thoroughly and add rinsings to the spray tank.

Once the spray solution has been prepared for use it should be used immediately and not left standing in the spray tank.

Wash out sprayer thoroughly after use.

Crops should not be re-entered until spray residues are dry.

If tank rinse nozzles are fitted: Add clean water to the spray tank until it is approximately a quarter (25%) full then mix in a cleaning agent recommended for sprayer decontamination at the correct dilution

rate for the volume of liquid in the spray tank. No further water needs to be added. Use the tank rinsing nozzles and recirculate for 2 minutes and spray out as described in Disposal of Washings (below).

If tank rinse nozzles are not fitted: Add clean water to the spray tank until it is approximately a quarter (25%) full then mix in a cleaning agent recommended for sprayer decontamination at the correct dilution rate for the full volume of liquid in the spray tank. Fill the sprayer tank to the brim and then recirculate for 2 minutes and spray out as described in Disposal of Washings (below).

Disposal of washings

Spray equipment should not be drained or flushed onto land planted with or intended for planting with trees or crops other than oilseed rape, potatoes, beans and peas.

COMPANY ADVISORY SECTION

This section is not part of the Product Label under Regulation 1107/2009 (GB/NI) and provides additional advice on product use.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Marketed by:

ADAMA Agricultural Solutions UK Ltd

Third Floor East

1410 Arlington Business Park

Theale

Reading

RG7 4SA

Tel: 01635 860555

Technical Helpline: 01635 876622

www.adama.com

Email: ukenquiries@adama.com

Blanco™ is a trademark of a company of the ADAMA Group.

© ADAMA Agricultural Solutions UK Ltd

SAMPLE