



PIOLI™



ADAMA



FUNGICIDE

Emulsifiable concentrate

PCS 07603

FOR PROFESSIONAL USE ONLY

PROTECT FROM FROST

SHAKE WELL BEFORE USE



Ingredients

● Fluxapyroxad 62.5 g/litre (6% w/w)



Crops

A fungicide for use in winter and spring wheat (including durum wheat), winter and spring barley, oats, rye and triticale.



SAFETY INFORMATION

Warning

Causes serious eye irritation.

Harmful if inhaled.

May cause respiratory irritation.

May cause harm to breast-fed children.

Toxic to aquatic life with long lasting effects.

Contains 2-ethylhexyl (2S) -2-hydroxy propionate and benzyl alcohol. May cause allergic reaction.

Contains N,N-Dimethyldodecan-1-amide.

If medical advice is needed, have product container or label at hand.

Use only outdoors or in a well-ventilated area.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do, continue rinsing.

IF exposed or concerned: please call a POISON CENTRE or doctor/physician.

Store in a well-ventilated place.

Keep container tightly closed.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed, empty clean containers which can be disposed of as non-hazardous.

To avoid risks to human health and the environment, comply with the instructions for use.



PCS 07603

L30827IPFR_01



Scan for Safety Data Sheet

Or use the weblink : <https://bit.ly/3P2vpE0>

Alternatively, contact your supplier.



R 3.0 Feb 26

NO VARNISH AREA

5Le

Product name	PIOLI™		
Country	IE		
Package size	5L		
Label code	L30827IPFR_01		
Label dimensions (w x h)	170mm x 150mm		
Label date	19:00:55 26.02.26	Name	
Smallest font on page	8.01 pt	Status	
		Template	Rect_5L_ILang_NA_UK-IE.indt



PIOLI™



ADAMA

FUNGICIDE

Emulsifiable concentrate

PCS 07603

FOR PROFESSIONAL USE ONLY

PROTECT FROM FROST

SHAKE WELL BEFORE USE

Ingredients

● Fluxapyroxad 62.5 g/litre (6% w/w)

Crops

A fungicide for use in winter and spring wheat (including durum wheat), winter and spring barley, oats, rye and triticale.

IMPORTANT INFORMATION FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE	
Crops:	Winter & spring wheat, winter & spring barley, durum wheat, triticale, rye, oats
Maximum individual dose:	2 litres product per hectare
Maximum number of applications:	2 per crop
Maximum total dose	4 litres per hectare
Latest time of application:	Up to and including flowering (anthesis) just complete (GS 69)
Other specific restrictions: A maximum of 2 foliar applications of product(s) containing SDHIs can be applied to any cereal crop. Apply SDHI fungicides always in mixtures. The mixture partner: ● should provide satisfactory disease control when used alone on the target disease ● must have a different mode of action	
READ ALL PRECAUTIONS BEFORE USE.	

Marketed by:
ADAMA Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park,
Theale, Reading, RG7 4SA
Tel: +44 (0)1635 860555
Technical Helpline:
+44 (0)1635 876622
www.adama.com
Email: ukenquiries@adama.com

Distributed in Ireland by:
Croplink (Ireland) Ltd,
Springhill, Carrigtwohill,
Co. Cork, T45 F685,
Tel: 021 4882806

Authorised to: BASF plc,
4th and 5th floors, 2 Stockport Exchange, Railway Road,
Stockport, SK1 3GG

Scan for Safety Data Sheet
Or use the weblink : <https://bit.ly/3P2vpE0>
Alternatively, contact your supplier.



R.3.0 Feb 26

UFI: YG2P-3D5K-DV08-W3DV



SAFETY PRECAUTIONS

Operator protection

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS before meals and after work.

IF SWALLOWED, seek medical advice immediately and show this container or label.

Environmental protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

Storage and disposal

DO NOT RE-USE CONTAINER for any purpose.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

STORE IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

Store in a suitable pesticide store. Keep dry and protect from frost.

In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital on 01 809 2166 or 01 837 9964 (8am - 10pm 7 days a week) or the UK National Chemical Emergency Centre on +44 (0)1865 407333 (24 hours, 7 days a week).

SAFETY INFORMATION

Warning

Causes serious eye irritation.

Harmful if inhaled.

May cause respiratory irritation.

May cause harm to breast-fed children.

Toxic to aquatic life with long lasting effects.

Contains 2-ethylhexyl (2S) -2-hydroxy propionate and benzyl alcohol. May cause allergic reaction.

Contains N,N-Dimethyldecane-1-amide.

If medical advice is needed, have product container or label at hand.

Use only outdoors or in a well-ventilated area.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do, continue rinsing.

IF exposed or concerned: please call a POISON CENTRE or doctor/physician.

Store in a well-ventilated place.

Keep container tightly closed.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed, empty clean containers which can be disposed of as non-hazardous.

To avoid risks to human health and the environment, comply with the instructions for use.



L308270CFR_01

GROUP 7 FUNGICIDE

4x5Le

Product name	PIOLI™		
Country	IE		
Package size	5L		
Label code	L308270CFR_01	Name	ADAMA
Label dimensions (w x h)	300mm x 165mm	Status	
Label date	19.02.26.02.26	Template	Rect_SLT_ILang_NA_UK-IE.indt
Smallest font on page	4.3 pt		



FUNGICIDE

Emulsifiable concentrate

PCS 07603

GROUP 7 FUNGICIDE

L30827IPBB_01

SAFETY PRECAUTIONS

Operator protection

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS before meals and after work.

IF SWALLOWED, seek medical advice immediately and show this container or label.

Environmental protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

Storage and disposal

DO NOT RE-USE CONTAINER for any purpose.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. KEEP OUT OF REACH OF CHILDREN.

STORE IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

Store in a suitable pesticide store. Keep dry and protect from frost.

In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital on 01 809 2166 or 01 837 9964 (8am - 10pm 7 days a week) or the UK National Chemical Emergency Centre on +44 (0)1865 407333 (24 hours, 7 days a week).

IMPORTANT INFORMATION FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE	
Crops:	Winter & spring wheat, winter & spring barley, durum wheat, triticale, rye, oats
Maximum individual dose:	2 litres product per hectare
Maximum number of applications:	2 per crop
Maximum total dose	4 litres per hectare
Latest time of application:	Up to and including flowering (anthesis) just complete (GS 69)
Other specific restrictions:	
A maximum of 2 foliar applications of product(s) containing SDHIs can be applied to any cereal crop.	
Apply SDHI fungicides always in mixtures.	
The mixture partner:	
<ul style="list-style-type: none"> ● should provide satisfactory disease control when used alone on the target disease ● must have a different mode of action 	
READ ALL PRECAUTIONS BEFORE USE.	

Marketed by:

ADAMA Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park,
Theale, Reading, RG7 4SA
Tel: +44 (0)1635 860555

Technical Helpline:

+44 (0)1635 876622
www.adama.com
Email: ukenquiries@adama.com

Distributed in Ireland by:

Croplink (Ireland) Ltd,
Springhill, Carrigtwohill,
Co. Cork, T45 F685,
Tel: 021 4882806

Authorised to: BASF plc,
4th and 5th floors, 2 Stockport
Exchange, Railway Road,
Stockport, SK1 3GG

UFI: YQ2P-3D5K-DV08-W3DV



5 056325 506555

R 3.0 Feb 26

Product name	PIOLI™		
Country	IE		
Package size	5L		
Label code	L30827IPBB_01		
Label dimensions (w x h)	170mm x 144mm		
Label date	19.03.25 26.02.26	Name	
Smallest font on page	5.59 pt	Status	
		Template	Rect_SLT_ILang_NA_UK-IE.indt



PIOLI™



ADAMA



FUNGICIDE

Emulsifiable concentrate

PCS 07603

GROUP 7 FUNGICIDE



SAFETY PRECAUTIONS

Operator protection

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS before meals and after work.

IF SWALLOWED, seek medical advice immediately and show this container or label.

Environmental protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

Storage and disposal

DO NOT RE-USE CONTAINER for any purpose.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. KEEP OUT OF REACH OF CHILDREN.

STORE IN ORIGINAL CONTAINER, tightly closed, in a safe place. RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

Store in a suitable pesticide store. Keep dry and protect from frost.

In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital on 01 809 2166 or 01 837 9964 (8am - 10pm 7 days a week) or the UK National Chemical Emergency Centre on +44 (0)1865 407333 (24 hours, 7 days a week).

L30827IPBB_01

IMPORTANT INFORMATION FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE	
Crops:	Winter & spring wheat, winter & spring barley, durum wheat, triticale, rye, oats
Maximum individual dose:	2 litres product per hectare
Maximum number of applications:	2 per crop
Maximum total dose	4 litres per hectare
Latest time of application:	Up to and including flowering (anthesis) just complete (GS 69)
Other specific restrictions: A maximum of 2 foliar applications of product(s) containing SDHIs can be applied to any cereal crop. Apply SDHI fungicides always in mixtures. The mixture partner: <ul style="list-style-type: none"> ● should provide satisfactory disease control when used alone on the target disease ● must have a different mode of action 	
READ ALL PRECAUTIONS BEFORE USE.	

Marketed by:
ADAMA Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park,
Theale, Reading, RG7 4SA
Tel: +44 (0)1635 860555

Technical Helpline:
+44 (0)1635 876622
www.adama.com
Email: ukenquiries@adama.com

Distributed in Ireland by:
Croplink (Ireland) Ltd,
Springhill, Carrigtwohill,
Co. Cork, T45 P685,
Tel: 021 4882806

Authorised to: BASF plc,
4th and 5th floors, 2 Stockport
Exchange, Railway Road,
Stockport, SK1 3GG

UFI: YQ2P-3D5K-DV08-W3DV



R 3.0 Feb 26



Open here

Product name	PIOLI™		
Country	IE		
Package size	5L		
Label code	L30827IPBB_01		
Label dimensions (w x h)	145mm x 150mm		
Label date	19:00:55 26.02.26	Name	
Smallest font on page	5.54 pt	Status	
		Template	Rect_5LT_1Lang_NA_UK-IE.indt

SAFETY INFORMATION

Warning

Causes serious eye irritation.

Harmful if inhaled.

May cause respiratory irritation.

May cause harm to breast-fed children.

Toxic to aquatic life with long lasting effects.

Contains 2-ethylhexyl (2S) -2-hydroxy propionate and benzyl alcohol. May cause allergic reaction.

Contains N,N-Dimethyldecane-1-amide.

If medical advice is needed, have product container or label at hand.

Use only outdoors or in a well-ventilated area.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do, continue rinsing.

IF exposed or concerned: please call a POISON CENTRE or doctor/physician.

Store in a well-ventilated place.

Keep container tightly closed.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed, empty clean containers which can be disposed of as non-hazardous.

To avoid risks to human health and the environment, comply with the instructions for use.



PCS 07603

PROTECT FROM FROST

SHAKE WELL BEFORE USE

This leaflet is part of the approved label.

PIOLI™ (PCS 07603) is an emulsifiable concentrate containing 62.5 g/litre (6 % w/w) fluxapyroxad. It is a fungicide for use in winter and spring wheat (including durum wheat), winter and spring barley, oats, rye and triticale.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS OR WARNINGS

For professional use only.

For use by tractor mounted/trailed sprayers only.

Avoid spray drift on to neighbouring crops

After spraying, wash out sprayer thoroughly according to manufacturer's guideline and dispose of washings and clean container according to local authority guidelines.

Crops should not be re-entered until spray residues are dry.

RESISTANCE

In order to minimise risk of resistance development, PIOLI™ should always be used in mixture with products with a different mode of action at a dose that will give robust control.

PIOLI should be used in accordance with the instructions for use for the target diseases at the specific growth stages indicated.

Use PIOLI as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control.

Product name	PIOLI™		
Country	IE		
Package size	5L		
Label code			
Label dimensions (w x h)	145mm x 150mm	Name	
Label date	19:00:55 26.02.26	Status	
Smallest font on page	6.89 pt	Template	Rect_SLT_1Lang_NA_UK-IE.indt

For further advice on resistance management contact your agronomist or specialist advisor and visit the FRAG-UK website.

PIOLI must always be tank mixed with another fungicide(s) from different cross resistance group(s)/different mode(s) of action, which when used alone, gives effective control of the same target disease(s). Products should be applied at robust rates. In any tank mix the SDHI should be applied in a balanced mixture

Isolates of *Septoria* leaf blotch with reduced sensitivity to SDHI fungicides have been detected.

Mixtures of two or more SDHI fungicides do not provide an anti-resistance strategy. Each application of such a mixture counts as one SDHI application.

DISEASES CONTROLLED

PIOLI is a fungicide with protectant and systemic properties for use in winter and spring wheat, durum wheat, winter and spring barley, oats, rye and triticale for disease control as summarised below.

	Wheat	Winter barley	Spring barley	Oats	Rye	Triticale
Septoria leaf blotch (<i>Zymoseptoria tritici</i>)	MC	-	-	-	-	MC
Brown rust	C	C	C	-	C	C
Yellow rust	MC	MC	MC	-	MC	MC
Crown rust	-	-	-	C	-	-
Net blotch	-	MC	MC	-	-	-
<i>Rhynchosporium secalis</i> (leaf scald)	-	MC	MC	-	MC	-
Powdery mildew	R	R	R	R	R	R
Tan spot	R	-	-	-	-	-

C = Full Control

MC = Moderate Control

R = Reduction

CROP SPECIFIC INFORMATION

Crops

For use in all varieties of winter and spring wheat, durum wheat, winter and spring barley, oats, rye and triticale.

Time of Application

Apply PIOLI at the start of disease attack. A maximum of 2 applications can be made before grain watery ripe stage in winter and spring wheat, durum wheat, winter and spring barley, rye, triticale, and oats.

Rate of Application

Apply PIOLI at 2 litres of product per hectare in a water volume of 150 to 300 litres of water per hectare.

PIOLI may be applied at 2 litres per hectare in 100 litres water per hectare although efficacy at this reduced volume has not been evaluated. Therefore application at 100 litres water per hectare is at user's risk with regard to biological efficacy.

FOLLOWING CROPS

After treating a cereal crop with PIOLI: barley, cabbages, carrots, chicory, clover, dwarf beans, field beans, fodder beet, grass, leeks, lettuces, linseed, maize, oats, oilseed rape, onions, peas, potatoes, radishes, ryegrass, soya beans, spinach, sugar beet, sunflowers, triticale and wheat may be sown as the following crop.

MIXING AND SPRAYING

Mixing

Half fill the spray tank with clean water and start agitation. Shake the container well before use and pour in the required amount of product. Rinse empty containers thoroughly and add rinsings to the spray tank. Add the remainder of the water and continue agitation until spraying is complete.

When tank mixes are to be used, each product should be added separately to the spray tank, taking due note of any instructions given as to the order of mixing.

Application

Apply as a MEDIUM spray as defined by BCPC.

Compatibility

Consult ADAMA Agricultural Solutions UK Ltd or your distributor for details on the products which may be mixed with Pioli.

Always read the label of the partner product carefully before use.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Marketed by:

Adama Agricultural Solutions UK Ltd
Third Floor East
1410 Arlington Business Park
Theale
Reading
RG7 4SA
Telephone: +44 (0)1635 860555
Technical Helpline: +44 (0)1635 876622
www.adama.com
Email: ukenquiries@adama.com

Distributed in Ireland by:

Croplink (Ireland) Ltd
Springhill
Carrigtwohill
Co. Cork
T45 F685
Ireland
Tel: 021 4882806

Authorisation Holder: BASF plc, 4th and 5th Floors,
2 Stockport Exchange, Railway Road, Stockport, SK1 3GG

Pioli™ is a trademark of BASF plc.