



GOLTIX TITAN[®]



ADAMA

H

LJ0585IPFR_04

10Le



HERBICIDE

Suspension concentrate

MAPP17301

This product must be sold and used only in England, Scotland and Wales.



Ingredients

- Quinmerac 40 g/l (3.4% w/w)
- Metamitron 525 g/l (45.1% w/w)



Crops

A pre- and post-emergence herbicide for use against annual broad-leaved weeds in sugar beet and fodder beet.



Warning

Toxic to aquatic life with long-lasting effects.
Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.

Keep out of reach of children.
Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



Scan for Safety Data Sheet

Or use the weblink : <https://bit.ly/45tPVku>
Alternatively, contact your supplier.



R 3.2 Jul 26

This label is compliant with the Voluntary Initiative Guidance.

NO VARNISH AREA

Product name	GOLTIX TITAN [®]		
Country	UK		
Package size	10L		
Label code	LJ0585IPFR_04		
Label dimensions (w x h)	170mm x 150mm		
Label date	16.10.19 16.07.26	Name	
Smallest font on page	6.23 pt	Status	
		Template	Rect_SLT_ILang_UK-IE.indt



GOLTIX TITAN[®]



ADAMA

H



HERBICIDE

Suspension
concentrate

MAPP17301

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable, in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

TAKE OFF IMMEDIATELY all contaminated clothing.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

Environmental protection

Do not contaminate water with the product

or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

For advice on medical emergencies, fires or major spills telephone the National Chemical Emergency Centre on 01865 407333.

SHAKE WELL BEFORE USE

This product must be sold and used only in England, Scotland and Wales.

Ingredients

Quinmerac 40 g/l (3.4% w/w)
Metamitron 525 g/l (45.1% w/w)

Crops

A pre- and post-emergence herbicide for use against annual broad-leaved weeds in sugar beet and fodder beet.

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops: Sugar beet and fodder beet

Maximum individual dose; Maximum total dose; Latest time of application and Other specific restrictions: full details are given in the information box within the leaflet attached to the product. READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

IMPORTANT: Note that goods treated under the terms of this Great Britain (GB) authorisation can be legally marketed in Northern Ireland if they are being moved under the Northern Ireland Retail Movement Scheme. All other treated goods can only be marketed in Northern Ireland if they are in accordance with the statutory EU Maximum Residue Level (MRL) set under Regulation No 396/2005. This may also apply to residues in animal products where treated crops are fed to livestock. Growers are advised to draw this to the attention of distributors and retailers so that EU MRL breaches and any associated enforcement against goods marketed in Northern Ireland are avoided.

ADAMA Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park,
Theale, Reading, RG7 4SA
Tel: 01635 860555
Technical Helpline: 01635 876622
ukenquiries@adama.com www.adama.com

SHAKE WELL BEFORE USE

Scan for Safety Data Sheet

Or use the weblink : <https://bit.ly/45tPVku>
Alternatively, contact your supplier.



R 3.2 Jul 26

This label is
compliant with the
Voluntary Initiative
Guidance.



GROUP 5 | 4 HERBICIDE

Warning

Toxic to aquatic life with long-lasting effects.

Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.

Keep out of reach of children.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment,
comply with the instructions for use.



L105850CFR_04

2x10Le

Product name	GOLTIX TITAN [®]	
Country	UK	
Package size	10L	
Label code	L105850CFR_04	
Label dimensions (w x h)	300mm x 165mm	Name
Label date	16-10-22 16:07:26	Status
Smallest font on page	4.52 pt	Template Rect_SLT_ILang_NA_UK-IE.indt



GOLTIX TITAN®



HERBICIDE

Suspension concentrate

MAPP 17301



SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable, in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

TAKE OFF IMMEDIATELY all contaminated clothing.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from

farmyards and roads.

GROUP 5 | 4 HERBICIDE

Storage and disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

For advice on medical emergencies, fires or major spills telephone the National Chemical Emergency Centre on 01865 407333.

SHAKE WELL BEFORE USE

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops: Sugar beet and fodder beet

Maximum individual dose; Maximum total dose; Latest time of application and Other specific restrictions: full details are given in the information box within the leaflet attached to the product.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

IMPORTANT: Note that goods treated under the terms of this Great Britain (GB) authorisation can be legally marketed in Northern Ireland if they are being moved under the Northern Ireland Retail Movement Scheme. All other treated goods can only be marketed in Northern Ireland if they are in accordance with the statutory EU Maximum Residue Level (MRL) set under Regulation No 396/2005. This may also apply to residues in animal products where treated crops are fed to livestock. Growers are advised to draw this to the attention of distributors and retailers so that EU MRL breaches and any associated enforcement against goods marketed in Northern Ireland are avoided.

ADAMA Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park,
Theale, Reading, RG7 4SA
Tel: 01635 860555

Technical Helpline: 01635 876622
ukenquiries@adama.com www.adama.com

SHAKE WELL BEFORE USE



5 056325 504742

R 3.2 Jul 26

Product name	GOLTIX TITAN®		
Country	UK		
Package size	10L		
Label code	L10585IPBB_04		
Label dimensions (w x h)	170mm x 144mm		
Label date	16.10.2016 16.07.26	Name	Status
Smallest font on page	6.81pt	Template	Rect_5LT_ILang_NA_UK-IE.indt



GOLTIX TITAN®



ADAMA



HERBICIDE

**Suspension
concentrate**

MAPP 17301



SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable, in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

TAKE OFF IMMEDIATELY all contaminated clothing.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from

farmyards
and roads.

Storage and disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

For advice on medical emergencies, fires or major spills telephone the National Chemical Emergency Centre on 01865 407333.

SHAKE WELL BEFORE USE

GROUP 5 | 4 HERBICIDE

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops: Sugar beet and fodder beet

Maximum individual dose; Maximum total dose; Latest time of application and Other specific restrictions: full details are given in the information box within the leaflet attached to the product.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

IMPORTANT: Note that goods treated under the terms of this Great Britain (GB) authorisation can be legally marketed in Northern Ireland if they are being moved under the Northern Ireland Retail Movement Scheme. All other treated goods can only be marketed in Northern Ireland if they are in accordance with the statutory EU Maximum Residue Level (MRL) set under Regulation No 396/2005. This may also apply to residues in agricultural products where treated crops are fed to livestock. Growers are advised to draw this to the attention of distributors and retailers so that EU MRL breaches and any associated enforcement against goods marketed in Northern Ireland are avoided.

ADAMA Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park,
Theale, Reading, RG7 4SA
Tel: 01635 860555
Technical Helpline: 01635 876622
ukenquiries@adama.com www.adama.com

SHAKE WELL BEFORE USE

R 3.2 Jul 26



**Directions For
Use**



5 056325 504742

Open here



Product name	GOLTIX TITAN®		
Country	UK		
Package size	10L		
Label code	LJ0585IPBB_04		
Label dimensions (w x h)	45mm x 150mm		
Label date	16.10.19 16.07.26	Name	
Smallest font on page	6.8 pt	Status	
		Template	Rect_SLT_1Lang_NA_UK-IE.indt

Warning

Toxic to aquatic life with long-lasting effects.
Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.

Keep out of reach of children.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



This leaflet is part of the approved label.

GOLTIX TITAN® (MAPP 17301) is a suspension concentrate formulation containing 40 g/L (3.4% w/w) quinmerac and 525 g/L (45.1% w/w) metamitron. It is a pre- and post-emergence herbicide for use against annual broad-leaved weeds in sugar beet and fodder beet.

IMPORTANT INFORMATION FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE

Crop	Maximum individual dose (L product/ha)	Maximum total dose (L product/ha)	Latest time of application
Sugar beet, fodder beet (Pre- and post-em programme)	Pre-emergence 3.0 AND Post-emergence 1.0	6.0 (3.0 pre-emergence and 3.0 post-emergence) <i>See other specific restriction no. 4</i>	9 leaves unfolded (BBCH 19)
Sugar beet, fodder beet (Post-em only)	2.0	6.0 <i>See other specific restriction no. 4</i>	9 leaves unfolded (BBCH 19)

Other specific restrictions:

1. Do not handle treated crops for at least 2 days after treatment.
2. Fodder beet must not be grazed by livestock or harvested for animal consumption until at least 90 days following the last application of GOLTIX TITAN.
3. The minimum interval between applications is 5 days.
4. To be used either as pre-emergence followed by post-emergence use, OR post-emergence use only.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

Product name	GOLTIX TITAN®		
Country	UK		
Package size	10L		
Label code			
Label dimensions (w x h)	145mm x 150mm		
Label date	16.18.59 16.07.26	Name	
Smallest font on page	7 pt	Status	
		Template	Rect_SLT_1Lang_NA_UK-IE.indt

DIRECTIONS FOR USE

***IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.*

RESTRICTIONS

Do not apply to crops suffering from stress. This may be due to a number of factors e.g. nutrient or trace element deficiency such as manganese, soil acidity, drought, periods of low temperatures or frost, substantial day to night time fluctuations, high light intensity, wind or hail damage, use after other herbicides, pest or disease attacks.

Do not apply when temperatures exceed 21°C or in strong sunlight. If treatment is essential, delay spraying until the temperature falls and light intensity reduces.

An interval of at least 5 days must elapse between applications of GOLTIX TITAN.

RESISTANCE

GOLTIX TITAN contains metamiltrone, a Group C1 herbicide and quinmerac a Group O herbicide according to the mode of action classification system of the Herbicide Resistance Action Committee. Quinmerac has a narrow weed spectrum, with the main target for control being cleavers.

A strategy for preventing and managing resistance should be adopted. This should include integrating herbicides with a programme of cultural control measures. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the AHDB, your distributor, crop adviser or product manufacturer.

Growers are advised to apply products containing herbicides with different modes of action in sequence, as co-formulations or in tank mix, where two or more components are active against the target weeds.

Use the recommended rate of GOLTIX TITAN and the correct application timing for the hardest to control weed species present in the field. The above should be used in conjunction with effective cropping rotation and cultivation techniques e.g. stale seedbed, cultivation and use of non-selective herbicide prior to drilling.

WEEDS CONTROLLED

GOLTIX TITAN can be used pre or post-emergence. The best results are achieved when weeds are between the cotyledon and two-leaf stages.

	Pre-emergence followed by post-emergence programme	Post-emergence only programme
Annual mercury	-	R
Black-bindweed	-	R
Black nightshade	MR	S
Chickweed	MS	S
Cleavers	MS	S
Common fumitory	MR	R
Common field-speedwell	-	S
Common orache	-	S
Red dead-nettle	S	S
Fat-hen	S	MR
Field pansy	MS	MS
Field penny-cress	-	MS
Fool's parsley	-	R
Groundsel	S	S
Knotgrass	-	R
Pale Persicaria	-	-
Redshank	-	S
Shepherd's purse	S	S

S = Susceptible MS = Moderately susceptible

MR = Moderately resistant R = Resistant

CROP SPECIFIC INFORMATION

GOLTIX TITAN may be safely used on all commonly grown sugar beet and fodder beet varieties under normal conditions. There may be a reduction in crop tolerance if the beet is under stress; factors such as application in high light intensity, high temperatures, nutrient deficiency.

RATE OF USE

Pre-emergence followed by post-emergence

Apply 3.0 L/ha of GOLTIX TITAN in 80-200 L/ha of water after drilling of the crop but before crop emergence. Three further applications should be applied post-emergence of the crop from seed leaf stage: cotyledons completely unfolded to 9 leaves unfolded (BBCH 10-19). The minimum interval between applications is 5 days and the maximum total dose is 6.0 L/ha (maximum 3.0 L/ha pre-emergence and 3 x 1.0 L/ha post emergence).

Post-emergence only

Three applications, each of 2.0 L/ha of GOLTIX TITAN in 80-200 L/ha of water should be applied post-emergence to the crop, from seed leaf stage: cotyledons completely unfolded to 9 leaves unfolded (BBCH 10-19). The minimum interval between applications is 5 days and the maximum total application rate is 6.0 L/ha (maximum 3 x 2.0 L/ha).

FOLLOWING CROPS

Beet crops may be sown at any time following the use of GOLTIX TITAN.

Providing 16 weeks elapse from the last application of GOLTIX TITAN, winter cereals may be sown in the same season. After normal harvest, any crop can be grown in the following spring. Deep and thorough cultivation is necessary before sowing or planting succeeding crops. Mouldboard ploughing to a minimum depth of 15 cm is recommended before seedbed preparation.

MIXING AND SPRAYING

Shake GOLTIX TITAN well before use. Fill the spray tank with two thirds of the required amount of water. With the mixer switched on, add the product and the remaining water. Rinse empty containers thoroughly and add rinsings to the spray tank.

Continue agitation until spraying is completed.

Wash out sprayer thoroughly after use.

Apply as a MEDIUM spray as defined by BCPC.

Crops should not be re-entered until spray residues are dry.

COMPATIBILITY

For up-to-date details of compatible tank-mixes, contact your distributor or call the ADAMA technical helpline: 01635 876622 or use the on-line ADAMA tank-mix database at <https://www.adama.com/uk/en/tank-mix-database>

Always read the label of the partner product carefully before use.

COMPANY ADVISORY INFORMATION

This section is not part of the Product Label under the Plant Protection Products Regulations and provides additional advice on product use at the discretion of ADAMA Agricultural Solutions UK Ltd.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Marketed by:
ADAMA Agricultural Solutions UK Ltd
Third Floor East
1410 Arlington Business Park
Theale
Reading
RG7 4SA

Tel: 01635 860555
Technical Helpline: 01635 876622
www.adama.com
Email: ukenquiries@adama.com

GOLTIX TITAN® is a registered trademark of a company of the ADAMA Group.
© ADAMA Agricultural Solutions UK Ltd