






FirstAct™ Herbicide for the Double Team™ Sorghum Cropping Solution

FirstAct herbicide delivers sorghum growers reliable broad spectrum grass control along with flexible rotation and high yields for more profitable returns per acre.



- High yield potential and protection
- · Improved weed control and strong postemergence grass control
- · Crop safety for Double Team sorghum
- · Convenient application



Use Directions

- a. Do not apply FirstAct until sorghum has reached 8-leaf growth stage (approximately 11-12" sorghum) or significant crop injury may occur
- **b.** It is recommended that an over-the-top broadleaf herbicide application (if required) be made at least a week prior to FirstAct application
- **c.** Apply 5-12 fl oz/acre (most FirstAct rates will be 8-10 oz/acre)
- **d.** Difficult-to-control perennial weeds will require a use rate of 12 oz/acre
- e. Recommended minimum carrier volume of 15 gal/ac.
- f. Add crop oil concentrate at 1-2% v/v
- **q.** Two applications are allowed with maximum rate of 21 oz/acre/year
- h. The last application must be made at least 45 days prior to harvest

Target Crop

Double Team Sorghum

Target Weeds

Volunteer Corn, Sandbur, Texas Millet/Panicum, Crabgrass, Foxtail and Barnyardgrass*

*See label for additional grasses

Mode of Action

ACCase inhibitor (WSSA Group 1)

Signal Word

Caution

Restricted Use

No



FirstAct™ Herbicide for the Double Team™ Sorghum Cropping Solution

Ground Application Parameters

- · Use spray nozzles that will deliver medium or larger spray droplets
- Do not use flood, rain drop, whirl chamber, or other nozzle types that produce coarse, large spray droplets
- · Use a minimum of 10 gal of water per acre or a minimum of 15 gal of water per acre in arid areas
- · Increase spray volume and pressure as weed or crop density and size increase

Aerial Application Parameters

- · Use spray nozzles that will deliver coarse or larger spray droplets
- Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage
- · Use a minimum of 3 gal of water per acre or a minimum of 5 gal of water per acre in arid areas.

Mixing Instructions

- 1. Fill the tank ¼ to ½ full of water.
- 2. While agitating, add the required amount of FirstAct. If FirstAct and a tank mix partner are to be applied together, consult the tank mix partner label for information on which should be added first (normally granules and powders are added first).
- **3.** Continue agitation until the FirstAct is fully dispersed, at least 5 minutes.
- **4.** Once the FirstAct is fully dispersed, maintain agitation and continue filling tank with water.
- **5.** As the tank is filling, add the required volume of spray additives, always add these to the spray tank last.
- **6.** Apply FirstAct spray mixture within a reasonable period of time of mixing to avoid product degradation (24 to 48 hrs). If the spray mixture stands for any period of time, thoroughly re-agitate before using.

Crop Rotations

Do not rotate to crops other than the following within 120 days after application:

- Cotton
- · Dry Beans (including Chickpea)
- Flax
- · Lentils
- Mint (Spearmint and Peppermint)
- · Peas (Dry and Succulent Peas)
- Rapeseed Crop Subgroup 20A [includes Borage, Canola, Crambe, Gold of Pleasure (Camelina), Cuphea, Echium,

Hare's Ear Mustard, Lesquerella, Lunaria, Meadowfoam, Milkweed, Mustard Seed, Oil Radish, Poppy Seed, Sesame, and Sweet Rocket]

- · Snap Beans
- Soybeans
- Sunflowers
- Sugarbeets
- · Quizalofop-tolerant field corn

To learn more about Double Team Sorghum Cropping Solution with FirstAct herbicide, talk to your local ADAMA sale rep or visit **adama.com**

SEED SOLD THROUGH SORGHUM PARTNERS RETAILERS.





FirstAct[™]



