

## Double Team™ FAQ

**Question: Are Double Team, igrowth® and Inzen™ technologies the same?**

Answer: No, they are three different herbicide technologies with different weed control spectrums. FirstAct™, used with Double Team, contains a Group 1 (ACCase inhibitor), while Inzen's Zest™ contains nicosulfuron and igrowth's IMIFLEX™ contains imazamox, both Group 2 (ALS inhibitor).

**Question: Will FirstAct herbicide control volunteer corn?**

Answer: Yes, unless the volunteer corn is Enlist™ corn which is also resistant to "FOP" (Group 1) herbicides. This corn type is uncommon.

**Question: Do I have to use Parallel® or Parallel® Plus as my preemergence herbicide?**

Answer: Parallel and Parallel Plus are the preferred herbicides, but you can use any product that contains metolachlor, dimethenamid-P, or acetochlor and is registered for sorghum.

**Question: Can I tank mix broadleaf herbicide and FirstAct herbicide?**

Answer: We will not allow tank mixes in 2022. Appropriate tank mix partners and rates are being determined. Broadleaf products are recommended for application seven days before FirstAct applications.

**Question: When can I make first application of FirstAct herbicide?**

Answer: Plant size is important to reduce crop response. There are two ways to determine if the crop is big enough to spray: plant height or leaf number. The plant height needs to be at least 11 inches tall, or the plant needs to be at the 8-leaf stage. See the FirstAct Tech sheet for additional information.

**Question: How many times per season can I spray FirstAct herbicide?**

Answer: Two applications are allowed with a maximum of 21oz of FirstAct per acre.

**Question: How late can I spray FirstAct herbicide?**

Answer: While the label allows for applications up to 45 days before harvest, generally the most effective weed control of nearly all weed species will occur early in the growing season.

**Question: Are adjuvants required with FirstAct herbicide applications?**

Answer: Yes, crop oil concentrate (COC) is recommended at 1% to 2% v/v. The higher rate is recommended when under drought stress.



**Question: What carrier spray volume is needed for FirstAct applications?**

Answer: Ground applications are labeled at a minimum of 10 gallons per acre but for better coverage, 15 gallons per acre is highly recommended, especially under drought stress.

**Question: Are Double Team hybrids SCA (sugarcane aphid)-tolerant?**

Answer: No, the first-generation hybrids will not be SCA-tolerant. In 2022, all Double Team seed will be treated with the highest label rate of Cruiser seed treatment, which should provide 60+ days of SCA control.

**Question: Can glyphosate herbicides be sprayed over the top of Double Team sorghum?**

Answer: No, glyphosate and products like Select (or any DIM chemistry) will kill Double Team sorghum. Glyphosate and other registered products can be used in burndowns prior to planting.

**Question: Are there grain channel requirements for 2022?**

Answer: No specific requirements are in place for 2022.

**Question: Is there a yield drag with this trait?**

Answer: No, yields have been equal to or higher than similar maturity hybrids.

**For more information about Double Team, visit [www.DoubleTeamSorghum.com](http://www.DoubleTeamSorghum.com)**

