



FUNGICIDE

MAXENTIS™ SC

The ultimate broad-spectrum fungicide with breakout performance against tar spot, northern corn leaf blight, grey leaf spot, frogeye leaf spot, and brown spot for growers looking to increase plant health overall.

What is Maxentis SC?

Maxentis SC is an optimized mixture of prothioconazole and azoxystrobin for use in row crops and more. It offers longer-lasting, broad-spectrum control of diseases so that growers can enhance overall plant health, leading to better yields and profit.

Benefits


- High-quality Formulation and Ready-to-use Premix Solution
- Long-lasting Disease Control
- High Efficacy on Target Diseases
- Flexible Application Window

Resistance Management Guidelines

Maxentis SC is a mixture of prothioconazole (FRAC Group 3: DMI) and azoxystrobin (FRAC Group 11: QoI) which controls target plant pathogenic fungi by disrupting sterol production, membrane synthesis, and fungal respiration. Any fungal population may contain individuals naturally resistant to Maxentis SC and other Group 3 and Group 11 fungicides.

Take the following steps to delay fungicide resistance:

- Rotate the use of Maxentis SC or other Group 3 and Group 11 fungicides with different groups that control the same pathogens
- Use at least the minimum application rate as labeled by the manufacturer
- Monitor efficacy of all fungicides used in the IPM program against the targeted pathogen and record factors that may influence fungicide performance and/or disease development
- Adopt an integrated disease management program that utilizes scouting and includes historical information related to pesticide use, and crop rotation



Active Ingredient

Prothioconazole +
Azoxystrobin

Formulation

Suspension Concentrate

Target Crops

Corn, Soybeans, Wheat

Target Pests

Tar Spot, Northern Corn Leaf Blight, Rusts, Frogeye Leaf Spot, Brown Spot, Powdery Mildew, Septoria Blotch, Tan Spot and more





Maxentis™ SC

Crop, Disease and Rate Information

The chart below provides an overview of the crops, diseases and rates that are recommended for Maxentis SC. Please refer to the label for full directions for use by specific crop groups.

Crop	Disease(s) Treated	Rate/Acre
Corn	Anthrachnose Leaf Blight, Eyespot, Gray Leaf Spot, Northern Corn Leaf Blight, Northern Corn Leaf Spot, Rust, Southern Corn Leaf Blight, Tar Spot	• 5 - 13 fl oz / A
Soybean*	Alternaria Leaf Spot, Anthracnose, Asian Soybean Rust, Brown Spot, Cercospora Blight, Frogeye Leaf Spot, Powdery Mildew, Rhizoctonia Aerial Blight	• 6 - 11 fl oz / A
Wheat	Powdery Mildew, Rusts, Septoria Leaf Blotch, Tan Spot	• 7 - 14 fl oz / A

* Not for use in California

Additional crops on the label but not listed above include:

- Barley
- Oats / Rye
- Bushberry
- Chickpea
- Cotton
- Cucurbit Vegetables
- Dried Shelled Peas and Beans (except soybeans)
- Lentils
- Cranberry
- Peanuts
- Rapeseed
- Rice
- Sugar Beet

Always read and follow label directions. Data on file. Maxentis is a trademark of an ADAMA Group Company. ©2024 ADAMA

Visit adama.com to learn more.