

Dicamba 4L DMA

A herbicide that controls a wide spectrum of broadleaf weeds.



What is Dicamba 4L DMA herbicide?

Dicamba 4L DMA is a selective pre- and postemergence herbicide for control of important broadleaf weeds infesting a range of crops including corn, sorghum, small grains and cotton.

Benefits

- Trusted performance growers expect
- Powerful mode of action
- Cost-effective control option

Corn Application Guidelines

See the full product label for all application/usage instructions and restrictions.

Preemergence or Preplant	Preemergence	Early Postemergence	Late Postemergence
<i>No-tillage</i>	<i>Conventional or Reduced Tillage</i>	<i>All Tillage Systems</i>	<i>All Tillage Systems</i>
n/a	n/a	Spike through 8" tall corn	8-36" tall corn
1 pint/A on medium and fine textured soils with 2% or greater organic matter	1 pint/A on medium and fine textured soils with 2% or greater organic matter	1 pint/A may be applied from corn emergence through the five-leaf stage or 8" tall, whichever comes first	1/2 pint/A may be made from 8" to 36" tall corn or 15 days before tassel emergence, whichever comes first
1/2 pint/A on coarse textured soils or medium and fine textured soils with less than 2% organic matter	Do not apply to course textured soil until after crop emergence	Reduce the rate to 1/2 pint/A if corn is growing on coarse textured soils	For best performance, make applications when weeds are less than 3" tall

Active Ingredient

Dimethylamine salt of dicamba

Formulation Type

Liquid

Target Crop

Crops including Cereals (corn, sorghum, small grains, etc.) and Cotton

Target Weeds

Annual, Biennial and Perennial broadleaf weeds

See label for full listing of weeds





Dicamba 4L DMA

Cotton Application Guidelines*

See the full product label for all application/usage instructions and restrictions.

Preplant	Tank Mixes	Restrictions
<i>Conventional or Conservation Tillage</i>	May be tank mixed with prometryn, paraquat, and glyphosate herbicides.	<i>Do not make preplant applications to geographic areas with average annual rainfall less than 25"</i>
Prior to planting cotton		
8 fl oz/A (0.25 lb ai/A)	It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use.	<i>Do not apply preplant to cotton west of the Rockies</i>
For best performance, apply when weeds are in the 2- to 4-leaf stage and rosettes are less than 2" across		

Sorghum Application Guidelines

See the full product label for all application/usage instructions and restrictions.

Preplant	Postemergence	Preharvest Uses
n/a	Spike stage but before 15" tall	Only in states of Texas and Oklahoma
1/2 pint/A broadcast rate	1/2 pint/A broadcast rate	1/2 pint/A broadcast rate
May be applied to emerged and actively growing weeds at least 15 days prior to planting	For best performance, make applications when sorghum is in the 3- to 5-leaf stage and weeds are small	May be applied for weed suppression any time after the sorghum has reached the soft dough stage. An agriculturally approved surfactant may be used to improve performance.

* Not registered in California

To learn more about, visit [ADAMA.COM](https://www.adama.com)