Precautions: Toxic to aquatic organisms and bees; do not contaminate water bodies and avoid usage when bees are foraging. Wear protective clothing when using this product. Do not you eat, drink or smoke when handling it. After work change clothing, wash exposed skin with soap before eating, drinking or smoking.

Handling/storage/disposal: Keep in original tightly-closed container in a dry, cool, safe place away from food and drinks; and beyond reach of children. Dispose of container by triple rinsing, puncturing and burying away from water sources.

First aid instructions: If Swallowed: Have victim sip a alass of water. Do not induce vomiting unless told to by a doctor. Never give anything by mouth to an unconscious person. If on skin: Take off contaminated clothing and wash affected skin area immediately with plenty of water and soap. If in eyes: Rinse eye with clean water for 10-15 minutes. If Inhaled: Move victim into fresh air. If victim is not breathing, call a doctor immediately and give artificial respiration, preferably mouth-tomouth. In all cases: Seek medical attention immediately; have the product container or label with you. Note to the doctor: There is no specific antidote; treat symptomatically and give supportive therapy.

WHO toxicity class: II (Moderately toxic)

INESECTICIDE



Imidachloprid 250 q/l + Bifenthrin 50 q/

Systemic and contact properties for longer and more effective control of insects.

Crops: Legumes, vegetables, cotton, ornamentals, fruit trees











500ml.

Batch No.	Date of Manufacture:	Expiry date:	
EPA Registration N°:		Scan for	

Adama Makhteshim Ltd. P. O. Box 60, Industrial Zone, Beer Sheva 84100, Israel. Tel/Fax: 972 8 6280304 / 6296602

Adama West Africa Ltd. No 7 Adembra Road, East Cantonments. P. O. Box KT 194, Accra Tel/Fax: + 233 (0) 302775030



Description:

A broad-spectrum insecticide combining systemic and contact properties for effective control of a wide range of insect pests on many crops.

DIRECTIONS FOR USE- Mixing:

Half-fill the spray tank with clean water. Add the required amount of Galil 300 SC to the water and gently agitate the machine. Fill the spray tank to the squired level while agitating continuously.

Dosage:

The recommended dosage is 15-20 ml per 15 L of water.

Usage:

Apply when signs of pest outbreaks are seen.

Pre-harvest interval:

7 days.

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN























