PRECAUTIONS: Toxic to aquatic organisms and bees. Wear protective clothing. Do not eat, drink or smoke when using it. Aft work, change clothing, wash exposed skin with soap before eating, drinking or smoking.

STORAGE/DISPOSAL: Keep in tightly-closed original container in dry and cool place, away from children. Dispose of cotainer by triple rinsing, puncturing and burying away from water sources. FIRST AID: If swallowed: Have victim sip water; do not induce vomiting. If in eyes: Flush eyes with clean water for 15-20 minutes If on skin: Remove clothing immediately and wash affected area with soap. If inhaled: Move victim into fresh air; give artificial respiration if necessary. In all cases: Consult medical facility if necessary. To physician: No known antidote; treat symptomatically and give supportive therapy.









INSECTICIDE

PYRINEX QUICK[™] 424 FC

Chlorpyrifos 400 + deltamethrin 24 q/l

Broad-spectrum, with residual and fast-acting contact properties to control insect pests in field crops, ornamentals and in-doors; also for wood protection in buildings. Crops: Yam, Cassava, Legumes, Water melon,

Lettuce, Okra, Pepper, Tomato, Pineapple & Cotton,



Manufacturer Adama Makhteshim Ltd P. O. Box 60. Industrial Zone Beer Sheva 84100. Israel Tel: 972-8-6280304

Distributor: Adama West Africa Ltd No 7 Adembra Road. East Cantonments, Accra. Tel: + 233 (0) 302775030

ADAMA

DIRECTIONS FOR LISE, Son Table for details

| Crops/sites | Major pests | Dosages | Methods of application |
|--|--|--------------------------|---|
| Lawns, legumes, ornamentals, vegetables, pineapple | Crickets, defoliators, mealybugs, aphids, ants, termites, other soil-borne pests, leaf miners | 40-70 ml/15L of water | Lawns: Use 400 ml of solution per m2. Water lawn thereafter. Ornamentals: Spray to drip. Pineapple: Dip sucker in solution before planting. Drench base thereafter. |
| Soi l treatment | Crickets, termites, ants, other soil-borne pests | 80 ml/15L of water | Drench soil for 5-7 days prior to planting. Repeat when necessary. |
| Wood preservation | Termites, wood borers | 60-80 ml/15L of water | Brush or spray to drip every surface of sawn timber and leave to dry. |
| Protection of buildings in termite endemic areas | Termites | 60-80 ml/15L of water | Dig shallowly along walls. Drench furrow with solution and cover. Repeat as necessary. |

Mixing: Fill spray tank half way with water. Pre-mix required Pyrinex-quick. Pour mixture into spray tank and agitate sprayer. Fill tank with water to the required level. Usage: Apply Pyrinex-quick when pests are noticed. Agitate sprayer before spraying and every 5 minutes during course of spraying. Pre-harvest interval: 14 days

P. O. Box KT 194, Accra, Ghana WHO Toxicity Class: II (Moderately toxic)

Batch No. Date of Manuf: Expiry date: EPA Registration N°:





















