

Precautions: Toxic to mammals, birds and bees. Wear protective clothing. Never eat, drink or smoke when using the product. After work, change clothing, wash hands and exposed skin with soap.

HANDLING/STORAGE/DISPOSAL: Handle with care; keep in original tightly-closed container in a dry, cool, safe place away from food, drinks and animal feed-stuffs; and out of reach of children. Never re-use container for any other purpose.

FIRST AID INSTRUCTIONS: Symptoms of poisoning include headache and salivation. If **swallowed:** Do not induce vomiting, wash out mouth with plenty of water. If **in eyes:** Flush eyes immediately with plenty of water for 10-15 minutes. If **on skin:** Remove clothing and wash affected area. If **inhaled:** Move victim into fresh air; give artificial respiration if necessary. In **all cases:** Consult a medical facility if victim's conditions so demand.

Note for physician: No specific antidote known; treat symptomatically and give supportive therapy.

WHO toxicity class: II (Moderately toxic)

ADAMA

Rimon™

10 EC

Novaluron 100 g/L

Controls larvae of pest insects belonging to moths/butterflies, beetles, bugs and flies. Good for IPM.

Crops: Cabbage, Tomato, Legumes, Mango, Citrus & Maize.

150ml.

Manufacturer:

Adama Makhteshim Ltd,
P. O. Box 60, Industrial Zone,
Beer Sheva 84100, Israel.
Tel/Fax: 972 8 6280304 /
6296602

Distributor:

Adama West Africa Ltd,
No 7 Adembra Road, East
Cantonments,
P. O. Box KT 194, Accra



INSECTICIDE

Description: Rimon 10 EC is an Insect Growth Regulator (IGR), which inhibits chitin formation and causes abortive moulting, thereby disrupting development in insects.

Mixing: Fill spray tank with half the required amount of clean water. Add the recommended amount of Rimon 10 EC and mix thoroughly. Top up the spray tank with water to the required level and agitate well to ensure thorough mixture.

Usage: Start application when first signs of pest infestation appear. Repeat application every 7-10 days till threats of pests are over.

Pre-harvest interval: 7 days



Batch No.	Date of Manufacture:	Expiry date:	EPA Registration N°:

DIRECTIONS FOR USE in Crops, Pests and Application Rates:

Crops	Pests	Rates	Remarks
Maize	Stem borers & defoliators	40-60 ml in 15 L water	Ensure good coverage of foliage/other plant surfaces.
Citrus & mango	Leaf miners, defoliators & fruit flies	40-60 ml in 15 L water	Ensure good coverage of foliage/other plant surfaces.
Vegetables	Diamond back moth, caterpillars, thrips, leaf miners, aphids & white flies	40-60 ml in 15 L water	Ensure good coverage of foliage/other plant surfaces.



HARMFUL

