

## READ THE LABEL BEFORE USE

**Safety precautions: KEEP OUT OF REACH OF CHILDREN.**

**WEAR** suitable protective clothing, including hand gloves, face mask, eye and nose masks when handling the concentrate and during application. Wash hands thoroughly after handling.

**DO NOT** eat, drink or smoke when using this product. **WASH** all protective clothing thoroughly after use, especially the inner part of gloves.

**Storage and disposal:** Keep in original container, tightly closed in a safe place. Do not re-use container for any purpose.

**Warning:** Very toxic to aquatic life with long lasting effects.

**FIRST AID INSTRUCTIONS: If swallowed:** give activated charcoal in a glass of water. Never give anything by mouth to an unconscious person. Do not induce vomiting. **If in eyes:** flush eyes immediately with plenty of clean water for at least 15 minutes. **If on skin:** remove clothing immediately and wash affected area well with soap and clean water. **If inhaled:** move victim into fresh air; keep him/her warm and at rest, give artificial respiration if necessary; preferably mouth-to-mouth.

**In all cases:** consult a medical facility if victim's conditions so Demand. **Antidote: Note to physician:** no specific antidote known; treat symptomatically and give supportive therapy.

# FUNGICIDE

## BANJO FORTE™

### 400 SC

200g/l dimethomorph +  
200g/l fluazinam

Suspension concentrate



ADAMA

Effective non-copper fungicide for  
the Control of  
Black pod Disease on Cocoa

Batch No.	Date of Manuf:	Expiry date:

250ml.

**Manufacturer:**  
ADAMA Makhteshim Ltd,  
Industrial Zone, P.O. Box  
60, Beer-Sheva 84100, Israel

**Distributor:**  
Adama West Africa Ltd,  
# 4 Fourth Norla Street, North  
Labone, P. O. Box KT 194, Accra,  
Ghana. Tel: 0302 775030

## PRODUCT DESCRIPTION

Banjo Forte 400 SC is a non-copperfoliar fungicide with protectant And translaminar activity. It is effective against black pod disease caused by *phytophthora* on cocoa and on other crops.

**DIRECTIONS FOR USE: Mixing and application** Half-fill the spray tank with clean water. Add the recommended amount of Banjo forte 400 SC and the remainder of the water while agitating the spraying machine continuously. Spray to drip on pods, including cherrels.

**Dosage:** 75 ml of Banjo forte 400 SC in 15 L of water (i.e., 1.125 L of Banjo Forte in 225 L of water/ha).

**Timing and frequency of application:** Apply at 4 weekly intervals from April/May (depending on the start of the rains) to September/ November Annually.

**Pre-harvest interval:** 7 days.

**Re-entry period:** At least 24 hrs after treatment.



UNTREATED



TREATED

EPA Registration N°:



## CAUTION

