PRECAUTION/HAZARD MEASURES:

Dangerous to domestic animals, birds, bees and fish; do not contaminate water bodies. Avoid breathing in vapour. Wear protective clothing, including gloves, when using the product. Do not eat, drink or smoke when using the product. After work, change clothing, wash hands and exposed skin with soap before eating, drinking or smoking. HANDLING/STORAGE/DISPOSAL:

Handle with care; keep in original tightly-closed container in a dry, cool, safe place away from food, drinks and animal feed-stuffs; and out of reach of children. Dispose of container by emptying completely, puncturing and burying away from water sources. Never re-use container for any other purpose.

FIRST AID INSTRUCTIONS:

Symptoms of poisoning may include excessive sweating, headache, weakness, faintness, nausea, stomach pains and blurred vision. If swallowed: Give artificial charcoal; induce vomiting. Never give anything by mouth to an unconscious person. If in eyes: Flush eyes immediately with plenty of clean water for at least 15 minutes.

If on skin: Remove clothing immediately and wash affected area well with soap and clean water. If inhaled: Move victim into fresh air; keep him/her warm and at rest, give artificial respiration if necessary; preferably mouth-to-mouth.

In all cases: consult a medical facility if victim's conditions so demand.

NOTE FOR PHYSICIAN: No specific antidote known; treat symptomatically

WHO toxicity class: II (Moderately toxic)

INSECTICIDE

Deltapaz

1.25 EC

Deltamethrin 12.5 g/L

Suitable for controlling a wide range of insect pests in field crops and ornamentals.

Crops: Cabbage, Okra, Legumes, Mango, Citrus & Tomato











Batch No.	Date of Manufacture:	Expiry date:

EPA Registration N°:

Manufacturer: Adama Makhteshim Ltd. P. O. Box 60. Industrial Zone. Beer Sheva 84100, Israel. Tel/Fax: 972 8 6280304 /

Distributor: Adama West Africa Ltd. No 7 Adembra Road, East Cantonments. P. O. Box KT 194, Accra

Brocheur

Scan for

DESCRIPTION:

Deltapaz 1.25 EC is a broad-spectrum pyrethroid insecticide, with contact and ingestion activity; has quick knock-down effect with short persistence. It is compatible with most commonly used pesticides.

DIRECTIONS FOR USE per Crops, Pests and Application Rates:

Crops	Pests	Rates	Remarks
Legumes	Defoliators, Pod borers, aphids & white flies	40-60 ml in 15 L water	Ensure good coverage of foliage/other plant surfaces
Citrus & mango	Mealy bugs, aphids, ants & defoliators	40-60 ml in 15 L water	Ensure good coverage of foliage/other plant surfaces
Cabbage	Diamond back moth, aphids & thrips,	40-60 ml in 15 L water	Ensure good coverage of foliage/other plant surfaces
Okra	Leaf miners, defoliators, white flies & aphids	40-60 ml in 15 L water	Ensure good coverage of foliage/other plant surfaces
Tomato	Aphids, thrips & white flies	40-60 ml in 15 L water	Ensure good coverage of foliage/other plant surfaces

Mixing: Fill the spray tank with half the required amount of water. Add the recommended amount of Deltapaz EC and mix thoroughly. Top up the spray tank with water to the required level and agitate well to ensure thorough mixture.

Usage: Start application when first signs of pest infestation appear. Repeat application every 7-10 days till threats of pests are over. To avoid resistance development in pests, apply Deltapaz 1.25 EC in a spray program that incorporates other products with different modes of action.

PRE-HARVEST INTERVAL: 1-2 days

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN























