

Precaution/hazard measures:

Harmful to fish; do not contaminate water bodies.

Avoid breathing in vapour. Wear protective clothing, including gloves, when using the product.

Do not eat, drink or smoke when using the product.

After work, change clothing, wash hands and exposed skin with soap before eating, drinking or smoking.

Handling/storage/disposal:

Keep in original tightly-closed container in a dry, cool, safe place away from food, drinks and animal feed-stuffs; and out of reach of children. Dispose of container by emptying completely, puncturing and burying away from water sources. Never re-use container for any other purpose

First aid instructions:

If swallowed – give a large amount of water to drink; induce vomiting.

If in eyes – flush eyes immediately with plenty of clean water for at least 15 minutes.

If on skin – remove clothing immediately and wash affected area well with soap and clean water.

If inhaled – move victim into fresh air; give artificial respiration if necessary; preferably mouth-to-mouth.

In all cases – consult a medical facility if victim's conditions so demand.

Note for physician:

No specific antidote known; treat symptomatically.

WHO toxicity class: III (Slightly Toxic)

FUNGICIDE

Folpan™

50WP

Folpet 500g/Kg

Broad-spectrum, protective, organic fungicide used on food crops and ornamentals against a wide range of fungal pathogens.

Crops:

Cauliflower, Tomato, Onion, Pepper, Citrus



ADAMA



1Kg

EPA Registration N°:		
Batch No.	Date of Manufacture:	Expiry date:

Manufacturer:
Adama Makhteshim Ltd,
P. O. Box 60, Industrial Zone,
Beer Sheva 84100, Israel.
Tel/Fax: 972 8 6296713

Distributor:
Adama West Africa Ltd,
No 7 Adembra Road, East
Cantonments,
P. O. Box KT 194, Accra

Scan for
Brochure



Description:

Folpan 50 WP is a contact fungicide containing Folpet in wettable powder formulation. It is compatible with most commonly used pesticides except those of alkaline nature and oils; not to be applied closely after oil application.

Direction for General agricultural use:

Apply **Folpan 50 WP** at
2-3 kg/ha in 200-400 L of water
OR
250-300g/15 L of water (knapsack sprayer).

Mixing:

Fill the spray tank with half the required amount of water. Add the recommended amount of Folpan 50 WP and mix thoroughly. Top up the spray tank with water to the required level and agitate well to ensure thorough mixture.

Usage:

Start application when first signs of disease infection appear OR when weather conditions favour disease development. Repeat regularly every 7 days until threat of disease is over; the interval may be reduced during periods of intense shoot growth or when serious infections are likely. Thorough coverage of the crop, including upper and lower leaves, is recommended.

Pre-harvest interval: 10 days

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN



CAUTION

