

Precaution/hazard measures:

Harmful to fish; do not contaminate water bodies.

Avoid breathing in vapour. Wear protective clothing, including gloves, when using the product.

Do not eat, drink or smoke when using the product.

After work, change clothing, wash hands and exposed skin with soap before eating, drinking or smoking.

Handling/storage/disposal:

Keep in original tightly-closed container in a dry, cool, safe place away from food, drinks and animal feed-stuffs; and out of reach of children. Dispose of container by emptying completely, puncturing and burying away from water sources. Never re-use container for any other purpose

First aid instructions:

If swallowed – give a large amount of water to drink; induce vomiting.

If in eyes – flush eyes immediately with plenty of clean water for at least 15 minutes.

If on skin – remove clothing immediately and wash affected area well with soap and clean water.

If inhaled – move victim into fresh air; give artificial respiration if necessary; preferably mouth-to-mouth.

In all cases – consult a medical facility if victim's conditions so demand.

Note for physician:

No specific antidote known; treat symptomatically.

WHO toxicity class: IV (Practically non-toxic)

FUNGICIDE

Merpan™

50WP

Captan 500g/Kg

Broad-spectrum, fungicide for use against heart rot, damping off, moulds, blight, Cercospora and Anthracnose.

Crops:

Fruits, Seeds, Coffee, Vegetables & Ornamentals.



1Kg

EPA Registration N°:		
Batch No.	Date of Manufacture:	Expiry date:

Manufacturer:
Adama Makhteshim Ltd,
P. O. Box 60, Industrial Zone,
Beer Sheva 84100, Israel.
Tel/Fax: 972 8 6296713

Distributor:
Adama West Africa Ltd,
No 7 Adembra Road, East
Cantonments,
P. O. Box KT 194, Accra

Scan for
Brochure



Description:

An organic, wettable powder, protective fungicide; very effective for the control of various fungal diseases on many crops.

DIRECTIONS FOR USE:

Foliar Application

Quantity of product	Volume of water
2-3 kg/ha	200 L
150-220g	15 L

Dipping treatment of pineapple suckers, seeds, grains and cuttings

Dip seed, seedling or sucker into a solution of 50g/10L of water for at least 2 minutes before planting or storage.

Special Uses

Pre-planting soil treatment:

Broadcast onto the ground and work into top 10 cm of the soil.

Post-harvest dip of fruits:

Dip fruits for 2 minutes into 0.6% solution before storage.

Mixing:

Mix required amount of Merpan 50 WP with little amount of water. Add this mixture to the bulk of required volume of water in the spray tank and agitate well.

Usage:

Merpan 50 WP is a protective fungicide, so the first application should be made just before the disease is expected to appear. Repeat application at 6-7 day intervals for as long as protection is required.

Pre-harvest interval: 7 days

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN



CAUTION

