

2026-2285
2026-06-26

Container Label

GROUP	14	HERBICIDE
--------------	-----------	------------------

SAFACYN®

Herbicide Solution

COMMERCIAL (AGRICULTURAL)

For pre-seed, pre-emergent and chemfallow application for control of broadleaf weeds, for pre-harvest weed management in wheat, barley, and triticale, and for harvest aid in canola, dry common beans, chickpeas, red lentil varieties, dry field peas, faba beans, flax, mustard, soybeans and sunflowers IN MANITOBA, SASKATCHEWAN, ALBERTA AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION).

For sale for use on raspberry and blackberry IN EASTERN CANADA AND BRITISH COLUMBIA (EXCLUDING THE PEACE RIVER REGION), AND ON ALL OTHER LABELLED CROPS IN EASTERN CANADA.

ACTIVE INGREDIENT: Saflufenacil 160 g/L

DANGER – CORROSIVE TO EYES
POTENTIAL SKIN SENSITIZER

REGISTRATION NO. 36039 PEST CONTROL PRODUCTS ACT

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

NET CONTENTS: 0.5 L – 1050 L, Bulk

ADAMA Agricultural Solutions Canada Ltd.
300 – 191 Lombard Avenue
Winnipeg, MB R3B 0X1
1-855-264-6262

For emergency medical help and health/safety inquiries call ProPharma Group at 1.877.250.9291 (24 hours a day).
For spill, leak or fire call INFOTRAC at 1.800.535.5053 (24 hours a day).

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT take internally. Harmful if swallowed.

Avoid contact with skin, eyes or clothing. Severely irritating to the eye. **DO NOT** get in eyes. Potential skin sensitizer.

Avoid inhalation of vapor or spray. Use with adequate ventilation.

DO NOT eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, using the toilet or handling mobile devices. Aircraft cockpit and vehicle cabs **MUST** be decontaminated regularly.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Personal Protective Equipment (PPE)

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair.

For application using ground equipment: Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Gloves are not required during application within a closed cab.

If the product comes in contact with clothing/PPE, remove all contaminated clothing/PPE, wash the outside of gloves with soap and water before removing, wash contaminated skin with soap and water, and then dress in clean clothing. Launder contaminated clothing/PPE regularly and separately from other laundry. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. **DO NOT** reuse them.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then

give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic plants and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce run-off from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

STORAGE

Store this product away from food or feed.

DO NOT ship or store the product near food, feed, seed or fertilizers.

Store in original container with the lid tightly closed, in a cool, secure, well-ventilated area.

DISPOSAL

Recyclable Containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers (containers that can be refilled for the user by the distributor/dealer)

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any

2026-2285
2026-06-26

other purpose.

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Booklet

GROUP	14	HERBICIDE
--------------	-----------	------------------

SAFACYN®

Herbicide Solution

COMMERCIAL (AGRICULTURAL)

For pre-seed, pre-emergent and chemfallow application for control of broadleaf weeds, for pre-harvest weed management in wheat, barley, and triticale, and for harvest aid in canola, dry common beans, chickpeas, red lentil varieties, dry field peas, faba beans, flax, mustard, soybeans and sunflowers IN MANITOBA, SASKATCHEWAN, ALBERTA AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION).

For sale for use on raspberry and blackberry IN EASTERN CANADA AND BRITISH COLUMBIA (EXCLUDING THE PEACE RIVER REGION), AND ON ALL OTHER LABELLED CROPS IN EASTERN CANADA.

ACTIVE INGREDIENT: Saflufenacil 160 g/L

DANGER – CORROSIVE TO EYES
POTENTIAL SKIN SENSITIZER

REGISTRATION NO. 36039 PEST CONTROL PRODUCTS ACT

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

NET CONTENTS: 0.5 L – 1050 L, Bulk

ADAMA Agricultural Solutions Canada Ltd.
300 – 191 Lombard Avenue
Winnipeg, MB R3B 0X1
1-855-264-6262

For emergency medical help and health/safety inquiries call ProPharma Group at 1.877.250.9291 (24 hours a day)
For spill, leak or fire call INFOTRAC at 1.800.535.5053 (24 hours a day)

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT take internally. Harmful if swallowed.

Avoid contact with skin, eyes or clothing. Severely irritating to the eye. **DO NOT** get in eyes. Potential skin sensitizer.

Avoid inhalation of vapor or spray. Use with adequate ventilation.

DO NOT eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, using the toilet or handling mobile devices. Aircraft cockpit and vehicle cabs **MUST** be decontaminated regularly.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Personal Protective Equipment (PPE)

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair.

For application using ground equipment: Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Gloves are not required during application within a closed cab.

If the product comes in contact with clothing/PPE, remove all contaminated clothing/PPE, wash the outside of gloves with soap and water before removing, wash contaminated skin with soap and water, and then dress in clean clothing. Launder contaminated clothing/PPE regularly and separately from other laundry. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. **DO NOT** reuse them.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then

give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic plants and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

STORAGE

Store this product away from food or feed.

DO NOT ship or store the product near food, feed, seed or fertilizers.

Store in original container with the lid tightly closed, in a cool, secure, well-ventilated area.

DISPOSAL

Recyclable Containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers (containers that can be refilled for the user by the distributor/dealer)

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any

other purpose.

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

DIRECTIONS FOR USE

GENERAL INFORMATION

SAFACYN® is a herbicide for broadleaf weed control.

SAFACYN® is rapidly absorbed by root and foliar uptake; once absorbed it exhibits mobility in plants. **SAFACYN®** is a potent inhibitor of protoporphyrinogen oxidase (PPO). Cell membrane damage induced by inhibition of PPO leads to plant death. Susceptible weeds develop injury symptoms within hours of application under active growing conditions; plant death occurs within 3 to 5 days depending upon growing conditions.

SAFACYN® is recommended for pre-seed, pre-plant and pre-emergent applications. **SAFACYN®** may also be applied in fallow crop lands and post-harvest and as a pre-harvest weed management treatment and as a harvest aid in registered crops. For control of emerged broadleaf weeds, **SAFACYN®** may be applied as a banded application directed to the base of raspberry and blackberry prior to the plants breaking winter dormancy.

SAFACYN® does not control grass weeds. **SAFACYN®** should always be tank mixed with glyphosate (and/or an appropriate herbicide with graminicide activity as needed) for broad spectrum weed control. Refer to the **SAFACYN® + GLYPHOSATE TANK MIX CROP USE RATES** section for additional detailed instructions.

SAFACYN® is a broad spectrum weed resistance management tool for activity on a range of broadleaf weeds.

For good coverage and optimal control use a minimum of 100 L/ha (field sprayer application) and 50 L/ha (aerial application) of water volume.

MIXING INSTRUCTIONS

1. When using **SAFACYN®**, always start with a clean sprayer. Thoroughly clean the sprayer by flushing the system with water containing detergent. Refer to previously applied product labels for specific cleaning instructions.
2. Completely fill spray tank with water, leaving sufficient room for the addition of products. Start agitation system.
3. Add **SAFACYN®** first and continue to agitate until thoroughly mixed.
4. When tank mixing, add tank-mix partner and continue agitation.
5. Add the correct amount of adjuvant (a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS)

- (e.g. Adama Adjuvant 80) at 0.25% v/v) to the tank last.
6. Continue agitation while filling the remainder of the tank with water necessary to fill the spray tank.
 7. Continue to agitate or run the by-pass system.
 8. After any break in spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to remix the spray materials. Use tank mixtures immediately, or as soon as possible, after mixing with constant and adequate agitation. Do not allow the mixture to sit overnight.
 9. If a white residue starts to build up in the tank, drain it and clean the tank with strong detergent solution.
 10. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

Dispose of all rinsings in accordance with provincial regulations.

FOLLOW CROPPING

The crops listed below can be safely grown after a spring application of **SAFACYN®**.

Plant Back Crops In case of crop failure, the following crops can be planted in the same season ¹	Rotational Crops The following crops can be planted anytime in the following season
Barley Canary seed Chickpeas Corn (field and sweet) Lentils* Oats Dry field peas Soybean* Wheat (spring, winter & durum)	Barley (spring, winter, malting) Canary seed Canola Chickpeas Corn (field and sweet) Dry common beans Flax Lentils Mustard Oats Dry field peas Soybean Triticale Wheat (spring, winter & durum)
* Rate restrictions apply. Lentils and soybeans can only be grown as plant back crops provided that a maximum product use rate of 113 mL/ha and 156 mL/ha, respectively, was used in the previous crop.	

¹ A second application of **SAFACYN®** cannot be made in the rescue crop.

The crops listed below can be safely grown after a fall application of **SAFACYN®**.

Rotational crops that can be planted in the following spring after application	Rotational crops that can be planted in the second spring after application
---	--

Barley Canary seed Canola Chickpeas Corn (field and sweet) Flax Lentils Oats Dry field peas Soybeans Wheat (spring, winter, and durum)	All crops
--	-----------

SPRAYING INSTRUCTIONS

Thoroughly clean all screens to prevent nozzle clogging. Select proper nozzle to avoid spraying fine mist. Apply in a water volume of 100 to 200 L/ha. For applications to dense weed infestations and thick canopies, use the higher water volume. Consult nozzle manufacturer's recommendation for spray pressures for specific nozzles.

Better coverage of the product results in enhanced control of weeds.

RESTRICTIONS AND LIMITATIONS

1. Wash sprayer thoroughly after use to avoid damage to the next crop sprayed.
2. Field corn – Corn forage and silage can be harvested, used as feed or grazed 60 or more days after application of **SAFACYN®**.
3. Legume forage (chickpeas, faba beans, field peas and lentils) may be used as feed or grazed 60 or more days after application of **SAFACYN®**. Desiccation-treated pea vines may be grazed or fed to livestock.
4. Small grains (wheat, barley and oats) – forage and hay can be used as feed or grazed 30 or more days after application of **SAFACYN®**. Pre-harvest treated barley, wheat and triticale straw may be grazed or fed to livestock.
5. Soybeans may be used as feed or grazed 60 or more days after application of **SAFACYN®**.
6. Soybeans, chickpeas and dry common beans – Do not graze or feed treated hay or straw to livestock when **SAFACYN®** is used as a harvest aid.
7. DO NOT apply directly to water. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
8. As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
9. This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

SPRAY BUFFER ZONES

Field sprayer application: **DO NOT** apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. **DO NOT** apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) Medium classification. Boom height must be 60 cm or less above the crop or ground.

Conventionally piloted aircraft application: **DO NOT** apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) Medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

Apply only by aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate specified for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices or a Global Positioning System (GPS).

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, a long-sleeved shirt, long pants, coveralls, shoes plus socks, and goggles or face shield during mixing/loading, cleanup and repair. Applicators must wear long-sleeved shirt, long pants, and shoes plus socks. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label

recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the registrant at 1-855-264-6262 or obtain technical advice from the distributor or your provincial or territorial agricultural representative.

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,
- low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop/Site		Spray Buffer Zones (metres) Required for the Protection of:				
			Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:		Terrestrial Habitat:
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Lentils (pre-seed or pre-emergent)		0	0	0	0	15
	Soybean (pre-seed or pre-emergent), raspberry, blackberry		0	0	0	0	15
	Barley, canary seed, chickpea, corn, faba beans, oats, dried field peas, wheat (pre-seed or pre-emergent); Chemfallow; Canola, faba beans, flax, lentil, chickpea, soybean, mustard, dried field peas, sunflower, dry common beans, wheat, barley, triticale (harvest aid)		1	0	0	0	15
Aerial	Canola, faba beans, flax, lentil, chickpea, mustard, soybean, dried field peas, sunflower, dry common bean, wheat, barley, triticale (harvest aid)	Fixed wing	0	0	0	0	200
		Rotary wing	0	0	0	0	175

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest

2026-2285
2026-06-26

(most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

WESTERN CANADA DIRECTIONS

**FOR USE IN MANITOBA, SASKATCHEWAN, ALBERTA AND INTERIOR OF
BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY**

REGISTERED USES AND CROPS

PRE-SEED OR PRE-EMERGENT

SAFACYN® is registered for use prior to the following crops as a pre-seed or pre-emergent application.

Barley
Canary seed
Chickpeas
Faba beans
Lentils*
Oats
Peas (dried field)
Wheat (spring, winter and durum)
Corn
Soybeans*

* Rate restrictions apply. Refer to crop specific section for details.

CHEMFALLOW HARVEST AID

SAFACYN® is registered for use as a desiccant in the following crops:

Desiccant
Canola (all types)
Chickpeas
Dry common beans
Faba beans
Flax
Lentils (red lentil varieties only) ¹
Mustard ²
Peas (dried field)
Soybeans
Sunflower

¹ Always tank mix **SAFACYN®** with glyphosate when applying as a desiccant to red lentils. Refer to the **SAFACYN® + GLYPHOSATE TANK MIX CROP USE RATES** section for additional detailed instructions.

² All classes, including brown, oriental, canola quality *Brassica juncea*, *Brassica juncea* varieties with the Clearfield® trait, and yellow mustard.

PREHARVEST WEED MANAGEMENT

SAFACYN® is registered for use as a pre-harvest treatment to improve dry down of volunteer canola (all types including Roundup Ready®), common ragweed, Canada fleabane, redroot pigweed and wild buckwheat in the following crops:

Barley (including feed varieties)
Triticale
Wheat (including durum, spring and winter wheat)

USE RATES

For all **SAFACYN®** solo applications applied pre-seed, pre-emergent or as a chemfallow treatment, use a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.

PRE-SEED OR PRE-EMERGENT

Crop	Use Rate (mL/ha)
Lentils ¹	113
Soybean ²	113 - 156
Barley	113 – 312
Canary seed	113 – 312
Chickpea Kabuli	113 - 312
Corn (field, sweet ³)	113 - 312
Faba beans	113 - 312
Oats	113 - 312
Peas (dried field)	113 - 312
Wheat (spring, durum, winter)	113 - 312
Chemfallow	113 - 312

¹ Rate restrictions apply. Do not use rates higher than 113 mL/ha or injury could result. See crop specific section for additional details.

² Rate restrictions apply. Do not use rates higher than 156 mL/ha or injury could result. See crop specific section for additional details. Some soybeans cultivars may be more sensitive to saflufenacil and injury might occur.

³ Some sweet corn hybrids may be more sensitive to saflufenacil and injury might occur.

CHEMFALLOW

Site	Use Rate (mL/ha)	Application Timing
Chemfallow	113 - 312	Post-emergence in fallow croplands and post-harvest.

HARVEST AID

SAFACYN® may be used as a harvest aid to accelerate the rate of crop dry down and improve crop uniformity to facilitate direct combining. Early application may result in yield and/or seed quality loss.

The dry down of crops will be best under favorable environmental conditions like warm temperatures, good moisture conditions and low humidity.

Harvesting of crops can be done when plant material is dry and seed moisture level allows efficient harvesting. Under ideal conditions, harvest can normally commence within 7-14 days after desiccation when applied at the appropriate crop stage recommendation. Adverse weather conditions such as rainfall, cool temperatures, shorter day length and high humidity may slow the plant desiccation and keep seed moisture levels high which can delay commencement of harvest after the **SAFACYN®** application. Consult your ADAMA representative for further information on the timing of harvest after a pre-harvest application.

Crop	Use Rate (mL/ha)
Canola (all types)	227
Faba beans	
Chickpeas	
Dry common beans	
Flax	
Lentils ¹	
Mustard ²	
Peas (dried field),	
Soybeans	
Sunflower	

¹ Apply only to red lentil varieties. Always tank mix **SAFACYN®** with glyphosate when applying as a desiccant to red lentils. Refer to the **SAFACYN® + GLYPHOSATE TANK MIX CROP USE RATES** section for additional detailed instructions.

² All classes, including brown, oriental, canola quality *Brassica juncea*, *Brassica juncea* varieties with the Clearfield® trait, and yellow mustard.

Aerial Application

SAFACYN® can be used for aerial application when used as a harvest aid or a pre-harvest weed management in the registered crops.

HARVEST AID - CROP SPECIFIC RECOMMENDATIONS STAND-ALONE

DESICCATION - DRY COMMON BEANS / SOYBEANS

Rate	227 mL/ha of SAFACYN® + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when stems are green to brown in colour and pods are mature (yellow – brown) and 80 – 90% of the original leaves have dropped.

Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application, use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
Remarks	Do not graze or feed treated hay or straw to livestock.

DESICCATION - CHICKPEAS

Rate	227 mL/ha of SAFACYN® + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	For Desi type, apply at the time swathing would normally commence, when the majority of plants are yellow and most pods are mature and seeds have turned from green to yellow or brown. Upper part of plant may still be green. For Kabuli type, apply when the majority of plants and pods are ripe and dry with seeds turned from green to white or tan, and detached from the pods. Dry down is less complete in Kabuli type due to its thick pod wall.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
Remarks	Do not graze or feed treated hay or straw to livestock.

DESICCATION – FIELD PEAS

Rate	227 mL/ha of SAFACYN® + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when the majority of pods are brown (70 – 80%).
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
Remarks	Desiccation treated pea vines may be grazed or fed to livestock.

DESICCATION – SUNFLOWER

Rate	227 mL/ha of SAFACYN® + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic

	surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when the backs of the heads and bracts are turning yellow, and seed moisture is 20 – 30%.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.

DESICCATION – CANOLA (ALL TYPES) AND MUSTARD¹

Rate	227 mL/ha of SAFACYN® + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when 65 – 80% of seeds have changed colour.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
Remarks	It is recommended that the application of SAFACYN® as a desiccant in canola and mustard be made to shatter resistant varieties. Under ideal conditions, harvest can normally commence within 14-21 days after desiccation when applied at the appropriate crop stage recommendation. Adverse weather conditions such as rainfall, cool temperatures, shorter day length and high humidity may slow the plant desiccation and keep seed moisture levels high which can delay commencement of harvest after the SAFACYN® application.

¹ All classes, including brown, oriental, canola quality *Brassica juncea*, *Brassica juncea* varieties with the Clearfield® trait, and yellow mustard.

DESICCATION – FLAX

Rate	227 mL/ha of SAFACYN® + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when 75% of bolls have turned colour.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.

DESICCATION – FABA BEANS

Rate	227 mL/ha of SAFACYN® + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when 80% of lower pods have turned black, middle pods have turned yellow/tan, and top green pods have firm seed.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
Remarks	Do not graze or feed treated hay or straw to livestock.

PRE-HARVEST WEED MANAGEMENT

SAFACYN® may be used as a pre-harvest treatment in wheat (including durum, spring, and winter wheat), barley (including feed varieties) and triticale to improve dry down of volunteer canola (all types including Roundup Ready®), common ragweed, Canada fleabane, redroot pigweed and wild buckwheat and to facilitate direct combining. Early application may result in yield loss.

SAFACYN® when used alone as a pre-harvest treatment will not affect the seed germination if applied according to label recommendations.

WHEAT, BARLEY AND TRITICALE

Rate	156 – 227 mL/ha of SAFACYN® + adjuvant
Water Volume	100 – 200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Hard dough stage; a thumbnail impression remains on seed; less than 30% moisture.
Weeds Controlled	Improved dry down of volunteer canola (all types including Roundup Ready®), common ragweed, Canada fleabane, redroot pigweed and wild buckwheat.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
Remarks	Treated barley, wheat and triticale straw may be grazed or fed to livestock. Use higher water volume for dense crop stands and higher weed pressure.

WEEDS CONTROLLED

SAFACYN® can be applied pre-seed or pre-emergence to the crop to control weeds listed below.

SAFACYN® should always be applied in combination with glyphosate for broad spectrum weed control including grassy weeds. Refer to the **SAFACYN® + GLYPHOSATE TANK MIX CROP USE RATES** section for additional detailed instructions.

SAFACYN® ALONE

For Rapid Burndown (Pre-seed or Pre-emergent)

Use Rate: 113 mL/ha SAFACYN®

Weed	Maximum Weed Stage
Kochia	15 cm
Canada fleabane	8 leaf
Cleavers	4-whorl stage
Lamb's-quarters	8 leaf
Narrow-leaved hawk's beard	8 cm
Redroot pigweed	8 leaf
Round-leaved mallow	8 leaf
Stinkweed	8 leaf
Volunteer canola (all types including Roundup Ready®)	8 leaf
Wild buckwheat	8 leaf
Wild mustard	8 leaf

Use Rate: 156 mL/ha SAFACYN®

SAFACYN® applied pre-seed or pre-emergence at 156 mL/ha will provide rapid burndown control of the following weeds in addition to those listed above:

Weed	Maximum Weed Stage
Common ragweed	8 leaf
Giant ragweed	8 leaf
Lady's-thumb	6 leaf
Perennial sow-thistle (top growth burndown control)	8 leaf
Prickly lettuce (top growth only)	9 leaf
Shepherd's-purse	Full flower

For Rapid Burndown and Suppression of Secondary Weed Flushes (Pre-seed or Pre-emergent)

Use Rate: 227 - 312 mL/ha SAFACYN®

Weed	Maximum Weed Stage
Redroot pigweed	8 leaf

Stinkweed	8 leaf
Volunteer canola (all types including Roundup Ready®)	8 leaf
Wild buckwheat	8 leaf
Wild mustard	8 leaf

TANK MIXTURES

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

TANK MIXTURES – PRE-SEED, PRE-EMERGENCE AND CHEMFALLOW

SAFACYN® + GLYPHOSATE

For broad spectrum weed control, **SAFACYN®** should always be tank mixed with glyphosate present as isopropylamine salt, di-ammonium salt or potassium salt. **SAFACYN®** is compatible with all liquid glyphosate formulations in which glyphosate is present as isopropylamine salt, di-ammonium salt or potassium salt.

Tank mixing **SAFACYN®** with glyphosate will provide control of all weeds controlled by glyphosate in addition to the broadleaf weeds listed on the **SAFACYN®** label. Consult the glyphosate label for a complete list of weeds controlled by glyphosate.

Tank mixing is a recognized strategy to delay herbicide resistance as well as improve the weed spectrum controlled.

Refer to the **SAFACYN® + GLYPHOSATE TANK MIX CROP USE RATES** section for additional detailed instructions.

Do not exceed 900 g a.e./ha of GLYPHOSATE rate when tank mixing with **SAFACYN®**.

TANK MIX CROP USE RATES

SAFACYN® + GLYPHOSATE

Crop	Use Rate	
	SAFACYN® (mL/ha)	GLYPHOSATE ⁴ (g a.e./ha)

Lentils ¹	113	450 – 900
Soybean ²	113 - 156	450 – 900
Barley	113 - 312	450 – 900
Canary seed	113 - 312	450 – 900
Chickpea Kabuli	113 - 312	450 – 900
Corn (field, sweet ³)	113 - 312	450 – 900
Faba beans	113 - 312	450 – 900
Oats	113 - 312	450 – 900
Peas (dried field)	113 - 312	450 – 900
Wheat (spring, durum, winter)	113 - 312	450 – 900
Chemfallow	113 - 312	450 – 900

¹ Rate restrictions apply. Do not use rates higher than 113 mL/ha or crop injury may result. See crop specific section for additional details.

² Rate restrictions apply. Do not use rates higher than 156 mL/ha or crop injury may result. See crop specific section for additional details. Some soybeans cultivars may be more sensitive to saflufenacil and injury might occur.

³ Some sweet corn hybrids may be more sensitive to saflufenacil and injury might occur.

⁴ Must use a minimum of 100 L/ha (ground application) of water volume and a maximum GLYPHOSATE rate of 900 g a.e./ha.

For tank mix applications of **SAFACYN®** with glyphosate, use a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.

Do not apply pre-seed, pre-emergent and chemfallow applications using aerial application equipment.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

TANK MIX WEEDS CONTROLLED

Rapid Burndown (Pre-seed or Pre-emergent)

SAFACYN® when tank mixed with glyphosate will provide rapid burndown of the following weeds in addition to those weeds controlled by **SAFACYN®** applied alone.

Use Rate: 113 mL/ha SAFACYN® + 450 – 900 g a.e./ha GLYPHOSATE

Weed	Maximum Weed Stage
Dandelion (top growth burndown control only)	15 cm
Flixweed	8 leaf

CROP SPECIFIC RECOMMENDATIONS - PRE-SEED OR PRE-EMERGENT

CHICKPEAS, FABA BEANS AND PEAS (dried field)

Timing	Pre-seed or Pre-emergent
Rate	113 – 312 mL/ha of SAFACYN® + 450 - 900 g a.e./ha glyphosate + adjuvant
Water Volume	Must use a minimum of 100 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Weeds Controlled	Refer to the Weeds Controlled section for a complete list of weeds controlled by SAFACYN® . Refer to the glyphosate label for weeds controlled in addition to those controlled by SAFACYN® .

CEREALS – BARLEY, CANARY SEED, OATS AND WHEAT (spring, durum, winter)

Timing	Pre-seed or Pre-emergent
Rate	113 – 312 mL/ha of SAFACYN® + 450 - 900 g a.e./ha glyphosate + adjuvant
Water Volume	Must use a minimum of 100 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Weeds Controlled	Refer to the Weeds Controlled section for a complete list of weeds controlled by SAFACYN® . Refer to the glyphosate label for weeds controlled in addition to those controlled by SAFACYN® .

SOYBEANS

Timing	Pre-seed or Pre-emergent
Rate	113 - 156 mL/ha of SAFACYN® + 450 - 900 g a.e./ha glyphosate + adjuvant
Water Volume	Must use a minimum of 100 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Weeds Controlled	Refer to the Weeds Controlled section for a complete list of weeds controlled by SAFACYN® . Refer to the glyphosate label for weeds controlled in addition to those controlled by SAFACYN® .
Remarks	Some soybean cultivars may be more sensitive to saflufenacil and injury might occur. When applying 156 mL/ha pre-emergent to soybeans, DO NOT apply to coarse textured soils with less than 2% organic matter. When applying pre-emergent to soybeans, apply prior to when the soybeans cause the ground to crack and no more than 3 days after planting.

LENTILS (including Clearfield® Lentils)

Timing	Pre-seed or Pre-emergent
Rate	113 mL/ha of SAFACYN® + 450 - 900 g a.e./ha glyphosate + adjuvant
Water Volume	Must use a minimum of 100 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Weeds Controlled	Refer to the Weeds Controlled section for a complete list of weeds controlled by SAFACYN® . Refer to the glyphosate label for weeds controlled in addition to those controlled by SAFACYN® .
Remarks	Rainfall shortly after product application can result in slight injury to the crop. Lentils will be more susceptible to injury on coarse texture and low organic matter soils. Injury will usually appear as leaf tissue necrosis on the outer edges of the leaves. Lentils will grow out of injury symptoms, and yield will not be impacted at recommended rates. The user should contact ADAMA before applying any other soil applied herbicide with, before, or after applications of SAFACYN® + glyphosate. The addition of other soil applied herbicides may increase the sensitivity of lentils to SAFACYN® and injury may result.

CORN (field, sweet)

Timing	Pre-seed or Pre-emergent
Rate	113 – 312 mL/ha of SAFACYN® + 450 - 900 g a.e./ha glyphosate + adjuvant
Water Volume	Must use a minimum of 100 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Weeds Controlled	Refer to the Weeds Controlled section for a complete list of weeds controlled by SAFACYN® . Refer to the glyphosate label for weeds controlled in addition to those controlled by SAFACYN® .
Remarks	Some sweet corn hybrids may be more sensitive to saflufenacil and injury might occur.

CHEMFALLOW

Timing	Chemfallow
Rate	113 – 312 mL/ha of SAFACYN® + 450 - 900 g a.e./ha glyphosate + adjuvant
Water Volume	Must use a minimum of 100 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Weeds Controlled	Refer to the Weeds Controlled section for a complete list of weeds controlled by SAFACYN® . Refer to the glyphosate label for weeds

	controlled in addition to those controlled by SAFACYN® .
Remarks	Apply to actively growing weeds less than 15 cm in height. Better coverage of the product results in enhanced control of weeds. For application to larger weeds or dense weed infestations, use minimum water volume of 100 L per hectare.

HARVEST AID - TANK-MIX CROP SPECIFIC RECOMMENDATIONS

SAFACYN® may be used as a harvest aid to accelerate the rate of crop dry down and improve crop uniformity to facilitate direct combining. Early application may result in yield and/or seed quality loss.

The dry down of crops will be best under favorable environmental conditions like warm temperatures, good moisture conditions and low humidity.

Harvesting of crops can be done when plant material is dry and seed moisture level allows efficient harvesting. Under ideal conditions, harvest can normally commence within 7-14 days after desiccation when applied at the appropriate crop stage recommendation. Adverse weather conditions such as rainfall, cool temperatures, shorter day length and high humidity may slow the plant desiccation and keep seed moisture levels high which can delay commencement of harvest after the **SAFACYN®** application. Consult your ADAMA representative for further information on the timing of harvest after a pre-harvest application.

SAFACYN® may be applied in tank mix with glyphosate for additional pre-harvest weed control.

DESICCATION - DRY COMMON BEANS / SOYBEANS

Timing	Desiccation of DRY COMMON BEANS / SOYBEANS
Rate	227 mL/ha of SAFACYN® + 900 g a.e./ha glyphosate* (e.g. 1.67 L/ha of 540 g/L glyphosate formulation) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when stems are green to brown in colour and pods are mature (yellow – brown) and 80 – 90% of the original leaves have dropped.
Directions for Use – Ground ONLY	Ground application may be done with a standard boom sprayer.
Remarks	Do not graze or feed treated hay or straw to livestock. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.

DESICCATION - CHICKPEAS

Timing	Desiccation of CHICKPEAS
--------	--------------------------

Rate	227 mL/ha of SAFACYN® + 900 g a.e./ha glyphosate (e.g. 1.67 L/ha of 540 g/L glyphosate formulation) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	For Desi type, apply at the time swathing would normally commence, when the majority of plants are yellow and most pods are mature and seeds have turned from green to yellow or brown. Upper part of plant may still be green. For Kabuli type, apply when the majority of plants and pods are ripe and dry with seeds turned from green to white or tan, and detached from the pods. Dry down is less complete in Kabuli type due to its thick pod wall.
Directions for Use – Ground ONLY	Ground application may be done with a standard boom sprayer.
Remarks	Do not graze or feed treated hay or straw to livestock. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.

DESICCATION - RED LENTIL VARIETIES

Timing	Desiccation of RED LENTIL VARIETIES
Rate	227 mL/ha of SAFACYN® + 900 g a.e./ha glyphosate (e.g. 1.67 L/ha of 540 g/L glyphosate formulation) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when lowermost pods (bottom 15%) are brown and rattle when shaken.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Remarks	Apply only to red lentil varieties. Do not graze or feed treated hay or straw to livestock. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.

DESICCATION – FIELD PEAS

Timing	Desiccation of FIELD PEAS
Rate	227 mL/ha of SAFACYN® + 900 g a.e./ha glyphosate (e.g. 1.67 L/ha of 540 g/L glyphosate formulation) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate

	(COC) (e.g. Merge Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when the majority of pods are brown (70–80%).
Directions for Use – Ground ONLY	Ground application may be done with a standard boom sprayer.
Remarks	Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.

DESICCATION – CANOLA (ALL TYPES) AND MUSTARD¹

Timing	Desiccation of CANOLA (all types) and MUSTARD
Rate	227 mL/ha of SAFACYN® + 900 g a.e./ha glyphosate (e.g. 1.67 L/ha of 540 g/L glyphosate formulation) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when 65 – 80% of seeds have changed colour.
Directions for Use – Ground ONLY	Ground application may be done with a standard boom sprayer.
Remarks	It is recommended that the application of SAFACYN® as a desiccant in canola and mustard be made to shatter resistant varieties. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes. Under ideal conditions, harvest can normally commence within 14-21 days after desiccation when applied at the appropriate crop stage recommendation. Adverse weather conditions such as rainfall, cool temperatures, shorter day length and high humidity may slow the plant desiccation and keep seed moisture levels high which can delay commencement of harvest after the SAFACYN® application.

¹ All classes, including brown, oriental, canola quality *Brassica juncea*, *Brassica juncea* varieties with the Clearfield® trait, and yellow mustard.

DESICCATION – FLAX

Timing	Desiccation of FLAX
Rate	227 mL/ha of SAFACYN® + 900 g a.e./ha glyphosate (e.g. 1.67 L/ha of 540 g/L glyphosate formulation) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when 75% of bolls have turned colour.
Directions for Use – Ground ONLY	Ground application may be done with a standard boom sprayer.

Remarks	Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.
---------	--

DESICCATION – FABA BEANS

Timing	Desiccation of FABA BEANS
Rate	227 mL/ha of SAFACYN® + 900 g a.e./ha glyphosate (e.g. 1.67 L/ha of 540 g/L glyphosate formulation) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when 80% of lower pods have turned black, middle pods have turned yellow/tan, and top green pods have firm seed.
Directions for Use – Ground ONLY	Ground application may be done with a standard boom sprayer.
Remarks	Do not graze or feed treated hay or straw to livestock. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.

PRE-HARVEST WEED MANAGEMENT – TANK MIX CROP SPECIFIC RECOMMENDATIONS

SAFACYN® may be used as a pre-harvest treatment in wheat (including durum, spring and winter wheat), barley (including feed varieties) and triticale to improve dry down of volunteer canola (all types including Roundup Ready), common ragweed, Canada fleabane, redroot pigweed and wild buckwheat. Tank mixing **SAFACYN®** with glyphosate will provide additional pre-harvest weed management to the crop dry down provided by glyphosate.

The tank mix with glyphosate may affect the seed germination. Tank mixing with glyphosate is not recommended when harvested grain is to be used for seed.

WHEAT AND BARLEY

Rate	156 – 227 mL/ha of SAFACYN® + 900 g a.e./ha of glyphosate + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Hard dough stage; a thumbnail impression remains on seed; less than 30% moisture.
Weeds Controlled	Improved dry down of volunteer canola (all types including Roundup Ready®), common ragweed, Canada fleabane, redroot pigweed and wild

	buckwheat. Consult the glyphosate label regarding additional pre-harvest weed control.
Directions for Use – Ground ONLY	Ground application may be done with a standard boom sprayer.
Remarks	Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes. Use higher water volume for dense crop stands and higher weed pressure. Tank mixing with glyphosate is not recommended when harvested grain is to be used for seed.

PRE-HARVEST INTERVAL (PHI)

The following pre-harvest intervals should be observed for respective crops when **SAFACYN®** is used as a pre-seed or pre-emergent application.

Crop	PHI (days)
Barley	60
Canary seed	60
Chickpea	60
Corn (field, sweet)	60
Faba beans	60
Lentils	60
Oats	60
Peas (dried field)	60
Soybean	60
Wheat (spring, winter, durum)	60

The following pre-harvest intervals should be observed for respective crops when **SAFACYN®** is used as a harvest aid or a pre-harvest treatment for weed management.

Crop	PHI (days)
Barley	3
Canola	3
Chickpeas	2
Dry common beans	2
Faba beans	2
Flax	3
Lentils	3
Mustard	3
Peas (dried field)	3
Soybeans	3
Sunflower	7
Triticale	3
Wheat	3

EASTERN CANADA DIRECTIONS

FOR SALE FOR USE ON RASPBERRY AND BLACKBERRY IN EASTERN CANADA AND BRITISH COLUMBIA (EXCLUDING THE PEACE RIVER REGION), AND ON ALL OTHER LABELLED CROPS IN EASTERN CANADA ONLY

REGISTERED USES AND CROPS

SAFACYN® is registered for use in the following crops as a pre-plant or pre-emergent application. SAFACYN® may also be used as a harvest aid in soybeans, dry beans and canola, and as a pre-harvest weed management treatment in wheat, barley and triticale.

Pre-plant or Pre-emergent
Barley
Oats
Wheat (spring, winter and durum)
Corn (field and sweet)

Pre-plant
Soybean

Harvest Aid
Desiccation of soybeans
Desiccation of dry beans
Desiccation of canola (all types)

Pre-harvest Weed Management
Wheat (including durum, spring and winter wheat)
Barley (including feed varieties)
Triticale

DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT apply by air.

EARLY SEASON WEED CONTROL APPLICATIONS

CROP USE RATES – SAFACYN®

Pre-plant or Pre-emergent

Crop	Use Rate (mL/ha)	Application Timing
Barley	156 - 312	Pre-plant or Pre-emergent
Oats	156 - 312	Pre-plant or Pre-emergent
Wheat (spring, winter, durum)	156 - 312	Pre-plant or Pre-emergent
Soybean	156*	Pre-plant ONLY
Corn	312**	Pre-plant or Pre-emergent

* Do not use rates higher than 156 mL/ha or crop injury may result. For use in combination with glyphosate. Refer to the Crop Specific Recommendations section of this label for additional details. Some soybeans cultivars may be more sensitive to saflufenacil and injury might occur.

** 312 mL/ha will only provide short season weed control and should be used in a program where a later application of a herbicide is anticipated. Refer to the Crop Specific Recommendations section of this label for additional details. Some sweet corn hybrids may be more sensitive to saflufenacil and injury might occur.

Refer to Weeds Controlled section below for further application information.

WEEDS CONTROLLED

Pre-plant or Pre-emergent Burndown - CEREALS and Pre-plant - SOYBEAN

SAFACYN® may be applied pre-plant or pre-emergent to cereals (barley, oats, wheat) and pre-plant to soybeans for control (burndown) of the following emerged broadleaf weeds. It is recommended that SAFACYN® is always tank mixed with glyphosate for broad spectrum weed control. Refer to the SAFACYN® + GLYPHOSATE TANK MIX CROP USE RATES section for additional detailed instructions.

Use Rate: 156 mL/ha SAFACYN®

Weed	Maximum Weed Stage
Broadleaf plantain ¹	beginning of flowering
Canada fleabane ²	20 leaf
Common ragweed ²	8 leaf
Dandelion (suppression only)	6 leaf
Giant ragweed ^{1,2}	8 leaf
Lady's-thumb ¹	6 leaf
Lamb's-quarters	8 leaf
Perennial sow thistle (top growth burndown control) ¹	8 leaf
Prickly lettuce (top growth only) ¹	9 leaf
Redroot pigweed	8 leaf
Shepherd's-purse ¹	full flower
Stinkweed ¹	full flower
Wild buckwheat ¹	6 leaf
Wild mustard	8 leaf

¹ Controlled with a tank mix of SAFACYN® and glyphosate

² Includes glyphosate-resistant biotypes

Pre-plant Burndown and Suppression of Secondary Weed Flushes – CEREALS

Use Rate: 312 mL/ha SAFACYN®

Weed	Maximum Weed Stage
Lamb's-quarters	8 leaf
Redroot pigweed	8 leaf
Stinkweed	8 leaf
Wild buckwheat	8 leaf
Wild mustard	8 leaf

Pre-plant or Pre-emergent Burndown – CORN

SAFACYN® at 312 mL/ha may be tank mixed with glyphosate and applied to corn prior to planting or after planting prior to crop emergence.

SAFACYN® + glyphosate at 900 g a.e./ha will provide control (burndown) of the following emerged broadleaf weeds in addition to those controlled by glyphosate.

Use Rate: 312 mL/ha SAFACYN®

Weed	Maximum Weed Stage
Common ragweed*	8 leaf
Lamb's-quarters	8 leaf
Redroot pigweed	8 leaf
Velvetleaf	8 leaf
Wild buckwheat	8 leaf
Wild mustard	8 leaf

*Includes glyphosate-resistant biotypes

TANK MIXTURES

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

For broad spectrum burndown of weeds in soybeans, barley, oats and wheat (spring, winter and durum), SAFACYN® should always be tank mixed with glyphosate present as isopropylamine

salt, di-ammonium salt or potassium salt. Refer to the **SAFACYN® + GLYPHOSATE TANK MIX CROP USE RATES** section for additional detailed instructions. **SAFACYN®** is compatible with all liquid glyphosate formulations, in which glyphosate is present as isopropylamine salt, di-ammonium salt or potassium salt, as a tank-mix partner. Apply to weeds less than 15 cm in height, according to the glyphosate label. For extended grass and broadleaf control in soybeans, Pursuit® Herbicide or Phantom® 240 SL may be tank mixed with **SAFACYN®** and glyphosate.

Tank mixing is a recognized strategy to delay herbicide resistance as well as improve the weed spectrum controlled.

See the Crop Specific Recommendations section of this label for further details, including rates, adjuvant recommendations and restrictions.

Refer to the label of the tank mix partner for further precautions, use instructions and crop rotation restrictions. When a tank mix is used, consult the labels of the tank mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

CROP SPECIFIC RECOMMENDATIONS - PRE-PLANT OR PRE-EMERGENT

CEREALS – BARLEY, OATS AND WHEAT (spring, winter and durum)

Timing	Pre-plant or Pre-emergent
Rate	156 - 312 mL/ha of SAFACYN® + 900 g a.e./ha of glyphosate
Water Volume	100 – 200 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Weeds Controlled	Refer to the Weeds Controlled section of this label for a complete list of weeds controlled by SAFACYN® . Refer to the glyphosate label for weeds controlled in addition to those controlled by SAFACYN® .

SOYBEANS

SAFACYN® + Glyphosate

Timing	Pre-plant ONLY
Rate	156 mL/ha of SAFACYN® + 900 g a.e./ha of glyphosate
Water Volume	100 – 200 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Weeds Controlled	Refer to the Weeds Controlled section of this label for a complete list of weeds controlled by SAFACYN® . Refer to the glyphosate label for weeds controlled in addition to those controlled by SAFACYN® .
Remarks	Some soybeans cultivars may be more sensitive to saflufenacil and injury

	might occur. For Pre-plant timing, apply pre-plant from 21 days prior to planting up to the day of planting.
--	---

SAFACYN® + Glyphosate + Imazethapyr

Timing	Pre-plant ONLY
Rate	156 mL/ha of SAFACYN® + 900 g a.e./ha of glyphosate + 420 mL/ha Pursuit® Herbicide or Phantom® 240 SL.
Water Volume	100 – 200 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Weeds Controlled	Refer to the Weeds Controlled section of this label for a complete list of weeds controlled by SAFACYN® . Refer to the glyphosate label for weeds controlled in addition to those controlled by SAFACYN® .
Remarks	Some soybeans cultivars may be more sensitive to saflufenacil and injury might occur. For Pre-plant timing, apply pre-plant from 21 days prior to planting up to the day of planting.

CORN (field and sweet)

SAFACYN®

Timing	Pre-emergent
Rate	312 – 624 mL/ha of SAFACYN®
Water Volume	100 – 200 L/ha
Weeds Controlled	Refer to the Weeds Controlled section of this label for a complete list of weeds controlled by SAFACYN® .
Remarks	Some sweet corn hybrids may be more sensitive to saflufenacil and injury might occur.

SAFACYN® + Glyphosate

Timing	Pre-plant or Pre-emergent Burndown
Rate	312 mL/ha of SAFACYN® + 900 g a.e./ha of glyphosate
Water Volume	100 – 200 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Weeds Controlled	Refer to the Weeds Controlled section of this label for a complete list of weeds controlled by SAFACYN® . Refer to the glyphosate label for weeds controlled in addition to those controlled by SAFACYN® . 312 mL/ha will only provide short season weed control and should be

	used in a program where a later application of a herbicide is anticipated.
Remarks	Apply tank mixture prior to or after planting but before crop emergence. Some sweet corn hybrids may be more sensitive to saflufenacil and injury might occur.

HARVEST AID / DESICCATION IN SOYBEANS, DRY BEANS AND CANOLA (ALL TYPES)

SAFACYN® may be used as a harvest aid to accelerate the rate of dry down for crop and green weeds and improve crop uniformity to facilitate direct combining of soybeans, dry beans and canola. Early application may result in yield loss.

The dry down of crops will be best under favorable environmental conditions like warm temperatures, good moisture conditions and high humidity.

Harvesting of crops can be done when plant material is dry and seed moisture level allows efficient harvesting. For most crops, harvest can normally commence within 3-10 days after desiccation. Adverse weather conditions such as rainfall, cool temperatures and high humidity may slow the plant desiccation and keep seed moisture levels high which can delay commencement of harvest after the **SAFACYN®** application.

SAFACYN® when used as a harvest aid will not affect seed germination if applied according to label recommendations. Applications sprayed prior to label application timing may impact seed germination.

For the harvest aid/desiccation in soybeans and dry beans, do not graze or feed treated hay or straw to livestock.

SAFACYN® – Soybeans and Dry Beans

Timing	Desiccation of Soybeans and Dry Beans
Rate	156 - 312 mL/ha of SAFACYN®
Water Volume	200 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when 90% of pods have changed colour from green to yellow or brown.
Remarks	Ground application may be done with a standard boom sprayer. Do not graze or feed treated hay or straw to livestock.

SAFACYN® + Glyphosate – Soybeans and Dry Beans

SAFACYN® may be applied in tank mix with glyphosate for additional pre-harvest weed control.

The tank mix with glyphosate may affect seed germination. DO NOT tank mix with

glyphosate on fields grown for seed production.

Timing	Desiccation of Soybeans and Dry Beans
Rate	156 - 312 mL/ha of SAFACYN® + 900 g a.e./ha of glyphosate
Water Volume	200 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when 90% of pods have changed colour from green to yellow or brown.
Remarks	Ground application may be done with a standard boom sprayer. Do not graze or feed treated hay or straw to livestock. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.

SAFACYN® – CANOLA (All types)

Timing	Desiccation of Canola (all types)
Rate	156 - 312 mL/ha of SAFACYN®
Water Volume	200 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when 60 – 75% of seeds have changed colour.
Remarks	The use of SAFACYN® as a desiccant in canola may cause shattering. Ground application may be done with a standard boom sprayer.

SAFACYN® + Glyphosate – CANOLA (All types)

SAFACYN® may be applied in tank mix with glyphosate for additional pre-harvest weed control.

The tank mix with glyphosate may affect seed germination. DO NOT tank mix with glyphosate on fields grown for seed production.

Timing	Desiccation of Canola (all types)
Rate	156 - 312 mL/ha of SAFACYN® + 900 g a.e./ha of glyphosate
Water Volume	200 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Apply when 60 – 75% of seeds have changed colour.
Remarks	The use of SAFACYN® as a desiccant in canola may cause shattering. Ground application may be done with a standard boom sprayer. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling

	applies to tank mixes.
--	------------------------

PRE-HARVEST WEED MANAGEMENT IN WHEAT, BARLEY, AND TRITICALE

SAFACYN® may be used as a pre-harvest weed management treatment in wheat (including spring, durum, and winter wheat), barley (including feed varieties) and triticale to improve dry down of volunteer canola (all types including Roundup Ready®), common ragweed, Canada fleabane, redroot pigweed and wild buckwheat and to facilitate direct combining. Early application may result in yield loss.

SAFACYN® when used alone as a pre-harvest weed management treatment will not affect seed germination if applied according to label recommendations. Applications sprayed prior to label application timing may impact seed germination.

SAFACYN® – Wheat, Barley and Triticale

Rate	156 - 312 mL/ha of SAFACYN®
Water Volume	200 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Hard dough stage; a thumbnail impression remains on seed; less than 30% moisture.
Weeds Controlled	Improved dry down of volunteer canola (all types including Roundup Ready®), common ragweed*, Canada fleabane*, redroot pigweed and wild buckwheat.
Remarks	Ground application may be done with a standard boom sprayer. Use higher water volume for dense crop stands and higher weed pressure. Treated barley, wheat and triticale straw may be grazed or fed to livestock.

SAFACYN® + Glyphosate – Wheat and Barley

SAFACYN® may be applied in tank mix with glyphosate for additional pre-harvest weed management to the crop dry down provided by glyphosate herbicide.

The tank mix with glyphosate may affect seed germination. DO NOT tank mix with glyphosate on fields grown for seed production.

Rate	156 - 312 mL/ha of SAFACYN® + 900 g a.e./ha of glyphosate
Water Volume	100 – 200 L/ha
Surfactant/Adjuvant	Add a 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge® Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Application Timing	Hard dough stage; a thumbnail impression remains on seed; less than 30% moisture.
Weeds Controlled	Improved dry down of volunteer canola (all types including Roundup

	Ready®), common ragweed*, Canada fleabane*, redroot pigweed and wild buckwheat.
Remarks	Ground application may be done with a standard boom sprayer. Use higher water volume for dense crop stands and higher weed pressure. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.

* Includes glyphosate-resistant biotypes

MINOR USE CROPS – RASPBERRY AND BLACKBERRY

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than ADAMA Agricultural Solutions Canada Ltd. under the User Requested Minor Use Label Expansion program. For these uses, ADAMA Agricultural Solutions Canada Ltd. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

Raspberry and Blackberry (Eastern Canada and British Columbia, excluding the Peace River Region)

Use as a directed spray. Apply to dormant raspberry and blackberry as a banded application directed at the base of the crops while targeting emerged broadleaf weeds. **DO NOT allow spray drift to contact dormant canes as severe crop injury will result.** Use ground equipment only. DO NOT apply banded treatments using high pressure hand wands. Maximum of 1 application per season.

Rate	156 mL/ha of SAFACYN®
Surfactant/Adjuvant	Add 30% phosphate ester surfactant, MSO adjuvant, Crop Oil Concentrate (COC) (e.g. Merge Adjuvant) at 0.5 – 1% v/v or a non-ionic surfactant (NIS) (e.g. Adama Adjuvant 80) at 0.25% v/v.
Water Volume	Minimum 100 L/ha
Application Timing	Early post-emergent application directed to the base of the raspberry or blackberry plant prior to the plants breaking winter dormancy. One application per season. DO NOT allow spray drift to contact dormant canes.
Weeds Controlled	Labelled broadleaf weeds as listed in the Weeds Controlled section of this label (see weeds controlled at the 156 mL/ha use rate).

PRE-HARVEST AND GRAZING INTERVALS (PHI)

The following pre-harvest and grazing intervals should be observed for respective crops when SAFACYN® is applied as an early season weed control application.

Crop	PHI (days)
Barley	60 ¹
Corn (field, sweet)	60 ²
Soybean	60 ³
Oats	60 ¹
Wheat (spring, winter, durum)	60 ¹

¹ For barley, oats and wheat, forage and hay can be used as feed or grazed 30 days or more after an early season weed control application of **SAFACYN®**.

² Corn forage and silage can be used as feed or grazed 60 days or more after an early season weed control application of **SAFACYN®**.

³ Soybeans may be fed or grazed 60 days or more after an early season weed control application of **SAFACYN®**.

Harvest Aid and Pre-harvest Weed Management

The following pre-harvest intervals should be observed for respective crops when **SAFACYN®** is used as a harvest aid and pre-harvest weed management.

Crop	PHI (days)
Dry beans	2 ¹
Soybeans	3 ¹
Canola	3
Wheat	3 ²
Barley	3 ²
Triticale	3 ²

¹ Do not graze or feed treated hay or straw to livestock.

² For wheat, barley and triticale, straw can be used as feed or grazed 3 days or more after a pre-harvest weed management application of **SAFACYN®**.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, **SAFACYN®** is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to **SAFACYN®** and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **SAFACYN®** or other Group 14 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as

effectively as the more resistance-prone partner.

- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 or at adama.com/canada.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

SAFACYN is a registered trademark of an ADAMA Group Company.
All other trademarks are the property of their respective owners.