

Active Ingredient:

Azoxystrobin 143 g/L and Propiconazole 124 g/L = SC

Application Rates and Acres Treated:

Rate: 210 – 620 ml/ac

Acres Treated: 14 - 40 ac/jug

Packaging:

Case: 2 × 8.6 L jugs

Water Volume:

Ground: 40 L/ac

(10 US gal/ac)

Aerial: 20 L/ac (5 US gal/ac)

Rainfastness:

Avoid applying when heavy rainfall is in the forecast.

KEY DISEASES CONTROLLED:

- Anthracnose
- Ascochyta Blight
- Barley Leaf Rust
- Crown Rust
- Mycosphaerella Blight
- Net and Spot Blotches
- Powdery Mildew
- ¹Suppression only

- Scald
- Septoria Spot
- Stripe Rust
- Tan Spot
- Wheat Leaf Rust
- White Mould¹

HOW IT WORKS:

Used as both a curative and preventative fungicide, $TOPNOTCH^{\text{\tiny M}}$ has broad-spectrum, systemic and contact activity.

APPLICATION TIMING AND CROP STAGING:

Crop	Diseases	Application Timing	Rate
Barley	Barley Net Blotch, Barley Scald, Septoria Leaf Spot, Stripe Rust, Barley Leaf Rust, Tan Spot	Apply once between stem elongation and half-head emergence.	210 ml/ac
Beans, Field peas, Lentils, Soybeans	Mycosphaerella Blight, Anthracnose	Make the first application at the first sign of disease. Apply the high rate only under conditions of high disease pressures. A second application 14 days later may be needed if conditions persist. Good spray coverage and canopy penetration are important for best results.	310 – 620 ml/ac
	Powdery Mildew, White Mould (suppression only)		310 ml/ac
Oats	Barley Net Blotch, Crown Rust, Septoria Leaf Spot	Apply once between stem elongation and half-head emergence.	210 ml/ac
Rye	Septoria Leaf Spot, Barley Scald, Tan Spot	Apply once between stem elongation and half-head emergence.	210 ml/ac
Triticale	Septoria Leaf Spot, Tan Spot		
Wheat	Septoria Leaf Spot, Tan Spot, Stripe Rust, Wheat Leaf Rust		
Durum Wheat	Septoria Leaf Spot, Tan Spot, Stripe Rust		

Timing based on using the product alone. Please refer to tank-mix partners label for crop timing.



TOPNOTCH™

REGISTERED AND SUPPORTED TANK MIXES:

Herbicides:

- ARROW ALL IN®
- BRAZEN® II
- Broadband®
- LADDER ALL IN®
- LEOPARD®
- Glufosinate
- Post Ultra®
- Sierra® 3.0
- Traxion[™]
- Traxos[®]

Insecticides:

- Coragen®
- Decis®
- SILENCER® 120 EC
- Voliam Xpress[®]
- ZIVATA®

Fungicides:

Quadris®

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

- 1. Fill spray tank $\frac{1}{2} \frac{2}{3}$ full with water.
- 2. With agitator running, add required amount of TOPNOTCH™ and continue agitating while adding remainder of the water.
- Begin application after TOPNOTCH™ is completely dispersed into the mix water, and maintain agitation during spraying operation.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

CROP ROTATIONS:

Do not plant any other crop intended for food, grazing or any component of animal feed or bedding within 105 days of application.

PRE-HARVEST INTERVALS:

- Cereals: 45 days
- · Beans, Field Peas, Lentils, Soybeans: 30 days

GRAZING RESTRICTIONS:

Do not graze pea vines. 30 days for all other crops.

STORAGE:

Do not freeze.

CLICK HERE FOR FULL PRODUCT DETAILS.





Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

ARROW ALL IN®, BRAZEN®, LADDER ALL IN®, LEOPARD®, SILENCER®, and ZIVATA®, are registered trademarks, and TOPNOTCH® is a trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. ® 2025 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS

Good spray coverage and canopy penetration are important to achieve the best results.