

Fungicide Groups:

Group 3 & Group 11

Active Ingredient:

Azoxystrobin 143 g/L and Propiconazole 124 g/L = SC

Application Rates and Acres Treated:

Rate: 210 - 620 ml/ac

Acres Treated: 14 - 40 ac/jug

Packaging:

Case: 2 × 8.6 L jugs

Water Volume:

Ground: 40 L/ac (10 US gal/ac)

Aerial: 20 L/ac (5 US gal/ac)

Rainfastness:

Avoid applying when heavy rainfall is in the forecast.

KEY BENEFITS:

- Multiple modes of action for resistance management
- · Curative and preventative systemic action
- Can use on multiple crops, with exceptional crop safety
- · Registered for both ground and aerial spraying
- Excellent crown rust control in oats

KEY DISEASES CONTROLLED:

- Anthracnose
- Ascochyta blight
- Barley leaf rust
- Crown rust
- Mycosphaerella blight
- Net and spot blotches
- · Powdery mildew

- Scald
- Septoria spot
- Stripe rust
- Tan spot
- Wheat leaf rust
- White mould¹

HOW IT WORKS:

Used as both a curative and preventative fungicide, TOPNOTCH $^{\text{\tiny M}}$ has broad-spectrum, systemic and contact activity.


APPLICATION TIMING AND CROP STAGING:

Crop	Diseases	Application Timing	Rate	
Barley	Barley net blotch, Barley scald, Septoria leaf spot, Stripe rust, Barley leaf rust, Tan spot	Apply once between stem elongation and half-head emergence (Growth stage 29 – 55).	210 ml/ac	
Beans, Field peas, Lentils, Soybeans	Mycosphaerella blight, Anthracnose	Make the first application at the first sign of disease. Apply the high rate only under conditions of high disease pressures. A second application 14	310 – 620 ml/ac	
	Powdery mildew, White mould (suppression only)	days later may be needed if conditions persist. Good spray coverage and canopy penetration are important for best results.	310 ml/ac	
Oats Barley net blotch, Crown rust, Septoria leaf spot		Apply once between stem elongation and half-head emergence (Growth stage 29 – 55).	210 ml/ac	

¹Suppression only

TOPNOTCH™

Crop	Diseases	Application Timing	Rate
Rye	Septoria leaf spot, Barley scald, Tan spot		210 ml/ac
Triticale	Septoria leaf spot, Tan spot	A - I I - I I I I I I - I - I	
Wheat	Septoria leaf spot, Tan spot, Stripe rust, Wheat leaf rust	Apply once between stem elongation and half-head emergence (Growth stage 29 – 55).	
Durum wheat	Septoria leaf spot, Tan spot, Stripe rust		



TOPNOTCH™

LENTILS GROWTH STAGES








GROWTH STAGE	PRE-SEED	EMERGENCE	1	2	3	FLOWERING	PRE-HARVEST
PRODUCT							
TOPNOTCH™ ————							





TOPNOTCHTM

SUPPORTED TANK MIXES:

Herbicides:

- ARROW® 240 EC
- ARROW ALL IN®
- BRAZEN™ II
- Broadband®
- LADDER ALL IN®
- LEOPARD®
- Glufosinate
- Post Ultra®
- Sierra® 3.0
- Traxion™
- Traxos®

Insecticides:

- Coragen[®]
- Decis®
- SILENCER® 120 EC
- Voliam Xpress®

Fungicides:

Quadris®

MIXING INSTRUCTIONS:

- 1. Fill spray tank $\frac{1}{2} \frac{2}{3}$ full with water.
- With agitator running, add required amount of TOPNOTCH™ and continue agitating while adding remainder of the water.
- **3.** Begin application after TOPNOTCH™ is completely dispersed into the mix water, and maintain agitation during spraying operation.

CROP ROTATIONS:

Do not plant any other crop intended for food, grazing or any component of animal feed or bedding within 105 days of application.

PRE-HARVEST INTERVALS:

· Cereals: 45 days

• Field peas, Beans, Soybeans: 30 days

• Lentils: 30 days

GRAZING RESTRICTIONS:

No restrictions.

STORAGE:

Do not freeze.

Quick Tips

Good spray coverage and canopy penetration are important to achieve the best results.



Always read and follow label directions. Toll-free: 1.855.264.6262

Website: ADAMA.COM