



INVOLVE® 50 WDG

ADAMA's Group 2 herbicide for control of broadleaf and grassy weed pre-seed, in-crop, post-harvest and summerfallow applications.



GROUP 2

Active Ingredient:

50% Tribenuron-methyl = WDG

Application Rates and Acres Treated:

Rate: 6 g/ac

Acres Treated: 80 ac/bottle;
800 ac/case

Packaging:

Case: 10 × 480 g bottles/case

Water Volume:

Ground: 22 – 44 L/ac
(5 – 12 US gal/ac)

Aerial: Do not apply.

Rainfastness:

Rain within 6 hours may reduce control

REGISTERED CROPS:

Pre-seed/Post-harvest:

- Canary Seed
- Dry Beans
- Faba Bean
- Field Peas
- Lupin
- Oats
- Soybean
- Spring Barley
- Wheat (spring, durum, winter)

INVOLVE® 50 WDG may also be used as a summerfallow herbicide application.

WEEDS CONTROLLED:

Refer to registered label for complete list of weeds controlled and for tank-mix information.

WHEN TANK MIXED WITH 2,4-D ESTER:

- | | | |
|---------------------------------------|---|-------------------------------|
| • Annual Sunflower | • Lamb's-Quarters | • Stinkweed ¹ |
| • Ball Mustard | • Narrow-Leaved Hawk's Beard ¹ | • Sweet Clover |
| • Canada Thistle (top growth control) | • Prickly Lettuce | • Thyme-Leaved Spurge |
| • Flixweed ¹ | • Redroot Pigweed | • Tumble Mustard |
| • Hare's-Ear Mustard | • Russian Pigweed | • Wild Buckwheat ² |
| • Indian Mustard | • Russian Thistle | • Wild Mustard |
| • Kochia (2–10 leaf) | • Shepard's Purse ¹ | • Wild Radish |
| | | • Wormseed Mustard |

WHEN TANK MIXED WITH 0.5 REL/AC GLYPHOSATE:

Broadleaf control:

- | | | |
|-------------------------------|------------------------------|--|
| • Canada Fleabane | • Lady's Thumb | • Volunteer Canola (including glyphosate-tolerant varieties) |
| • Canada Thistle ² | • Lamb's-Quarters | • Volunteer Flax |
| • Common Ragweed | • Narrow-Leaved Hawk's Beard | • White Cockle ¹ |
| • Cow Cockle | • Redroot Pigweed | • Wild Mustard |
| • Dandelion | • Russian Thistle | • Wild Buckwheat |
| • Flixweed | • Stinkweed | |
| • Hempnettle | | |
| • Kochia | | |

Grass control:

- | | | |
|-----------------|--------------------|-------------|
| • Downy Brome | • Persian Darnel | • Wild Oats |
| • Giant Foxtail | • Volunteer Barley | |
| • Green Foxtail | • Volunteer Wheat | |

¹ Fall rosettes and spring seedlings

² Suppressed

INVOLVE® 50 WDG

HOW IT WORKS:

INVOLVE® 50 WDG inhibits the production of the ALS enzyme, quickly causing plants to stop growing and become discoloured (red, yellow, purple) at the growing point and spreading to the entire plant within 1–3 weeks.

CROP STAGING:

- Pre-seed
- In crop/post-emergent
- Summerfallow

PRE-SEED REGISTERED AND SUPPORTED TANK MIXES:

- 2,4-D Ester
- AIM® EC
- Authority® 480
- Glyphosate

POST-EMERGENT REGISTERED AND SUPPORTED TANK MIXES:

- 2,4-D Ester
- AIM® EC
- Assert® 300 SC
- Authority® 480
- Dicamba
- Everest® 3.0
- Puma® Advance

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

ADJUVANT RATE:

- Agral® 90 @ 0.35% v/v
- Not all tank mixes require an adjuvant, see label for details.

PRE-SEED CROP ROTATIONS:

- 24 hours after application: spring wheat (including durum), winter wheat, spring barley, oats, canary seed or pulse crops (including dry bean, faba bean, field pea, lupin and soybean), alfalfa, red clover or alsike clover, smooth brome grass, meadow brome grass, timothy and creeping red fescue.
- 60 days after application: canola, flax and lentils.
- Post-harvest application in the fall may be seeded in the spring to: spring wheat (including durum), spring barley, oats, field corn, canary seed or pulse crops (including dry bean, faba bean, field pea, lupin and soybean), canola, flax, lentils, alfalfa, red clover or alsike clover, smooth brome grass, meadow brome grass, timothy and creeping red fescue or fields may be summerfallowed.

MIXING INSTRUCTIONS:

1. Fill clean spray tank ½ full of clean water, start agitation.
2. Add the required amount of INVOLVE® 50 WDG and agitate until product is completely dispersed.
3. Add the required amount of the tank-mix partner.
4. Add the required amount of surfactant, then fill tank with remaining water.
5. For repeat tank loads, empty the spray tank completely to avoid INVOLVE® 50 WDG from not dispersing or add to tank as a pre-slurry in 5–10 L of water.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

PRE-HARVEST INTERVAL:

30 Days

STORAGE:

May be stored at any temperature.



[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

INVOLVE® is a registered trademarks of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS:

Degree of control and duration of effect depend on weed sensitivity, weed size, spray coverage and growing conditions. Activity of the herbicide mixture may be delayed by cold, dry conditions after application. Injury to pulse crops may occur on coarse-textured soils, low in organic matter (less than 3%), or in fields with variable soils, gravelly areas, sandy areas or eroded knolls. Avoid planting pulse crops in soils containing more than 50% sand.