



# LADDER ALL IN®

Trusted graminicide with a built-in surfactant for wheat. Spend less time mixing and more time on the field.



## GROUP 1

### Active Ingredients:

Clodinafop-propargyl 80 g/L = EC

### Application Rates & Acres Treated:

Rate: 283-356 ml/ac

Acres Treated: 15-20 ac/jug;  
255-320 ac/drum

### Packaging:

Case: 2 x 5.66 L jugs

Bulk: 90.6 L drums

### Water Volume:

Ground: 20 – 40 L/ac  
(5 – 10 US gal/ac)

Aerial: 12 L/ac (3 US gal/ac)

### Rainfastness:

30 minutes

### REGISTERED CROPS AND STAGING:

- Wheat (spring, durum)<sup>1</sup>

### WEEDS CONTROLLED:

#### 1 – 5 leaf stage

##### on main stem:

- Barnyard Grass<sup>2</sup>
- Foxtail (green, yellow)<sup>3</sup>
- Persian Darnel<sup>4\*</sup>

#### 1 – 6 leaf stage

##### on main stem:

- Voluntary Canary Seed<sup>5</sup>
- Wild Oats<sup>5</sup>

#### 3 – 6 leaf stage

##### on main stem:

- Volunteer Oats (tame)<sup>5</sup>

\*Use higher rates when targeting Persian Darnel.

<sup>1</sup> Prior to emergence of 4th tiller

<sup>2</sup> For optimum control apply before tillering, and while barnyard grass is actively growing

<sup>3</sup> For optimum control apply prior to emergence of the 3rd tiller and while foxtail is actively growing

<sup>4</sup> For optimum control apply before tillering, and while Persian darnel is actively growing

<sup>5</sup> Prior to emergence of 4th tiller

### HOW IT WORKS:

LADDER ALL IN® is absorbed by the leaves and is rapidly translocated to the growing points of leaves and stems. Yellowing in 1–3 weeks. Complete control in 3–5 weeks after application.



[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



# LADDER ALL IN®

## REGISTERED AND SUPPORTED TANK MIXES:

### Herbicides:

- 2,4-D amine
- Ally®
- BADGE®
- Barricade® II
- BROMOTRIL®
- Curtail® M
- Dicamba
- Dichlorprop-DX
- Estaprop® XT
- ESTEEM®
- ESTEEM ALL IN®
- FORCEFIGHTER ALL IN®
- FORCEFIGHTER® M
- Infinity®
- Infinity® FX
- Lontrel™ XC
- MCPA amine
- MCPA Ester 600
- MCPA sodium salt 300
- Mecoprop-P
- OUTSHINE®<sup>1</sup>
- OUTSHINE ALL IN®
- Pixxaro®
- Pulsar®
- Refine® SG
- Retain®SG
- RUSH® 24
- RUSH 24 ALL IN®
- Target®
- THRASHER®
- Travallas®
- Trophy®

### Insecticides:

- Decis®
- SILENCER® 120 EC
- ZIVATA®

### Fungicides:

- BUMPER® 432 EC

<sup>1</sup>Normal weed pressure use with high rate of LADDER ALL IN® low rate only with low populations and early application.

When tank mixing with a broadleaf herbicide, insecticide, or a fungicide, always refer to the label of the broadleaf partner, insecticide partner or fungicide partner prior to use.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

## MIXING INSTRUCTIONS:

1. Clean spray tank and ½ fill with clean water. Start agitation or bypass system.
2. If a broadleaf herbicide, insecticide or fungicide is to be used, add the product FIRST prior to adding LADDER ALL IN® and agitate for 2–3 minutes.
3. Add correct amount of LADDER ALL IN®. Agitate for 3–5 minutes before adding remainder of water and then maintain constant agitation.
4. After any break in spraying operations, agitate thoroughly before spraying again.
5. Use the spray suspension as soon as it is prepared.
6. If an oil film starts to build up in the tank, drain tank and then clean with a detergent.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

## PRE-HARVEST INTERVAL:

60 days

## CROP ROTATION:

No restrictions.

## GRAZING RESTRICTIONS:

Observe a minimum of 3 days before grazing livestock on treated crops.

## STORAGE:

May be stored at any temperature.



Always read and follow label directions.  
Toll-free: 1.855.264.6262 | Website: ADAMA.COM/CANADA

BADGE®, BROMOTRIL®, BUMPER®, ESTEEM®, ESTEEM ALL IN®, FORCEFIGHTER®, FORCEFIGHTER ALL IN®, LADDER ALL IN®, OUTSHINE®, OUTSHINE ALL IN®, RUSH®, RUSH 24 ALL IN®, SILENCER®, THRASHER®, TOPNOTCH® and ZIVATA® are registered trademarks of an ADAMA Group Company. All other products are trademarks of their respective companies.  
© 2025 ADAMA Agricultural Solutions Canada Ltd.

## Quick Tips

- Avoid application when heavy rain is forecasted.
- Use higher application rate when targeting Persian dandelion or in cases of heavy grassy weed infestation.
- LADDER ALL IN® contains an internal adjuvant; do not add an external surfactant.