



ADAMA

THRASHER®

Provides excellent dual modes of action for control of 26 tough broadleaf weeds (including Group 2 resistant kochia) in wheat and barley.



HERBICIDE

GROUP 4 & 6

Active Ingredients:

Bromoxynil Octanoic Ester
225 g/L and 2,4-D 2 EH
Ester 225 g/L = EC

Application Rates and Acres Treated:

Rate: 500 ml/ac

Acres Treated: 20 ac/jug;
240 ac/drum; 2,000 ac/tote

Packaging:

Case: 2 x 10 L jug
Bulk: 120 L drums

Water Volume:

Ground: 20 – 40 L/ac
(5 – 10 US gal/ac)

Aerial: 12 – 16 L/ac
(3 – 4 US gal/ac)

Rainfastness:

1 hour

REGISTERED CROPS:

- Barley
- Wheat (spring, durum)

WEEDS CONTROLLED (4-leaf stage unless otherwise noted):

- American nightshade
- Ball mustard
- Bluebur
- Cocklebur
- Common buckwheat (8-leaf)
- Common groundsel (8-leaf)
- Common ragweed
- Cow cockle
- Flixweed
- Green smartweed
- Kochia (2 inches high or 1–12 leaf)
- Lady's thumb
- Lamb's quarters (8-leaf)
- Night-flowering catchfly
- Pale smartweed
- Redroot pigweed
- Russian thistle (2–12 leaf)
- Shepherd's purse
- Stinkweed (8-leaf)
- Tartary buckwheat (8-leaf)
- Triazine-resistant pigweed
- Velvetleaf (3 inches high)
- Volunteer canola
- Volunteer sunflower
- Wild buckwheat (8-leaf)
- Wild mustard (8-leaf)

HOW IT WORKS:

A combination of systemic and contact activity with weeds yellowing within 2–4 days and exhibiting abnormal growth (twisting and cupping of leaves) in 2–10 days.



THRASHER[®]

CROP STAGING:

4-leaf to early flag leaf.

REGISTERED AND SUPPORTED TANK MIXES:

Wheat (spring, durum) and barley:

- BISON[®] 400 L

Wheat (spring, durum) only:

- Everest[®] 3.0
- LADDER ALL IN[®]

MIXING INSTRUCTIONS:

1. Fill the spray tank ½ full with water.
2. Add the required amount of THRASHER[®] and agitate thoroughly.
3. Fill the tank with remaining water and agitate again before use.
4. If tank mixing with a grassy weed herbicide, read both labels and follow the more stringent directions for tank mixing.

CROP ROTATIONS:

No restrictions the year after application.

PRE-HARVEST INTERVAL:

30 days

GRAZING RESTRICTIONS:

- Do not permit livestock to graze fields within 30 days of application.
- Do not harvest, forage or cut for hay within 30 days of application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.

STORAGE:

- May be stored at any temperature.
- Shake well before using.

QUICK TIPS:

For best results when there is a heavy crop canopy, or when the majority of the weeds are cow cockle, smartweed or pigweed, use higher water volumes. Spray when weeds are in the seedling stage. Apply in good growing conditions. Application must be made before the crop canopy shields the weeds.



ADAMA

Always read and follow label directions.
Toll-free: 1.855.264.6262
Website: ADAMA.COM

BISON[®], LADDER ALL IN[®] and THRASHER[®] are registered trademarks of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies. © 2023 ADAMA Agricultural Solutions Canada Ltd.