



## **Active Ingredients:**

Florasulam 50 g/L = SC and MCPA 2-Ethylhexyl Ester 600 g/L = EC

# Application Rates and Acres Treated:

#### Rate:

Florasulam: 40 ml/ac; MCPA Ester 600: 230 ml/ac

Acres Treated: 40 ac/case

### Packaging:

#### Case:

Florasulam: 1 × 1.6 L jug; MCPA Ester 600: 1 × 9.33 L jug

### Water Volume:

Ground: 40 L/ac (10 US gal/ac)

Aerial: Do not apply.

## Rainfastness:

2 hours

#### **REGISTERED CROPS:**

Barley

 Wheat (spring, durum)

## WEEDS CONTROLLED:

#### 1-4 leaf stage:

Ball Mustard

Burdock

Oats

 Common Chickweed

Cleavers

Cow Cockle

Flixweed

Hempnettle<sup>1</sup>

Lamb's-Quarters

Redroot Pigweed

Russian Pigweed

• Prickly Lettuce

Ragweed

Shepherd's Purse

Smartweed

Stinkweed

• Sunflower (annual)

Volunteer Canola<sup>2</sup>

Wild Buckwheat

Wild Mustard

Stork's Bill<sup>1</sup>

## Suppressed:

Annual Sow Thistle

stie

Canada Thistle<sup>1</sup>
 Dandelion<sup>1,3</sup>

Plantain<sup>4</sup>
Perennial Sow

Thistle

<sup>1</sup> For heavy infestations, add 47 ml/ac of MCPA ester for improved control.

<sup>&</sup>lt;sup>2</sup> Including all herbicide-resistant varieties.

<sup>&</sup>lt;sup>3</sup> Seedlings and overwintered rosettes less than 6 inches.

<sup>&</sup>lt;sup>4</sup> Top growth control only.



## **TOPLINE®**

#### **HOW IT WORKS:**

The Group 2 herbicide inhibits the production of the ALS enzyme in plants. This enzyme is essential for the production of certain amino acids required for plant growth. The Group 4 herbicide disrupts normal plant growth, resulting in epinasty and death of susceptible plants.

#### **CROP STAGING:**

Expanded 3-leaf up to the 6-leaf stage.

#### **REGISTERED AND SUPPORTED TANK MIXES:**

#### Wheat and Barley:

- Assert®
- BRAZEN® II

## Wheat only:

- Everest® 3.0
- LADDER ALL IN®
- Simplicity® GoDRI

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

#### **MIXING INSTRUCTIONS:**

- 1. After filling the spray tank ½ full with water, and with agitation running, add the required amount of Florasulam 50 SC, followed by the required amount of MCPA 2 EH Ester 600.
- 2. Fill tank with remaining water.

NOTE: Do not add a surfactant to this mixture.

#### **CROP ROTATIONS:**

Wheat, barley, oats, canola and field peas may be grown the year following an application.

#### STORAGE:

- · May be stored at any temperature.
- Shake well before use.

#### **PRE-HARVEST INTERVAL:**

60 days

#### **GRAZING RESTRICTIONS:**

- Do not cut for feed or hay or allow lactating dairy animals to graze treated crops or within 7 days of application.
- Withdraw meat animals from treated feed 3 days prior to marketing.





Always read and follow label directions.

Toll-free: 1.855.264.6262 | Website: ADAMA.COM/CANADA

BRAZEN®, LADDER ALL IN® and TOPLINE® are registered trademarks, of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd.

## **QUICK TIPS**

TOPLINE® is well suited to dark brown, black and grey soil zones where Cleavers, Hempnettle, Wild Buckwheat and Volunteer Canola are main concerns.

Strong on weeds susceptible to Group 2's (Chickweed, Hempnettle, Redroot Pigweed)