

# **Active Ingredient:**

Bromoxynil Octanoate Ester 240 g/L = EC

# Application Rates and Acres Treated:

Pre-seed:

Rate: 490 ml/ac

Acres Treated: 20 ac/jug

Post-emergent:

Rate: 490 - 570 ml/ac

Acres Treated: 17 - 20 ac/jug;

200-240 ac/drum

# Packaging:

Case: 2 × 9.7 L jugs Bulk: 116.4 L drums

# Water Volume:

### Pre-seed:

Ground: 20 – 40 L/ac (5 – 10 US gal/ac) Aerial: Do not apply.

#### Post-emergent:

Ground: 20 – 80 L/ac (5 – 20 US gal/ac)

Aerial: 8-16 L/ac (3-5 US gal/ac)

wheat and barley only

# Rainfastness:

30 minutes

# **REGISTERED CROPS:**

Crop	Stage
Barley, Oats, Wheat	Pre-seed burn-off with glyphosate

Crop	Crop Leaf Stage
Alfalfa (seedling)	2-6 trifoliate
Alfalfa (established)	Spring: before the crop begins to shield the weeds
Barley, Oats, Triticale, Wheat (spring, durum)	2 leaf to early flag
Winter wheat	Fall: 2 – 4 leaf Spring: first growth to early flag
Corn (field, sweet) with drop pipes	Beyond 8 leaf
Fall rye	Spring: from first growth to early flag
Flax	2-4 inches in height
Forage millet, Sorghum	4 leaf to 8 inches
Seedling grasses	2–4 leaf (year of establishment only)

# **WEEDS CONTROLLED:**

# Seedling up to 4-leaf stage:

American
 nightshade
 Bluebur
 Cocklebur
 Cow cockle¹
 Green smartweed
 Kochia²
 Lady's thumb

Common ragweed
Pale smartweed

Pigweed<sup>1</sup>
Russian thistle<sup>2</sup>
Stinkweed<sup>1</sup>

Velvetleaf<sup>3</sup>
Wild mustard<sup>1</sup>

# Seedling up to 8-leaf stage:

Common
Common groundsel
Tartary buckwheat
Lamb's quarters
Wild buckwheat

<sup>&</sup>lt;sup>1</sup> In normal conditions, it will be controlled up to 4-leaf stage. Plants beyond this stage are unlikely to be controlled; the higher rate generally gives better results.

<sup>&</sup>lt;sup>2</sup> Spray before plants are 2 inches high.

<sup>&</sup>lt;sup>3</sup> Spray before plants are 3 inches high.



# **BROMOTRIL®**

#### **HOW IT WORKS:**

BROMOTRIL® is a contact herbicide which controls Group 2 and Group 9 resistant biotypes. Leaves will yellow in 2-4 days with complete control in 7-14 days.

#### **REGISTERED AND SUPPORTED POST-EMERGENT TANK MIXES:**

#### Post-emergent Herbicides:

- Spring wheat: 2,4-D Ester, LADDER ALL IN®, MCPA Ester 600, BISON® 400 L, Traxos®
- Winter wheat: 2,4-D Ester, MCPA Ester 600, BISON® 400 L, Traxos®
- Barley: 2,4-D Ester, MCPA Ester 600, BISON® 400 L
- Oats: MCPA Ester 600
- Corn: Accent<sup>™</sup>, Atrazine, Banvel<sup>®</sup> (Dicamba), Ultim<sup>®</sup>
- Fall rye: MCPA Ester 600
- Flax: MCPA Ester 600 or MCPA-K
- Canary seed: MCPA Ester 600
- Seedling grasses: MCPA Ester 600

#### Pre-seed:

Glyphosate

#### Fungicides:

BUMPER® 432 EC

#### **MIXING INSTRUCTIONS:**

- 1. Fill spray tank ½ full with water.
- 2. Add required amount of BROMOTRIL®. Begin agitation.
- 3. If tank mixing, add the recommended amount of any tank-mix partners to the spray tank first, agitate and then add BROMOTRIL® (unless otherwise directed by the BROMOTRIL® and tank-mix partner label).
- 4. Add the remaining amount of water while agitation continues.

#### **CROP ROTATIONS:**

No restrictions.

# **PRE-HARVEST INTERVAL:**

30 days

#### STORAGE:

Do not freeze.

#### **GRAZING RESTRICTIONS:**

- Do not use treated crops for grazing of livestock or green feed until 30 days after application.
- Do not cut treated crops for forage until 30 days after application.

#### **QUICK TIPS:**

Avoid spraying if temperatures are above 25° C. Leaf scorching may occur in corn and flax if applied during or after adverse growing conditions, such as cool and wet or hot (above 27° C) and humid weather. For best results, spray when weeds are in the seedling stage.



Always read and follow label directions. Toll-free: 1.855.264.6262 Website: ADAMA.COM